

Lymphocytoma cutis: Cases linked with Lyme disease

Chodynicka, Bozena; Flisiak, Iwona; Okrasinska, Krystyna; Andrzejewska, Anna; Schwartz, Robert A .
Cutis ; Chatham Vol. 66, Iss. 4, (Oct 2000): 243-6.

[ProQuest document link](#)

FULL TEXT

Lymphocytoma cutis (LCC) is a cutaneous B-cell pseudolymphoma representing a wide variety of disorders that share clinical and histologic features.¹⁻ Although the cause in most cases is unknown, it can be induced by a variety of stimuli, including insect bites, *Borrelia burgdorferi* infection, trauma, vaccination, drug or antigen injections, and tattoos.^{1,-} When the bite is from a tick carrying Lyme disease, the result is sometimes called borrelial lymphocytoma (BL). We describe 2 children whose lesions were probably best classified in this latter category. LCC in endemic regions such as Connecticut, New Jersey, or central Europe should be evaluated as a possible sign of Lyme disease.

Case Reports

Patient 1-A 7-year-old boy from a forested region near Bialystok, Poland, was seen for cutaneous nodules of the periumbilical skin, pubic skin, and scrotum of 6 months duration. They had slowly enlarged and were asymptomatic. There was no history of a tick bite. On examination, 3 bluish-red soft painless well-demarcated nodules were evident, 0.5 to 1.5 cm in diameter, on periumbilical and pubic skin and anterior surface of scrotum (Figure 1). There was no significant lymphadenopathy. Ophthalmologic and neurologic evaluations were within normal limits.

Laboratory evaluations showed a normal complete blood cell count and differential. Indirect immunofluorescence assay showed IgM antibodies against *B burgdorferi* titered to 1:256 and by enzyme-linked immunosorbent assay (ELISA) antibodies. Histologic examination of the skin biopsy specimens showed acanthotic epidermal hyperplasia overlying an impressive superficial and deep infiltrate of small mature lymphocytes, with less numerous neutrophils and eosinophils. The formation of lymphoid follicles without distinct germinal centers was evident. In deeper sections, a perivascular pattern of these cells was seen. Skin appendages were preserved. Neither Warthin-Starry nor Steiner and Steiner staining was used. Erythromycin was given orally 250 mg twice a day for 14 days. The lesions resolved without any trace. Follow-up serologic tests, performed 1 and 3 years later, did not reveal any antibodies against *B burgdorferi*.

Patient 2-A 7-year-old boy from a forested region of western Sweden was seen with red slightly pruritic scrotal nodules 6 months after a tick bite at that site. An attached tick had been removed 5 months prior to onset. A cluster of bright red painless nodules was evident on the anterior surface of the scrotum. (Figure 2) There was no significant lymphadenopathy.

The blood cell count and peripheral smear were normal. ELISA did not show IgM or IgG antibodies against *B burgdorferi*. Histologic examination of the skin biopsy specimens showed a dense superficial and deep infiltrate of small mature lymphocytes, with scattered neutrophils and no eosinophils. The formation of lymphoid follicles was not evident. The infiltrate was more prominent in the upper and mid dermis than in the deep dermis. Few mitotic figures were observed. A proliferation of small blood vessels within the cellular infiltration was seen. In deeper sections, a perivascular pattern of these cells was noted. Skin appendages were preserved. Neither Warthin-Starry

nor Steiner and Steiner staining was used. The tissue was not examined under polarized light for retained tick mouth parts. Penicillin V 750 mg twice daily for 2 weeks was given. The lesions resolved without trace.

Discussion

LCC is an uncommon disorder still also known as pseudolymphoma of Spiegler and Fendt.¹ Each described the entity about a century ago."² The name lymphocytoma cubi was coined in 1921 in a description of childhood scrotal nodules by Kaufmann-- Wolf.² Bafverstedt²¹ classified Spiegler-Fendt disease as a lymphocytoma, which he called lymphadenosis benigna cutis, observing the formation of lymphoreticular tissue in skin. LCC tends to appear as a reddish blue nodule, plaque, or tumor. It is typically without symptoms, although it may be painful to touch and persist for months.

When associated with Lyme disease, LCC is usually solitary or a cluster of erythematous or violaceous nodules or plaques about 1 to 5 cm in diameter on the ear lobes or other head regions, the trunk (especially areolae), or extremities.²²⁻²⁸ A predilection for earlobe, nipple and areola mammae, nose, and scrotum probably reflects the preference of spirochetes for regions of low body temperature. This seems a valid concept with regard to the earlobe, but is less clear for the breasts. In one series of 36 patients, 17 had lesions on the earlobe and 15 had them on the breasts.⁶ The average age of those 17 patients with earlobe LCC was 12 years, whereas there were mainly adults with breast involvement. Another study of 8 children showed 6 with ear nodules.⁷ Benign lymphocytic infiltration (Jessner-Kanof disease), which usually appears as a erythematous facial nodule or plaque, may also be a form of borrelial lymphocytoma,^{2,3} as may the dispersed subcutaneous form of LCC.^{23,25} Because LCC is much more frequent in those who probably acquired a Lyme disease infection in Europe rather than North America, a distinct species or subspecies of the organism has been postulated. A recent study of 13 patients showed that BL is caused not only by the *B burgdorferi* genospecies *B afzelii* but also by another genomic group that is seen in North America as well. ¹⁸

In most patients with BL, BL is the first and only cutaneous sign of Lyme disease. In the minority of cases when BL is observed with erythema chronicum migrans (ECM), BL tends to appear weeks after the onset of ECM, and lasts longer. ^{5,8,29-31} However, BL and ECM may occasionally develop simultaneously, sometimes with nodules of BL within a patch of ECM. BL seems to be more common in children than adults. BL may represent an early localized infection or be a sign of an early stage of disseminated Lyme disease. It may appear at the site of the primary spirochete entry within weeks to months after the tick bite. Rarely, BL may be associated with acrodermatitis chronica atrophicans,^{11,11} a late cutaneous finding in Lyme disease." Systemic findings of Lyme disease may also be evident, including carditis, polyradiculitis, and facial palsy with meningitis.^{5,27,28} It has been suggested that every child with BL should receive an electrocardiogram.¹

The histologic features of LCC should prompt elicitation of a history of preceding erythema chronicum migrans or a tick bite, serum antibody titer to *B burgdorferi*, and histologic examination of skin biopsy specimens to demonstrate spirochetes."³⁵ Both Warthin-Starry and Steiner and Steiner methods can be used. Identification of the organism can also be made by polymerase chain reaction or by culture."³⁶ However, even culture cannot be relied upon, as it was positive in only 38.5% in a recent work." The presence of specific antibodies seems to confirm the diagnosis of Lyme disease in our first patient, despite no history of a tick bite. In the second one, the history of a tick bite, the development of a nodule after 15 weeks and its successful treatment with penicillin suggests this diagnosis too,^{11,33} despite a negative serology.³¹ In any case it is important to attempt to diagnose this complex multisystem disease because efficacious treatment is available, and delayed or inadequate therapy can be associated with unfortunate sequelae.⁴⁰

References

REFERENCES

References

1. Ploysangam T, Breneman DL, Mutasim DE Cutaneous pseudolymphoma. *J Am Acad Dermatol.* 1998;38:877-895.

2. Trevisan G, Cattonar P, Nobile C, et al. Dermatological manifestations of Lyme borreliosis. *Acta Derm Venerol (Ljubljana)*. 1996;5:101-107.
3. Abele DC, Anders KH. The many faces and phases of borreliosis II. *J Am Acad Dermatol*. 1990;23:401-410.
4. Kuflik AS, Schwartz RA: Lymphocytoma cutis. a series of five patients successfully treated with cryosurgery. *J Am Acad Dermatol*. 1992;26:449-452.
5. Pohl-Koppe A, Wilske B, Weiss M, et al. *Borrelia lymphocytoma* in childhood. *Ped Infect Dis J*. 1998;17:423-426.
6. Strle F, Pleterski-Rigler G, Stanek G, et al. Solitary borrelial lymphocytoma: report of 36 cases. *Infection*. 1992;20:201-206.

References

7. Miyamoto T, Iwasaki K, Mihara Y, et al. Lymphocytoma cutis induced by cobalt. *Br J Dermatol*. 1997;137:469-471.
8. Bagley MP, Schwartz RA, Lambert WC. Hyperplastic reaction developing within a tattoo. *Arch Dermatol*. 1987;123:1557-1558.
9. Magro CM, Crowson A. Drugs with antihistaminic proper

References

- ties as a cause of atypical lymphoid hyperplasia. *J Am Acad Dermatol*. 1995;32:419-428.
10. Zemtsov A, Cameron GS, Montalvo-Lugo V Nickel– induced lymphocytoma cutis of the earlobe. *Contact Dermatitis*. 1997;36:266.
 11. Asbrink E, Hovmark A, Olsson I. Lymphadenosis benigna cutis solitaria: borrelial lymphocytoma in Sweden. *Zentralbl Bakteriol [131 Suppl]*. 1989; 18:156-163.
 12. Asbrink E. Cutaneous manifestations of Lyme borreliosis: clinical definitions and differential diagnoses. *Scand J Infect Dis Suppl*. 1991;77:44-50.

References

13. Thyresson N. Historical notes on skin manifestations of Lyme borreliosis. *Stand J Infect Dis Suppl*. 1991;77:9-13.
14. Asbrink E, Hovmark A. Comments on the course and classification of Lyme borreliosis. *Stand J Infect Dis Suppl*. 1991;77:41-43.
15. Komatsu H, Aiba S, Mori S, et al. Lymphocytoma cutis involving the lower lip. *Contact Dermatitis*. 1997;36:167-169.
16. Magro CM, Crowson AN. Drug-induced immune dysregulation as a cause of atypical cutaneous lymphoid infiltrates: a hypothesis. *Human Pathol*. 1996;27:125-132.
17. Berger BW. Current aspects of Lyme disease and other *Borrelia burgdorferi* infections. *Dermatol Clin*. 1997;15:247-255.
18. Picken RN, Strle F, Ruzic-Sabljić E, et al. Molecular subtyping of *Borrelia burgdorferi* sensu lato isolates from five patients with solitary lymphocytoma. *J Invest Dermatol*. 1997;108:92-97.

References

19. Spiegler E. Ober die sogenannte Sarkomatosis cutis. *Arch f Dermatol u Syph (Wien)*. 1894;27:163-174.
20. Fendt H. Beitrage zur kenntniss der sogenannten sarcoiden geschwulste der haut. *Arch f Dermatol u Syph (Wien)*. 1900;53:213-242.
21. Kaufmann-Wolf M. Ober gutartige lymphocytare Neubildungen der Scrotalhaut des Kindes. *Arch f Dermatol u Syph (Berlin)*. 1921;130:425-435.
22. Bafversted B. Ober Lymphadenosis benigna cutis. Eine klinische and patologisch-anatomische studie. *Acta*

Derm Venereol (Stockh) Suppl. 1943;11:1-202.

23. Bafversted B. Lymphadenosis benigna cutis (LABC) its nature, course and prognosis. Acta Derm Venereol (Stockh). 1960;40:10-18.

References

24. Abele DC, Anders KH, Chandler FW. Benign lymphocytic infiltration (Jessner-Kanof): another manifestation of borreliosis? J Am Acad Dermatol. 1989;21:795-797.

25. Asbrink E, Hovmark A. Cutaneous manifestations of Ixodesborne borrelia spirochetosis. Intern J Dermatol. 1989;26:215-223.

26. Gottron H. Lymphadenosis cubs circumscripta im bereich der mamille bei gleichzeitiger acrodermatitis chron: atrophicans der extremitaten. Z Hautr. 1938;59:633.

27. Paschoud JM. Lymphocytom nach zeckenbiss. Dermatologica (Basel). 1954;108:435-437.

28. Hovmark A, Asbrink E, Olsson I. The spirochetal etiology of lymphadenosis benigna cutis solitaria. Acta Denn Venereol (Stockh). 1986;66:474-484.

References

29. Chodynicka B, Flisiak I, Lukaszuk C. Studies on Lyme borreliosis in Poland. Medicine 2000. 1996;7:30-33.

30. Flisiak I, Chodynicka B, Laudanska H, et al. A study of the clinical picture of erythema chronicum migrans. Przegl Dermatol. 1996;83:163-166.

31. Chodynicka B, Flisiak I, Lukaszuk C, et al. Serologic and clinical evaluation follow-up after treatment of erythema chronicum migrans. Przegl Dermatol. 1996;83:173-176.

32. Flisiak I, Schwartz RA, Chodynicka B. Clinical features and specific immunological response against Borrelia afzelii in patients with acrodermatitis chronica atrophicans. J Med. 1999;30:267-278.

References

33. Zanconati F, Cattonar P, Grandi G. Histochemical and immunohistochemical methods for demonstration of spirochetes in skin biopsies. Acta Derm Venerol (Ljubljana). 1994;3:99-104.

34. Hodl S, Soyer HP, Mullegger RR. Dermatopathologic diagnosis of Lyme borreliosis. Acta Dean Venerol (Ljubljana). 1996;5:123-129.

35. Albrecht S, Hofstadter S, Artsob H, et al. Lymphadenosis benigna cubs resulting from Borrelia infection (Borrelia lymphocytoma). J Am Acad Dermatol. 1991;24:621-625.

36. Schutzer SE, Schwartz RA. Diagnosing Lyme disease: often simple, often difficult. Cubs. 1991;47:229-232.

37. Strle F, Maraspin V, Pleterski-Rigler D. Treatment of borrelial lymphocytoma. Infection. 1996;24:80-84.

References

38. Krbkova L, Stanek G. Therapy of Lyme borreliosis in children. Infection. 1996;24:170-173.

39. Esquivias Gomez J, Miranda Romero A, Gonzalez Lopez A, et al. Asociacion de liquen escleroatrofico y linfadenosis benigna cutis: marcador cutaneo de infeccion por Borrelia? Med Cutan Iber Lat Am. 1996;24:302-306.

40. Tugwell P, Dennis DT, Weinstein A, et al. Laboratory evaluation in the diagnosis of Lyme disease. Ann Intern Med. 1997;127:1109-1123.

AuthorAffiliation

Bozena Chodynicka, MD, Bialystok, Poland

Iwona Flisiak, MD, Bialystok, Poland

Krystyna Okrasinska, MD, Trollhattan, Sweden

Anna Andrzejewska, MD, Bialystok, Poland

Robert A. Schwartz, MD, MPH, Newark, New Jersey

AuthorAffiliation

Drs. Chodynicka and Flisiak are from the Dermatology and Venereology Department, Medical University of Bialystok, Bialystok, Poland. Dr. Okrasinska is from the Department of Dermatology, TrollhAttan Hospital, Trollhattan, Sweden.

Dr. Andrzejewska is from the Department of Pathological Anatomy, Medical University of Bialystok, Bialystok, Poland. Dr. Schwartz is from the Department of Dermatology, New Jersey Medical School, Newark, New Jersey. REPRINT REQUESTS to Dermatology and Venereology Department, Medical University of Bialystok, ul. Sw. Rocha 3, 15-879 Bialystok, Poland (Dr. Chodynicka).

DETAILS

MeSH:	Animal, Bites &Stings -- complications, Case Report, Child, Diagnosis, Differential, Enzyme-Linked Immunosorbent Assay, Fluorescent Antibody Technique, Indirect, Human, IgM -- blood, Lyme Disease -- complications, Male, Pseudolymphoma -- diagnosis, Pseudolymphoma -- etiology, Scrotum -- pathology, Skin Diseases -- diagnosis, Skin Diseases -- etiology, Ticks
Substance:	IgM
Publication title:	Cutis; Chatham
Volume:	66
Issue:	4
Pages:	243-6
Number of pages:	4
Publication year:	2000
Publication date:	Oct 2000
Publisher:	Quadrant HealthCom Inc.
Place of publication:	Chatham
Country of publication:	United States, Chatham
Publication subject:	Medical Sciences--Dermatology And Venereology
ISSN:	00114162
CODEN:	CUTIBC
Source type:	Trade Journals
Language of publication:	English

Document type: Journal Article

Accession number: 11109143

ProQuest document ID: 202886529

Document URL: <http://ezproxy.liberty.edu/login?url=https://search.proquest.com/docview/202886529?accountid=12085>

Copyright: Copyright Quadrant HealthCom Inc. Oct 2000

Last updated: 2014-05-18

Database: ProQuest Central

LINKS

[Linking Service](#), [Check for full text via Journal Finder](#)

Database copyright © 2018 ProQuest LLC. All rights reserved.

[Terms and Conditions](#) [Contact ProQuest](#)