



UNIVERSIDAD DE GUAYAQUIL
DEPARTMENT OF CHEMICAL SCIENCES

Ciudadela Universitaria "Dr. Salvador Allende"

Telephone: 2293680, E-mail: fcquimic@ug.edu.ec

Guayaquil, Ecuador

FINAL REPORT

CODE: 19-05

TITLE:

Study of the acute oral toxicity of Babuna, originating from NUTRAMEDIX Laboratories, LLC, Florida, USA

OBJECTIVES:

To study adverse side effects produced by the administration of Babuna on body weight and different body systems

BACKGROUND:

Babuna will be used in humans because of the vital importance of carrying out these first-step tests. They will not only guarantee the quality of the product, but will also establish that there are no adverse side effects in humans who take the product.

As discussed in numerous international works, the study of acute toxicity is indispensable, and guarantees (within the margin of error associated with the technique) that the potential for toxicity from the compounds that will be ingested or that may enter into the system accidentally will be learned.

Describing oral acute toxicity in the international literature is a requirement that must be fulfilled for all products that are to be introduced in the market for the first time. (1, 2, 3, 4, 5)

UNIVERSIDAD DE GUAYAQUIL
DEPARTMENT OF CHEMICAL SCIENCES

Ciudadela Universitaria "Dr. Salvador Allende"

Telephone: 2293680, E-mail: fcquimic@ug.edu.ec

Guayaquil, Ecuador

FINAL REPORT

SCIENTIFIC, TECHNICAL, AND SOCIOECONOMIC BENEFITS:

Demonstration of the innocuousness of this product is important in that the product could produce undesirable reactions in individuals who use it. Demonstrating that it does not produce toxic effects can lead to other tests that will allow it to be registered as a new medicine.

VARIABLES TO MEASURE:

- Toxic effects produced by oral application only of this product
- Weight of the animal's day 1, 7 and 14
- Mortality rates and time of death
- When clinical symptoms appear and disappear
- Anatomic-pathological exams (if required)

PROCEDURES TO FOLLOW:

Acute toxicity via oral introduction was determined using the procedures described in the OECD (Organization for Economic Cooperation and Development, comprised of the 24 most developed nations in the world) TG (Test Guidelines) 423.

CHANGES IN THE STUDY PLAN:

Changes did not take place in protocol proposed to the Unity of Quality Guarantee, whose number is referred to on Page 1.

SAMPLE DATA:

Product Name: Babuna Sleep

Represented by: Ing. José Icaza

Entity that carried out the work: University of Guayaquil, Department of Chemical Sciences

UNIVERSIDAD DE GUAYAQUIL

DEPARTMENT OF CHEMICAL SCIENCES

Ciudadela Universitaria "Dr. Salvador Allende"

Telephone: 2293680, E-mail: fcquimic@ug.edu.ec

Guayaquil, Ecuador

FINAL REPORT

Address: Ciudadela Universitaria "Dr. Salvador Allende"

Represented by: Dr. Walter Herrera Arguello

Form of product presentation: glass bottle containing 30 ml

Storage: The product was stored at room temperature, was protected from light and kept under lock and key

INFORMATION WITH RESPECT TO HANDLING:

No special handling instructions were needed

PRODUCT COMPOSITION:

Chamomile flower extract

Mineral water

Ethanol (20 -25 %)

EXPERIMENTAL PROCEDURE:

INTRODUCTION:

This test was performed with the intention of determining the Acute Toxicity by oral intake of the product to be evaluated, given that this is one of the ways proposed for human intake.

DOSAGE USED IN THE TEST:

Data used indicates that:

Suggested use is 5 to 15 drops once a day 30 minutes before going to bed. Additionally these drops are dissolved in a 120 ml glass of water. With this data in mind, each mouse received 20 ml/kg of body weight, but without dissolving in water.

UNIVERSIDAD DE GUAYAQUIL
DEPARTMENT OF CHEMICAL SCIENCES

Ciudadela Universitaria "Dr. Salvador Allende"

Telephone: 2293680, E-mail: fcquimic@ug.edu.ec

Guayaquil, Ecuador

FINAL REPORT

PRINCIPAL TEST PROCEDURE FOLLOWED:

Those that are described in the norms of the OECD. (5)

METHODS AND TECHNIQUES:

Study Material: Babuna Sleep

Animal Model: The test was carried out with a species of rodent (mouse), with a minimum of 6 animals per test of the same sex. That is, 6 females were used in accordance with recommendations with a mean weight of $\pm 20\%$ (7), and belonging to the Wistar line, and originating from the Chemistry Department of the University of Guayaquil. These mice were appropriate for carrying out the study of acute toxicity via oral intake.

The animals were maintained in climate-controlled and quarantine conditions according to established procedures (8, 9), during a period of at least 5 days.

Access to food and water was "ad libitum." (10, 11)

Animals were distributed randomly among the different groups. (12)

Food was denied the mice 18 hours before exposure to the test substance.

The test lasted 19 days (5 days of acclimation, 14 test days)

METHOD DEVELOPMENT:

Two experimental groups were created. Only females were used, as is in accordance with OECD guidelines 423 of 2001, asserting that the female is more sensitive than the male, making it a more appropriate choice for acute toxicity studies.

UNIVERSIDAD DE GUAYAQUIL
DEPARTMENT OF CHEMICAL SCIENCES

Ciudadela Universitaria "Dr. Salvador Allende"

Telephone: 2293680, E-mail: fcquimic@ug.edu.ec

Guayaquil, Ecuador

FINAL REPORT

The evening before the experiment food was denied the animals with the testing carried out after this fasting. After the fasting all animals were weighed to determine the appropriate dosage.

The substance administered was the study product, in a dosage of 4 ml for each 200 grams of animal weight.

Two to three hours after the administration of the product the animals were allowed access to food again.

After the product's administration observations were conducted and systematically recorded for each individual animal, several times on the first day and at least once a day for the next 13 days.

Given that oral ingestion of the product could cause delayed toxic reactions, the animals were weighed on the first, seventh, and 14th days.

At the end of the experiment, the animals were euthanized in a saturated ether atmosphere. (13)

AUTOPSY OBSERVATIONS:

If any abnormality were detected during the examination of the organs (lungs, heart, kidneys and stomach or other organs that may have shown clinical symptoms during the clinical studies), samples were taken for pathological studies (14).

UNIVERSIDAD DE GUAYAQUIL
DEPARTMENT OF CHEMICAL SCIENCES

Ciudadela Universitaria "Dr. Salvador Allende"

Telephone: 2293680, E-mail: fcquimic@ug.edu.ec

Guayaquil, Ecuador

FINAL REPORT

RESULTS CALCULATIONS:

The weights of the mice at different times were statistically processed to obtain the mean and the standard deviation (2).

DESCRIPTION OF THE DOSAGE, ADMINISTRATION METHOD AND DURATION OF THE EXPERIMENT:

The experiment was conducted following the guidelines of OECD TG 423.

The method of administration was oral, using an intra-gastric cannula.

The experiment lasted 19 days (5 of acclimation and 14 of testing).

It is important to realize that this experiment was carried out using a volume of 20 ml per kilogram of body weight. In comparison, a human would be expected to ingest 15 drops, or approximately 0.45 ml dissolved in 120 ml of water. This means 0.00375 ml per 60 kg of body weight. If we convert to kilograms it would be 6.25×10^{-5} ml per kg of body weight. The mouse therefore receives 307,692 times the expected human dosage.

The experiment had a duration of 19 days (5 of acclimation and 14 of experiment).

ANALYTICAL RESULTS:

Results of the daily observations during the 14-day experimental period are recorded in Table 1.



UNIVERSIDAD DE GUAYAQUIL DEPARTMENT OF CHEMICAL SCIENCES

TABLE 1 - CLINICAL SYMPTOMS

PRODUCT: Babuna Sleep
DOSAGE: 20 ml/kg.
START DATE: 04/24/05.

PRODUCT ORIGIN: NutraMedix, LLC, Florida, USA
SEX: Female
END DATE: 05/08/05

CLINICAL SYMPTOMS	DAYS													
	1	2	3	4	5	6	7	8	9	10	11	12	13	14
EYES	-	-	-	-	-	-	-	-	-	-	-	-	-	-
MUCOUS MEMBRANES	-	-	-	-	-	-	-	-	-	-	-	-	-	-
RESPIRATORY SYS.	-	-	-	-	-	-	-	-	-	-	-	-	-	-
CIRCULATORY SYS.	-	-	-	-	-	-	-	-	-	-	-	-	-	-
AUTONOMO	-	-	-	-	-	-	-	-	-	-	-	-	-	-
CENTRAL NERV. SYS.	-	-	-	-	-	-	-	-	-	-	-	-	-	-
CHANGES IN HAIR	-	-	-	-	-	-	-	-	-	-	-	-	-	-
TREMBLING	-	-	-	-	-	-	-	-	-	-	-	-	-	-
CONVULSIONS	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SALIVATION	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SKIN	-	-	-	-	-	-	-	-	-	-	-	-	-	-
SEDATION	+	-	-	-	-	-	-	-	-	-	-	-	-	-
SOMULENCE	+	-	-	-	-	-	-	-	-	-	-	-	-	-
DEATH	-	-	-	-	-	-	-	-	-	-	-	-	-	-
OTHER	-	-	-	-	-	-	-	-	-	-	-	-	-	-

PLEASE NOTE: THE NUMBER OF ANIMALS WITH THE SYMPTOM IS NOTED ON THE CHART.

TEC. RUNNING THE EXPERIMENT:
PROFESSIONAL RESPONSIBLE: MsC. Gaston Garcia Simon

SIGNATURE:
SIGNATURE:

DATE DISSEMINATED: 05/08/05

UNIVERSIDAD DE GUAYAQUIL
DEPARTMENT OF CHEMICAL SCIENCES

Ciudadela Universitaria "Dr. Salvador Allende"

Telephone: 2293680, E-mail: fcquimic@ug.edu.ec

Guayaquil, Ecuador

FINAL REPORT

Clinical symptoms were noted in the study group, as recorded on Table 1. Sedation and somnolence may be due to the alcoholic content of the preparation.

BODY WEIGHT

Table 2 demonstrates the body weights of the animals on day 1, 7, and 14 of the experiment.

TABLE # 2. BODY WEIGHT (GRAMS) VARIATION AND STANDARD DEVIATION OF THE FEMALE ANIMALS IN THE EXPERIMENT OF ACUTE TOXICITY VIA ORAL INGESTION OF CUMANDA.			
GROUP	TIME (Days)		
	0	7	14
FEMALES I	136.6 ± 9.8	153 ± 21.3	169.6 ± 20.5
FEMALES II	144.0 ± 10.39	183.6 ± 4.04	210.0 ± 10.0

As can be seen from Table 2, the females treated with Babuna Sleep showed weight gain between the weighing sessions.

The following bar graphs were created with the values from Table 1.


UNIVERSIDAD DE GUAYAQUIL
DEPARTMENT OF CHEMICAL SCIENCES

Ciudadela Universitaria "Dr. Salvador Allende"

Telephone: 2293680, E-mail: fcquimic@ug.edu.ec

Guayaquil, Ecuador

FINAL REPORT



HISTO-PATHOLOGY:

Samples taken from selected organs showed no affects from under the microscope, thus the pathologist did not take history-pathological samples.

CONCLUSIONS:

- 1- Clinical symptoms were observed in the animals, presumably due to the alcoholic content of the preparation when 4 ml/200 grams of body weight was administered.
- 2- Autopsies revealed no affects to selected organs.
- 3- The product did not affect weight gain of the animals in the study.
- 4- No toxic effects are produced when administering Babuna Sleep in an acute form to the animals.

UNIVERSIDAD DE GUAYAQUIL

DEPARTMENT OF CHEMICAL SCIENCES

Ciudadela Universitaria "Dr. Salvador Allende"

Telephone: 2293680, E-mail: fcquimic@ug.edu.ec

Guayaquil, Ecuador

FINAL REPORT

- 5- Babuna Sleep is a compound whose potential for toxicity is very low based on the fact that its security margin is very high. LD50* cannot be calculated because the maximum dose that can be administered to a mouse is 4 ml per 200 grams of body weight, the dosage used in this experiment. In addition, the dosage used in humans is diluted in 120 ml of water, unlike with the mice that received an undiluted dosage. All this means that, taking into account that the total dissolved solids (1.5 mg/ml) that were administered per 30mg of animal weight, and that humans consume only 6.25×10^{-5} mg, the innocuousness of the product is confirmed.
- 6- The LD50 of Babuna is much higher than the 30 mg/kg administered to the mice, which in turn is much higher than what humans receive, again confirming the safety of the product.


GENERAL CONCLUSIONS:

Babuna Sleep did not produce toxic effects when used in accordance to the guidelines described in OECD TG 423, utilizing a volume of 20 ml/kg, thus the product is considered practically innocuous for human consumption.

PERSONNEL RESPONSIBLE FOR THE STUDY:

DIRECTOR OF THE STUDY:
DR. WALTER HERRERA:

SIGNATURE:



Responsible Professional:
MsC. Gaston Garcia Simon
Date: 05/05/05

SIGNATURE:

A handwritten signature in blue ink, consisting of a stylized 'G' followed by some less distinct characters.

* The dose at which 50 percent of the animals in the experiment suffer mortality.

UNIVERSIDAD DE GUAYAQUIL
DEPARTMENT OF CHEMICAL SCIENCES

Ciudadela Universitaria "Dr. Salvador Allende"

Telephone: 2293680, E-mail: fcquimic@ug.edu.ec

Guayaquil, Ecuador

FINAL REPORT

BIBLIOGRAPHY:

1. Hukeri VI, Kalyani GA, Kakarni (1988). Hypoglycemic activity of flavonoids of Phyllanthus fraternus in rat Fitoterapia 68-70.
2. Copra RN., Nayar SI, Copra LC (1956). Glossary of Indian medicinal plants. Editorial CSIR, Nueva Delhi pp 191.
3. Roig JT (1974) Plantas medicinales aromáticas o venenosas de Cuba. Ciencia y Técnica, la Habana, pp 404, 569, 808.
4. Hayes W., Principles and Methods of Toxicology. Principles and Methods for Acute Toxicity and Eye Irritancy Ed. Raven Press, Ltd. N.Y., 169-220, 1989
- 4A. Ibid, Statistic for toxicologist. 435-483, 1984
5. Organización para la Cooperación Económica y el Desarrollo. (OECD) Guidelines for testing of chemical. Paris. 423, 2001.
6. García G. Los estudios toxicológicos de primera barrera y la toxicología Alternativa. Tesis para optar por el grado académico de Maestro en Ciencias. La Habana, junio del 2000.
7. Procedure. Peso Corporal de las ratas.
8. Procedure. Guía para el cuidado de los animales de Laboratorio.
9. Procedure. Cuarentena .
10. Procedure. Suministro de Agua manual de rutina.
11. Procedure. Asignación aleatoria de las especies roedoras.
12. Procedure. Eutanasia.
13. Procedure. Técnicas de Anatomía Patológica.