

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/10594142>

# Anti- Herpes Simplex Virus Activity of Bidens pilosa and Houttuynia cordata

**Article** in *The American Journal of Chinese Medicine* · January 2003

DOI: 10.1142/S0192415X03001090 · Source: PubMed

CITATIONS

68

READS

104

5 authors, including:



Lien-Chai Chiang

Kaohsiung Medical University

98 PUBLICATIONS 3,756 CITATIONS

[SEE PROFILE](#)



Jung San Chang

Kaohsiung Medical University

39 PUBLICATIONS 781 CITATIONS

[SEE PROFILE](#)



Chun-Ching Lin

Kaohsiung Medical University

202 PUBLICATIONS 7,926 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



I will be retired after three months [View project](#)



Virally-safe and effective plasma and platelet therapies for developing countries [View project](#)

## ***Anti-Herpes Simplex Virus Activity of Bidens pilosa and Houttuynia cordata***

Lien-Chai Chiang,\* Jung-San Chang,† Chi-Chain Chen,†  
Lean-Teik Ng§ and Chun-Ching Lin‡

\*Department of Microbiology, College of Medicine, †Graduate Institute of Medicine, College of Medicine  
and ‡Graduate Institute of Natural Products, College of Pharmacy  
Kaohsiung Medical University, Kaohsiung 807, Taiwan

§Department of Food Sanitation, Tajen Institute of Technology, Ping-Tung 907, Taiwan

**Abstract:** The present study evaluated the antiviral activity of *Bidens pilosa* L. var. minor (Blume) Sherff and *Houttuynia cordata* Thunb., using cytotoxicity test with XTT-based colorimetric assay. BCC-1/KMC cells were infected with herpes simplex virus (HSV) and then were cultured with hot water extract of *B. pilosa* (HWBP) or *H. cordata* (HWHC). Results showed that HWBP significantly inhibited the replication of HSV at a concentration of 100 µg/ml (11.9% for HSV-1,  $p < 0.01$ ; 19.2% for HSV-2,  $p < 0.005$ ), whereas HWHC had the same effect at a concentration of 250 µg/ml (10.2% for HSV-1,  $p < 0.05$ ; 32.9% for HSV-2,  $p < 0.005$ ). The ED<sub>50</sub> of HSV type 1 (HSV-1) and HSV type 2 (HSV-2) for HWBP was 655.4 µg/ml and 960 µg/ml respectively, for HWHC it was 822.4 µg/ml and 362.5 µg/ml respectively. Both drugs had selective indexes above 1.04. *H. cordata* had better effect against HSV-2 than HSV-1, and had a low ED<sub>50</sub> against HSV-2. We suggest that *H. cordata* might be a useful medicinal plant against infection of HSV-2.

**Keywords:** *Bidens pilosa*; *Houttuynia cordata*; Herpes Simplex Virus.

### **Introduction**

During the past 50 years, a great deal of interest had been initiated in screening therapeutic agents from plants. *Bidens pilosa* L. was widely used as a traditional remedy for treating influenza, diabetes and gastroenteritis worldwide. *Houttuynia cordata* Thunb. was used in traditional Chinese medicine for treating infectious disease, refractory hemoptysis, malignant pleural effusion, nephrotic syndrome (Zheng *et al.*, 1998). Studies had shown that *B. pilosa* possessed antihyperglycemic (Ubillas *et al.*, 2000), antiulcerogenic (Alvarez *et al.*, 1999),

Correspondence to: Professor Chun-Ching Lin, College of Pharmacy, Kaohsiung Medical University, 100 Shih-Chuan 1st Road, Kaohsiung 807, Taiwan. Tel: (+886) 07-312-1101 (ext. 2122), Fax: (+886) 07-313-5215, E-mail: aalin@ms24.hinet.net

immunosuppressive (Pereira *et al.*, 1999), anti-inflammatory (Pereira *et al.*, 1999; Jager *et al.*, 1996; Chih *et al.*, 1995; Geissberger and Sequin, 1991), vasodilative (Dimo *et al.*, 1998), antimalarial (Brandao *et al.*, 1997), antibacterial (Geissberger and Sequin, 1991; Rabe and van Staden, 1997), and hepatoprotective activities (Chin *et al.*, 1996). However, there was no previous report of antiviral activity. Also co-carcinogenesis (Mirvish *et al.*, 1979; Mirvish *et al.*, 1985) and photocytotoxicity (Arnason *et al.*, 1980; Wat *et al.*, 1979) were reported with different species of *B. pilosa*.

*H. cordata* was reported to possess antimicrobial (Zheng *et al.*, 1998), antiviral (Zheng *et al.*, 1998; Hayashi *et al.*, 1995), immunostimulatory (Zheng *et al.*, 1998), diuretic (Zheng *et al.*, 1998), anticancer (Zheng *et al.*, 1998), sedative (Zheng *et al.*, 1998), anti-inflammatory (Zheng *et al.*, 1998) and antitussive effects (Zheng *et al.*, 1998). Although *H. cordata* was found to have activity against *herpes simplex* virus type 1 (HSV-1) (Hayashi *et al.*, 1995), no effect against *herpes simplex* virus type 2 (HSV-2) was reported (Zheng *et al.*, 1998). In searching natural crude drugs for potential anti-HSV activity, hot water extracts of whole plant of *B. pilosa* (HWBP) and *H. cordata* (HWHC) were tested.

## Materials and Methods

### *Preparation of Tested Drugs*

Hot water extracts of *B. pilosa* (HWBP) and *H. cordata* (HWHC) were prepared as follows: 100 g of dried whole plant of *B. pilosa* or *H. cordata* were added with 1000 ml reverse-osmotic water in a flask. They were boiled for 1 hour and then supernatant were collected. Three repetitions of these procedures were done. All these supernatants were mixed for filtration by a filter and for dryness in a vacuum. Dissolution and dilution of the powder with di-distilled water to concentrations of 100 µg/ml, 250 µg/ml and 500 µg/ml were done before experiments (Chang and Yeung, 1988). Acyclovir and dimethyl sulfoxide (DMSO) were purchased from Sigma Chemical Co. (USA). Acyclovir was suspended in DMSO and diluted with di-distilled water to concentrations of 0.1 µg/ml, 0.5 µg/ml and 1 µg/ml before use.

### *Cells*

BCC-1/KMC cell line (Chiang *et al.*, 1994) was used for viral culture and antiviral screening. Cells were cultured with RPMI 1640 medium supplemented with 10% fetal bovine serum and 1% antibiotics (10000 u/ml penicillin, 10 mg/ml streptomycin, 25 µg/ml amphotericin B; Sigma, USA) at 37°C in a humidified atmosphere containing 5% CO<sub>2</sub>.

### *Cytoprotective Activity Against HSV with XTT-based Colorimetric Assay*

BCC-1/KMC cells, treated by trypsin, were seeded into 96-well plates with a concentration of  $1 \times 10^5$  cells/ml and a volume of 70 µl per well. After incubation for 6 hours at 37°C with 5% CO<sub>2</sub>, 20 µl HSV-1 containing 25 TCID<sub>50</sub> or 20 µl HSV-2 containing 20 TCID<sub>50</sub> was

added. The mixtures were incubated for another 2 hours, followed by adding different concentrations of hot water extracts to the triplicate culture wells. DMSO was used as a negative control while acyclovir solution was used as a positive control. After incubation at 37°C with 5% CO<sub>2</sub> for three days, a mixture of 0.1 ml PMS (electron-coupling reagent) and 5 ml XTT (Sigma, USA) was added to each well in a volume of 50 µl. The trays were re-incubated for another 2 hours to allow XTT formazan production. The content of each plate was mixed and the optical density was determined with the ELISA reader (Multiskan EX, Labsystems) at 450 nm as test wavelength and 690 nm as reference wavelength.

Cytoprotection rate was calculated as  $(OD_{tv}-OD_{cv})/(OD_{cd}-OD_{cv}) \times 100\%$ . OD<sub>tv</sub> represents the absorbance of the test compounds with virus-infected cells. OD<sub>cv</sub> represents the absorbance of viral control. OD<sub>cd</sub> was the absorbance of the cell control only. The antiviral dose of 50% effectiveness (ED<sub>50</sub>) was defined as the concentration which achieved 50% cytoprotection against viral infection. The concentration of 50% cytotoxicity (CC<sub>50</sub>) of normal human lymphocytes ( $3 \times 10^6$  Cells/ml) was assayed and calculated by the above methods. The selective index was determined by the ratio of CC<sub>50</sub> to ED<sub>50</sub>.

### Statistical Analysis

Means and standard errors were calculated with the Excel software for Windows. Chi-square test with Yate's correction was used to calculate p value between control and samples by the SPSS Base 8.0 software for Windows. Difference with  $p < 0.05$  was considered as statistically significant.

### Results

Both HWBP and HWHC showed a significant inhibitory activity against infection of HSV (Tables 1 and 2). The inhibitory effects were dose-dependent (Figs. 1 and 2). At a concentration of 100 µg/ml HWBP, 11.9% ( $p < 0.01$ ) of cells were protected from infection of HSV-1 and 19.2% ( $p < 0.005$ ) from infection of HSV-2 (Fig. 1 and Table 1). In the case of HWHC, at a concentration of 250 µg/ml, 10.2% ( $p < 0.05$ ) of cells were protected from infection of HSV-1 and 32.9% ( $p < 0.005$ ) from infection of HSV-2 (Fig. 2 and Table 2). The inhibitory effect of HWHC against HSV-2 was comparable to that of acyclovir (0.5 µg/ml) and was better than that against HSV-1.

**Table 1. Cytoprotection Rates of *B. pilosa* Against HSV Infection**

	HWBP (µg/ml)				Acyclovir 0.5
	Solvent Control	100	250	500	
HSV-1 (%)	1.52	11.92*	23.82†	39.02†	45.07
HSV-2 (%)	0.26	19.21†	26.72†	33.07†	33.29

Data represented an average of three tests.

\* $p < 0.01$ ; † $p < 0.005$  (chi-square test with Yate's correction).

**Table 2. Cytoprotection Rates of *H. cordata* Against HSV Infection**

	HWHC ( $\mu\text{g/ml}$ )				Acyclovir 0.5
	Solvent Control	100	250	500	
HSV-1 (%)	1.52	4.71	10.23*	27.6 <sup>†</sup>	45.07
HSV-2 (%)	0.26	5.37	32.86 <sup>†</sup>	70.96 <sup>†</sup>	33.29

Data represented an average of three tests.

\* $p < 0.05$ ; <sup>†</sup> $p < 0.005$  (chi-square test with Yate's correction).




Figure 1. Cytoprotection rate of *B. pilosa* against HSV infection. Solvent was used as negative control. Acyclovir (0.5  $\mu\text{g/ml}$ ) was used as positive control (\*\* $p < 0.01$ ; \*\*\* $p < 0.005$ ).




Figure 2. Cytoprotection rate of *H. cordata* against HSV infection. Solvent was used as negative control. Acyclovir (0.5  $\mu\text{g/ml}$ ) was used as positive control (\* $p < 0.05$ ; \*\*\* $p < 0.005$ ).

HWBP had a better effect than that of HWHC in protecting cells from infection by HSV-1 ( $p < 0.025$ , Fig. 3 and Table 3) at concentration of 250  $\mu\text{g/ml}$ . The same result was also found in preventing infection from HSV-2 ( $p < 0.01$ , Fig. 4 and Table 3) at a lower concentration (100  $\mu\text{g/ml}$ ). However, HWBP exhibited less effect than that of HWHC in protecting cells from infection of HSV-2 ( $p < 0.005$ , Fig. 4 and Table 3) at a higher concentration (500  $\mu\text{g/ml}$ ). The cytoprotection rate of HWHC against infection of HSV-2 rapidly increased as the concentration rose (5.4% at a concentration of 100  $\mu\text{g/ml}$  to 71.0% at a concentration of 500  $\mu\text{g/ml}$ , Table 2). The  $\text{ED}_{50}$  of HWHC was 362.5  $\mu\text{g/ml}$  (Table 4).




Figure 3. Comparison of cytoprotection rate between *B. pilosa* and *H. cordata* against HSV-1 infection. *B. pilosa* had better effect than that of *H. cordata* at a concentration of 250  $\mu\text{g/ml}$  ( $*p < 0.025$ ). There was no difference between other concentrations ( $p > 0.05$ ).

**Table 3. Cytoprotection Rates of *B. pilosa* and *H. cordata* Against HSV Infection**

		HWBP and HWHC ( $\mu\text{g/ml}$ )		
		100	250	500
HSV-1 (%)	<i>B. pilosa</i>	11.92	23.82*	39.02
	<i>H. cordata</i>	4.71	10.23	27.6
HSV-2 (%)	<i>B. pilosa</i>	19.21 <sup>†</sup>	26.72	33.07
	<i>H. cordata</i>	5.37	32.86	70.96 <sup>‡</sup>

Data represented an average of three tests.

\* $p < 0.025$ ; <sup>†</sup> $p < 0.01$ ; <sup>‡</sup> $p < 0.005$  (chi-square test with Yate's correction).




Figure 4. Comparison of cytoprotection rate between *B. pilosa* and *H. cordata* against HSV-2 infection. The effect of *H. cordata* had rapidly increased when drug concentration rose (\*\* $p < 0.01$ ; \*\*\* $p < 0.005$ ).

Table 4. Antiviral Activity of *B. pilosa* and *H. cordata*

	<i>B. pilosa</i>		<i>H. cordata</i>		Acyclovir	
	ED <sub>50</sub> (µg/ml)	SI	ED <sub>50</sub> (µg/ml)	SI	ED <sub>50</sub> (µg/ml)	SI
<b>HSV-1</b>	655.44	>1.53	822.39	>1.22	0.61	17.26
<b>HSV-2</b>	960	>1.04	362.46	>2.76	0.81	12.9

ED<sub>50</sub>: Concentration of 50% cytoprotection.

SI: Selective index = CC<sub>50</sub>/ED<sub>50</sub>.

CC<sub>50</sub>: Concentration of 50% cytotoxicity. CC<sub>50</sub> of *B. pilosa* and *H. cordata* were > 1000 µg/ml. CC<sub>50</sub> of acyclovir was 10.44 µg/ml.

## Discussion

Much effort had been made to search for new antiviral therapeutic agents during the past 50 years. This had led to numerous studies of medicinal plants for clinical applications (Carter and Livingston, 1976; Marsoni and Wittes, 1984). *B. pilosa* and *H. cordata* had been widely used in the traditional medicine of different ethnic groups in the world. *H. cordata* was reported to have activity against HSV-1 (Hayashi *et al.*, 1995), but not HSV-2 (Zheng *et al.*, 1998). In this study, we demonstrated that both HWBP and HWHC possessed antiviral activity and the activity was dose-dependent (Figs. 1 and 2). In contrast to the findings of Zheng *et al.* (1998), *H. cordata* did have activity against HSV-1 and HSV-2. The activity against HSV-2 was better than that against HSV-1 (Fig. 2 and Table 2).

The ED<sub>50</sub> of *B. pilosa* and that of *H. cordata* were much higher than ED<sub>50</sub> of acyclovir (Table 4). We still believed that both plants had value for clinical application. Several reasons could explain the higher ED<sub>50</sub> of *B. pilosa* and *H. cordata*. The first, concentration of active

compounds of the plant materials might be too low or active components might be reduced during extraction. The second, they might become more active after transformation *in vivo* (Marsoni and Wittes, 1984; Double, 1992). Therefore, purification of the effective components from these crude drugs might be necessary.

High CC<sub>50</sub> (>1000 µg/ml) of HWBP and HWHC indicated a good tolerability by human cells. It will be possible to use these plants against infection of HSV in the future (Table 4).

According to our results, it is worthy to study *B. pilosa* and *H. cordata* further. The fractionation, separation of active components and clarification of their mechanism of action are currently under investigation.

## Conclusion

Hot water extracts of *B. pilosa* and *H. cordata* showed activity against HSV. The present study has demonstrated the potential clinical implication of *B. pilosa* and *H. cordata*. Hence, they are worthy of further investigation.

## References

- Alvarez, A., F. Pomar, M.A. Sevilla and M.J. Montero. Gastric antisecretory and antiulcer activities of an ethanolic extract of *Bidens pilosa* L. var. *radiata* Schult. Bip. *J. Ethnopharmacol.* 67: 333–340, 1999.
- Arnason, T., C.K. Wat, K. Downum, E. Yamamoto, E. Graham and G.H. Towers. Photosensitization of *Escherichia coli* and *Saccharomyces cerevisiae* by phenylheptatriene from *Bidens pilosa*. *Can. J. Microbiol.* 26: 698–705, 1980.
- Brandão, M.G.L., A.U. Krettli, L.S.R. Soares, C.G.C. Nery and H.C. Marinuzzi. Antimalarial activity of extracts and fractions from *Bidens* and other *Bidens* species (Asteraceae) correlated with the presence of acetylene and flavonoid compounds. *J. Ethnopharmacol.* 57: 131–138, 1997.
- Carter, S.K. and R.B. Livingston. Plant products in cancer chemotherapy. *Cancer Treat. Rep.* 60: 1141–1156, 1976.
- Chang, R.S. and H.W. Yeung. Inhibition of growth of human immuno-deficiency virus *in vitro* by crude extracts of Chinese medical herbs. *Antiviral Res.* 9: 163–175, 1988.
- Chiang, L.C, W. Chiang, H.S. Yu, H.M. Sheu and H.Y. Chen. Establishment and characterization of continuous human basal cell carcinoma cell line from facial skin. (I) Cytological behavior of early passages. *Kaohsiung J. Med. Sci.* 4: 170–176, 1994.
- Chih, H.W., C.C. Lin and K.S. Tang. Anti-inflammatory activity of Taiwan folk medicine “ham-hong-chho” in rats. *Am. J. Chin. Med.* 23: 273–278, 1995.
- Chin, H.W., C.C. Lin and K.S. Tang. Hepatoprotective effects of Taiwan folk medicine ham-hong-chho in rats. *Am. J. Chin. Med.* 24: 231–240, 1996.
- Dimo, T., S. Rakotonirina, R. Kamgang, P.V. Tan, A. Kamanyi and M. Bopelet. Effects of leaf aqueous extract of *Bidens pilosa* (Asteraceae) KCl- and norepinephrine-induced contractions of rat aorta. *J. Ethnopharmacol.* 60: 179–182, 1998.
- Double, J.A. Selectivity and potency: are we doing the right things to find anti-cancer agents with these properties? *Br. J. Cancer* 65: 143–144, 1992.



- Geissberger, P. and U. Sequin. Constituents of *Bidens pilosa* L.: do the components found should explain the use of this plant in traditional medicine? *Acta Trop.* 48: 251–261, 1991.
- Hayashi, K., M. Kamiya and T. Hayashi. Virucidal effects of the steam distillate from *Houttuynia cordata* and its components on HSV-I, influenza virus and HIV. *Planta Med.* 61: 237–241, 1995.
- Jager, A.K., A. Hutchings and J. van Staden. Screening of Zulu medicinal plants for prostaglandin-synthesis inhibitors. *J. Ethnopharmacol.* 52: 95–100, 1996.
- Marsoni, S. and R. Wittes. Clinical development of anticancer agents — a National Cancer Institute perspective. *Cancer Treat. Rep.* 68: 77–84, 1984.
- Mirvish, S.S., E.F. Rose and D.M. Sutherland. Studies on the esophagus. II. Enhancement of [<sup>3</sup>H] thymidine incorporation in the rat esophagus by *Bidens pilosa* (a plant in South Africa) and by croton oil. *Cancer Lett.* 6: 159–165, 1979.
- Mirvish, S.S., S. Salmasi, T.A. Lawson, P. Pour and D. Sutherland. Test of catechol, tannic acid, *Bidens pilosa*, croton oil, and phorbol for cocarcinogenesis of esophageal tumors induced by methyl-n-amyl nitrosamine. *J. Natl. Cancer Inst.* 74: 1283–1290, 1985.
- Pereira, R.L.C., T. Ibrahim, L. Lucchetti, A.J.R. da Silva and V.L.G. de Moraes. Immunosuppressive and anti-inflammatory effects of methanolic extract and the polyacetylene isolated from *Bidens pilosa* L. *Immuno-pharmacology* 43: 31–37, 1999.
- Rabe, T. and J. van Staden. Antibacterial activity of South African plants used for medicinal purposes. *J. Ethnopharmacol.* 56: 81–87, 1997.
- Ubillas, R.P., C.D. Mendez, S.D. Jolad, J. Luo, S.R. King, T.J. Carlson and D.M. Fort. Antihyperglycemic acetylenic glucosides from *Bidens pilosa*. *Planta Med.* 66: 82–83, 2000.
- Wat, C.K., R.K. Biswas, E.A. Graham, L. Bohm, G.H. Towers and E.R. Waygood. Ultraviolet-mediated cytotoxic activity of phenylheptatriene from *Bidens pilosa* L. *J. Nat. Prod.* 42: 103–111, 1979.
- Zheng, H.Z., Z.H. Dong and J. She (eds.). *Houttuynia cordata*. In: *Modern Study of Traditional Chinese Medicine, Vol. 3*. Xue Yuan Press, Beijing, 1998, pp. 2983–3003.