



## REVIEW

WILEY

# Biological, nutritional, and therapeutic significance of *Moringa oleifera* Lam

Ashok K. Dhakad<sup>1</sup> | Mohsin Ikram<sup>2</sup> | Shivani Sharma<sup>3</sup> | Salman Khan<sup>2</sup> |  
Vijay V. Pandey<sup>4</sup> | Avtar Singh<sup>1</sup>

<sup>1</sup>Department of Forestry and Natural Resources, Punjab Agricultural University, Ludhiana, India

<sup>2</sup>Forest Entomology Division, Forest Research Institute, Dehradun, India

<sup>3</sup>Department of Microbiology, Punjab Agricultural University, Ludhiana, India

<sup>4</sup>Forest Pathology Division, Forest Research Institute, Dehradun, India

## Correspondence

Dr. Ashok Kumar Dhakad, Department of Forestry and Natural Resources, Punjab Agricultural University, Ludhiana, India.  
Email: ashokdhakad@pau.edu

The genus *Moringa* Adans. comprises 13 species, of which *Moringa oleifera* Lam. native to India and cultivated across the world owing to its drought and frost resistance habit is widely used in traditional phytomedicine and as rich source of essential nutrients. Wide spectrum of phytochemical ingredients among leaf, flower, fruit, seed, seed oil, bark, and root depend on cultivar, season, and locality. The scientific studies provide insights on the use of *M. oleifera* with different aqueous, hydroalcoholic, alcoholic, and other organic solvent preparations of different parts for therapeutic activities, that is, antibioid, antitumor, antioxidant, anti-inflammatory, cardio-protective, hepato-protective, neuro-protective, tissue-protective, and other biological activities with a high degree of safety. A wide variety of alkaloid and sterol, polyphenols and phenolic acids, fatty acids, flavanoids and flavanol glycosides, glucosinolate and isothiocyanate, terpene, anthocyanins etc. are believed to be responsible for the pragmatic effects. Seeds are used with a view of low-cost biosorbent and coagulant agent for the removal of metals and microbial contamination from waste water. Thus, the present review explores the use of *M. oleifera* across disciplines for its prominent bioactive ingredients, nutraceutical, therapeutic uses and deals with agricultural, veterinarian, biosorbent, coagulation, biodiesel, and other industrial properties of this “Miracle Tree.”

## KEYWORDS

*Moringa oleifera*, nutritional properties, phytochemistry, therapeutic activities, water purification

## 1 | INTRODUCTION—RATIONALE AND OBJECTIVES

Approximately 80% of the total population depends only on plants for their well-being and healing (Ekor, 2014), and 25% of the integrated medications are made from medicinal and aromatic plants (Pan et al., 2013). The last few decades have seen chemical revolution, and most of things such as food, drugs, agriculture, and environment have been filled with chemicals. As the new discoveries are being made day by day and increasing dependency on chemical supplements and junk foods, the adverse effects of chemicals are being exposed and development of resistance against orthodox medicines;

thus, human attention has shifted towards the use of herbal plants, natural products, and organic farming (Igado & Olopade, 2016), especially because they are generally considered safe and sound. The traditional knowledge in Ayurveda is still in practice, which has made a good amalgam of traditional and scientific knowledge. Recently, *M. oleifera* tree has been used as a remarkable indigenous source of protein, crude fibers, minerals, and nutrients suitable for utilization in developing nations where undernourishment is a major concern (Verma & Nigam, 2014). *Moringa* tree has been used to combat malnutrition, especially among newborns and nursing mothers. Different organizations have promoted *Moringa* as “natural nutrition for the tropics” (Fahey, 2005).

*M. oleifera* (MO) belongs to monogeneric family *Moringaceae* (Marrufo et al., 2013) and known as sajjana, drumstick, and Horseradish tree. *Moringa* derives from a Tamil word, *murungai*, meaning “twisted pod,” alluding to the young fruit (Olsen, 2010). It is native to dry tropical forest of north-west India and foothills of the Himalaya (Lalas, Gortzi, Athanasiadis, Tsaknis, & Chinou, 2012). *M. oleifera*, *M. concanensis*, and *M. hildebrandtii* are scattered in the south Asia and Arabia; *M. drouhardii* in Madagascar; *M. stenopetala*, *M. arborea*, *M. borziana*, *M. longituba*, *M. pygmaea*, *M. rivae*, and *M. ruspoliana* in north-east and north-west Africa; *M. peregrina* in Red Sea, Arabia, Horn of Africa; *M. ovalifolia* in Namibia and South west Angola (Mark & Sylvain, 2006). Out of 13 species (Sengupta & Gupta, 1970), *M. oleifera* [syn. *M. pterygosperma* Gaertn. Nom. illeg, *Guilandina moringa* L. and *Hyperanthera moringa* (L.) Vahl] is extensively utilized and well-known species in most parts of Asia and Africa. Recorded evidences reveal that ancient rulers utilized *Moringa* leaves and natural product in their diet routine to keep up mental health and smooth skin (Mahmood, Mugall, & Haq, 2010). Each and all parts of *Moringa* are valuable for medicinal, useful food ingredients, nutraceuticals, water sanitization, and biofuel production (Saini, Sivanesan, & Keum, 2016). The flowers, unripened fruits, and leaves are utilized for cooking purposes worldwide (Stevens, Baiyeri, & Akininagbe, 2013). Specifically, they are consumed for human and animal nourishment and treatment of many disorders in Asian and African countries (Mbikay, 2012).

Despite its extensive uses in food and medicines, there are very few studies on clinical trials to demonstrate the efficacy for treating malnutrition and undeniable therapeutic action of *Moringa* leaf and other products in human beings. Owing medicinal significance, the seed oil of *Moringa* is widely used for medications due to high content of oleic acid, that is, 72% (Anwar & Rashid, 2007) in African and Asian countries. In spite of its (MO) self evident importance in practical utility and consisting food ingredients and natural remedial properties, there is an urgent need to screen the bioactive ingredients of *Moringa* for herbal therapy along with other ayurvedic treatments and geographical variations for nutrients worldwide. Thus, after reviewing the literature on phytochemical, nutritional, therapeutic properties, commercial use as biosorbent, and coagulant properties of *M. oleifera*, it is worthwhile to compile a comprehensive and updated review.

## 2 | METHODOLOGY

### 2.1 | Search strategem

The informations quoted in this review on *M. oleifera* Lam. along with its chemistry and plant part uses were searched and obtained using databases such as university online databases subscribed, Consortium for e-Resources in Agriculture, Krishikosh (An Institutional Repository of Indian National Agricultural Research System), Open Access e-Resources like PubMed, Science Direct, Scopus, MedLine, Research Gate, and Google Scholar. All were searched with the keyword “*Moringa oleifera* Lam.” along with “phytochemical constituents,” “nutritive value,” “food fortification,” “therapeutic activities,”

“agriculture and animal husbandry uses,” “coagulant/biosorbent/bio-diesel uses,” and “industrial uses.” No language restrictions were imposed. Research papers were assessed for the information about the extracts or fractions and isolated compounds of the plant or its parts, results of that particular study. The search process was also conducted by using a combination of different categories of keywords like “bacteria,” “fungi,” “parasites,” “nematodes,” and medical subject terms namely, “cancer,” “tumor,” “antioxidant,” “anti-inflammation,” “diabetes,” “hypoglycemic,” “hyperlipidemic,” “cholesterol,” and “blood sugar.” To increase the search sensitivity, quotation marks, parentheses, and asterisks were applied to search for the exact terms or expressions, locating a group of search terms and all of the words derived from one keyword, respectively. Finally, the reference lists of selected studies and previous reviews were manually screened to acquire other applicable publications.

### 2.2 | Data extraction

We extracted the following information from the full-text research papers and review articles of appropriate studies. Among the vast evidences, published articles listed in reference section were found in the databases that contain screening reports on the phytochemical and/or therapeutic activities of *M. oleifera* or its crude extracts/fractions or isolated compounds. The observed information in the included literatures has been summarized in the present review.

## 3 | RESULTS AND DISCUSSION

### 3.1 | Phytochemical profiling

Different parts of *M. oleifera* have been used for good source of unique glucosinolates, flavonoids and phenolic acids (Amaglo et al., 2010; Coppin et al., 2013), polyunsaturated unsaturated fatty acids (PUFAs), tocopherols, exceedingly minerals, and folate (Saini et al., 2016). Leaves aqueous extract contained 1.66% isothiocyanates and 3.82% total polyphenols (Waterman et al., 2014). Glucosinolates (glucomoringin) and isothiocyanates are the predominantly in the foliage, flowers, fruits, seeds, and bark of MO (Amaglo et al., 2010). Despite the fact that benzyl glucosinolate (glucotropaeolin) is prominently in roots, the highest noteworthy substance, that is, glucosinolate is reported in the leaves and seeds. The myrosinase enzymatic catabolism of glucosinolates produced isothiocyanates, nitriles, and thiocarbamates (Anwar, Latif, Ashraf, & Gilani, 2007). Among flavonol glycosides, quercetin and isorhamnetin are prevalently found in various plant parts except in seeds and roots. Refined gum exudates contain galactose, glucuronic acid, larabinose, mannose, rhamnose, and xylose (Shah, Jhade, & Chouksey, 2016). All-E-lutein is the main carotenoid in leaves and immature fruits, representing 53.6% and 52.0% of total carotenoids. Among different tissues, the most astounding substance of total carotenoids is reported in leaves (44.30–80.48 mg/100 g FW), followed by immature pods (29.66 mg/100 g FW), and flowers (5.44 mg/100 g FW). Singh et al. (2009) recorded total

phenolic content (105 mg gallic acid equivalents/100 g), flavonoid (31 mg quercetin equivalents/100 g), and ascorbic acid (107 mg/100 g) in leaves. It is an established fact that leaves are a rich source of omega-3 and omega-6 PUFAs, as linolenic acid (49–59%) and linoleic acid (6–13%). Palmitic acid is reported as major unsaturated fat, representing 16–18% of total unsaturated fats in leaves. Pods and inflorescences are described by a higher substance of total monounsaturated unsaturated fatty acids with the range of 16–30% and are low in PUFAs (34–47%), contrasted with the leaves (Saini, Shetty, & Giridhar, 2014). Interestingly, the seeds and seed oil contain a high amount of oleic (70–80%), palmitoleic (6–10%), stearic (4–10%), arachidic (2–4%), and low amount of linoleic and linolenic acids (Amaglo et al., 2010). MO leaves had high amount of total phenolic compounds ( $56.10 \pm 0.90$  mg gallic acid  $\text{gDW}^{-1}$ ), total flavonoid ( $38.90 \pm 1.90$  mg catechol  $\text{gDW}^{-1}$ ), and total anthocyanin ( $23.10 \pm 1.30$  mgcy-3-gluc  $\text{DW}^{-1}$ ), which are primary antioxidants (Al-Juhaimi, Ghafoor, Babiker, Matthaus, & Ozcan, 2017). Phytochemical constituents reported from different parts of *M. oleifera* are presented in Table 1 & 2.

### 3.1.1 | Chemical composition of leaf

Sixteen bioactive compounds have been identified in methanolic extract of leaves (Aja et al., 2014). The quercetin (0.07–1.26%) and kaempferol (0.05–0.67%) were major phytochemicals in leaves, and Indian MO varieties, that is, PKM-1 and PKM-2 had higher quercetin and kaempferol contrasted with the African indigenous varieties (Coppin et al., 2013). The quercetin, kaempferol, and apigenin were reported 47.0%, 30.0%, and 20.9% of the total flavonoids from leaf hydromethanolic extracts (Nouman et al., 2016). 5-Formyl-5,6,7,8-tetrahydrofolic acid, 5,6,7,8-tetrahydrofolic acid, 5-Methyl-5,6,7,8-tetrahydrofolic acid and 10-Formylfolic acid were the main folates reported from leaves (Saini et al., 2016). Chemical constituents varied with the growing environment and edaphic factors. Thirty-seven compounds were identified in leaves of MO grown at Madurai and 50 from Chennai locality (Chelliah, Ramakrishnan, & Antony, 2017). 1,30-triacontanediol (14.98%), octacosane (8.57%), Z-14-nonacosane (8.3%), and 2,2-dimethyl-1-oxa-2-silacyclotrid ecanone-13 (8.28%) were the major constituents isolated from Madurai locality. Nonacosane (15.55%) and  $\gamma$ -sitosterol (9.56%) were the major constituents from Chennai locality. Leone et al. (2015a) worked with the phenolic profiling through HPLC chromatogram of methanolic extracts of MO leaves of three different genotypes collected from Logone's Valley, Chad. As expected, flavonoids were the principal phenolic compounds. They identified 18 major compounds and two unidentified minor compounds. Quercetin-3-O-glucoside, quercetin-3-O-(6''-malonyl)glucoside, quercetin-3-O-(X''-malonyl)glucoside, kaempferol-3-O-glucoside, kaempferol-3-O-malonylglucoside were recognized in all genotypes with agreement of previous reports (Amaglo et al., 2010; Atawodi et al., 2010; Forster et al., 2015). In addition, glycosilated flavonoid, quercetin-3-O-glucoside-7-O-rhamnoside, methyl-O-quercetin-malonylglucoside, isorhamnetin-rhamnosylglucoside, rhamnosyl-glucoside, apigenin-C-glucoside, apigenin-O-glucoside, hexadecylferulate, 5-hydroxyferulic acid glucoside,

diferuloyl-(5OH-feruloyl)spermidine, glucosinolates, dicoumaroyl putrescine, di (dihydrocaffeoyl)spermidine, and two additional polyamine alkaloids were identified (Karthivashan, Fard, Arulselvan, Abas, & Fakurazi, 2013). The most abundant phenolic compounds (mg/100 g DW) in MO leaves were gallic acid ( $1.145 \pm 0.078$ ), 4-hydroxybenzoic acid ( $0.710 \pm 0.099$ ), and isorhamnetin ( $0.475 \pm 0.059$ ) followed by protocatechuic acid, catechin, caffeic acid, syringic acid, rutin trihydrate, trans *p*-coumaric acid, chlorogenic acid, trans ferulic acid, fisetin, trans resveratrol, quercetin, trans cinnamic acid, and naringenin (Al-Juhaimi, Ghafoor, Mohamed Ahmed, Babiker, & Ozcan, 2017).

### 3.1.2 | Chemical composition of leaf oil

*M. oleifera* leaves yielded 0.05% pale yellow color oil by hydrodistillation method (Marrufo et al., 2013). Chemical constituents of leaf oil of MO collected from Mozambique were oxygenated monoterpenes (linalool and  $\alpha$ -terpineol), phenolic compounds (*p*-vinylguaicol), oxygenated sesquiterpenes, that is, *cis*-dihydroagarofuran (0.1), Eudesm-11-en-4- $\alpha$ ,6 $\alpha$ -diol (0.6), and hydrocarbons, that is, eicosane (1.2), heneicosane (1.9), docosane (6.8), tricosane (8.1), tetracosane (9.7), pentacosane (13.3), hexacosane (13.9), heptacosane (11.4), octacosane (10.0), nonacosane (10.5), triacontane (1.1), 1-octadecene (0.3), octadecane (0.1), 5-Octadecin (0.3), *n*-hexadecanol (0.1), nonadecane (0.8), 1-eicosene (0.3), *n*-octadecanol (0.2), cyclopentadecanol (0.4), 1-docosene (0.4), *cis*-9-eicosen-1-ol (0.3), pseudo phytol (0.5), and two other compounds, that is, hexenyl propanoate and phenylethyl alcohol. Pentacosane (17.4%), hexacosane (11.2%), and (E)-phytol (7.7%) were main constituents of leaf oil grown in Taiwan (Chuang et al., 2007). Phytol (21.6%) and thymol (9.6%) have been reported as the most abundant compound in leaf oil of MO grown at Ceara, Brazil (Barreto et al., 2009). Mukunzi et al. (2011) recorded the correlation of chemical profile of MO leaves to the locality where the plants were grown at Rwanda and China. The Rwandan genotype contained 59 constituents, with hexanoic acid (19.8% of total volatiles) as the most abundant one, whereas the Chinese genotype contained mainly acetic acid (12.5%). Nonacosane (18.6%), 1,2,4-trimethyl-benzene (16.9%) and heptacosane (7.4%) were the main constituents in leaf oil (Zhao & Zhang, 2011). MO leaf oil contained 48.88% oleic acid, 18.09% linoleic acid, 9.32% linolenic acid, and stearic acid 2.77%. Apart from *M. oleifera*, fatty acid composition of *M. peregrina* leaf oil was investigated (Al-Juhaimi, Babiker, Ghafoor, & Ozcan, 2016). A total of 12 fatty acids were detected in the leaf oil and linolenic acid recorded the highest (32.53%).

### 3.1.3 | Chemical composition of seed

Seed oil is commercially known as "ben oil" or "behen oil" with nutty flavor and light yellow color and like olive oil (Nautiyal & Venhataraman, 1987). The seeds contain about 35% rosy yellow essential oil, 31.65% protein, 8.90% moisture, 7.54% fiber, and 6.53% ash substance (Mahmood et al., 2010). The seeds of *M. oleifera* contain 40.98% oil (Al-Juhaimi, Ghafoor, Babiker, et al., 2017), which is slight lower seed oil yield than seeds of *M. peregrina* (49.23%), of

**TABLE 1** Phytochemical constituents isolated from *Moringa oleifera* Lam.

S.N.	Compound name	Plant part	References
Flavanoids and flavanol glycosides			
1.	Rutin	Leaves	Elbatran et al. (2005); Devaraj, Krishna, and Viswanatha (2011); Habtemariam and Varghese (2015)
2.	Quercetin	Leaves	Elbatran et al. (2005); Devaraj et al. (2011); El-Alfy, Ezzat, Hegazy, Amer, and Kamel (2011)
3.	Isoquercetin	Leaves	Vongsak, Sithisarn, and Gritsanapan, (2014)
4.	Astragalín	Leaves	Vongsak et al. (2014)
5.	Isorhamnetin	Leaves	Leone et al. (2015)
6.	Kaempferol	Leaves	Manguro and Lemmen (2007); Devaraj et al. (2011); Leone, Spada, et al. (2015)
7.	Apigenin	Leaves,	Tahany et al. (2010); El-Alfy et al. (2011)
8.	Luteolin	Leaves	Leone, Fiorillo, et al. (2015)
9.	Genistein	Leaves	Leone, Fiorillo, et al. (2015)
10.	Daidzein	leaves	Leone, Fiorillo, et al. (2015)
11.	Myricetin	Leaves, seeds	Lalas and Tsaknis (2002b); Leone, Fiorillo, et al. (2015)
12.	Epicatechin	Leaves	Leone, Fiorillo, et al. (2015)
13.	Procyanidins	Roots, stem bark	Atawodi et al. (2010)
14.	Vicenín-2	Leaves	Muhammad, Arulseivan, Cheah, Abas, and Fakurazi (2016)
15.	Quercetin-3-O-glucoside	Leaves, seeds	Manguro and Lemmen (2007); Leone, Spada, et al. (2015); Maiyo, Moodley, and Singh (2016)
16.	Quercetin-3-O-(6''-malonyl) glucoside	Leaves	Leone, Fiorillo, et al. (2015)
17.	Kaempferol-3-O-glucoside	Leaves	Leone, Fiorillo, et al. (2015)
18.	Kaempferol-3-O-(6''-malonyl) glucoside	Leaves	Leone, Fiorillo, et al. (2015)
19.	Kaempferol-3-rutinoside	Leaves	Leone, Fiorillo, et al. (2015)
20.	Kaempferol-3-O- $\alpha$ -rhamnoside	Leaves	Manguro and Lemmen (2007)
21.	Kaempferide 3-O-(2'',3''-diacetylglucoside)	Leaves	Manguro and Lemmen (2007)
22.	Kaempferol-3-O-[[ $\beta$ -glucosyl-(1 $\rightarrow$ 2)]-[ $\alpha$ -rhamnosyl-(1 $\rightarrow$ 6)]- $\beta$ -glucoside-7-O- $\alpha$ -rhamnoside	Leaves	Manguro and Lemmen (2007)
23.	Kaempferide-3-O-(2''-galloylrhamnoside)	Leaves	Manguro and Lemmen (2007)
24.	Kaempferide-3-O-(2''-O-galloylrutinoside)-7-O- $\alpha$ -rhamnoside	Leaves	Manguro and Lemmen (2007)
25.	Kaempferol-3-O-[[ $\alpha$ -rhamnosyl-(1 $\rightarrow$ 2)]-[ $\alpha$ -rhamnosyl-(1 $\rightarrow$ 4)]- $\beta$ -glucoside-7-O- $\alpha$ -rhamnoside	Leaves	Manguro and Lemmen (2007)
Glucosinolate and isothiocyanate			
26.	4-( $\alpha$ -L-rhamnopyranosyloxy) benzyl glucosinolate (glucomoringin)	Leaves, seeds, root bark	Mekonnen and Drager (2003); Leone, Spada, et al. (2015); Tumer, Rojas-Silva, Poulev, Raskin, and Waterman (2015)
27.	4-[(2'-O-acetyl- $\alpha$ -L-rhamnosyloxy) benzyl] glucosinolate	Leaves	Leone, Spada, et al. (2015); Tumer et al. (2015)
28.	4-[(3'-O-acetyl- $\alpha$ -L-rhamnosyloxy) benzyl] glucosinolate	Leaves	Leone, Spada, et al. (2015); Tumer et al. (2015)
29.		Leaves	

(Continues)

TABLE 1 (Continued)

S.N.	Compound name	Plant part	References
	4-[(4'-O-Acetyl- $\alpha$ -L-rhamnosyloxy) benzyl] glucosinolate		Leone, Spada, et al. (2015); Tumer et al. (2015)
30.	4-[( $\alpha$ -L-rhamnosyloxy) benzyl] isothiocyanate	Leaves	Tumer et al. (2015)
31.	4-[(2'-O-acetyl- $\alpha$ -L-rhamnosyloxy) benzyl] isothiocyanate	Leaves	Tumer et al. (2015)
32.	4-[(3'-O-acetyl- $\alpha$ -L-rhamnosyloxy) benzyl] isothiocyanate	Leaves	Tumer et al. (2015)
33.	4-[(4'-O-acetyl- $\alpha$ -L-rhamnosyloxy) benzyl] isothiocyanate	Leaves	Mekonen and Gebreyesus (2000); Tumer et al. (2015)
34.	Benzyl glucosinolate (glucotropaeolin)	Seed	Saini et al. (2016)
35.	Sinalbin	Leaves	Leone, Spada, et al. (2015)
36.	4-[( $\beta$ -D-glucopyranosyl-1- > 4- $\alpha$ -L-rhamnopyranosyloxy) benzyl] isothiocyanate	Leaves, seeds	Maiyo et al. (2016)
Phenolic acid			
37.	Gallic acid	Leaves	Verma, Vijayakumar, Mathela, and Rao (2009)
38.	Salicylic acid	Leaves	Leone, Fiorillo, et al. (2015)
39.	Gentisic acid	Leaves	Leone, Fiorillo, et al. (2015)
40.	Syringic acid	Leaves	Leone, Fiorillo, et al. (2015)
41.	Ellagic acid	Leaves	Verma et al. (2009); Leone, Fiorillo, et al. (2015)
42.	Ferulic acid	Leaves	Verma et al. (2009); Leone, Fiorillo, et al. (2015)
43.	Caffeic acid	Leaves	Leone, Fiorillo, et al. (2015)
44.	<i>ortho</i> -Coumaric acid	Leaves	Leone, Fiorillo, et al. (2015)
45.	<i>p</i> -Coumaric acid	Leaves	Leone, Fiorillo, et al. (2015)
46.	Sinapic acid	Leaves	Leone, Fiorillo, et al. (2015)
47.	Chlorogenic acid	Leaves	Amaglo et al. (2010); Leone, Spada, et al. (2015)
48.	Cryptochlorogenic acid	Leaves	Vongsak et al. (2014)
Terpene			
49.	All-E-lutein	Pods	Teixeira, Carvalho, Neves, Silva, and Arantes-Pereira (2014); Saini et al. (2016)
50.	All-E-luteoxanthin	Pods	Saini et al. (2016)
51.	13-z-Lutein	Pods	Saini et al. (2016)
52.	15-z- $\beta$ -Carotene	Pods	Saini et al. (2016)
53.	All-E-zeaxanthin	Pods	Saini et al. (2016)
Alkaloid and sterol			
54.	4'-hydroxyphenylethanamide- $\alpha$ -L-rhamnopyranoside (Marumoside A)	Leaves	Sahakitpichan, Mahidol, Disadee, Ruchirawat, and Kanchanapoom (2011)
55.	3''-O- $\beta$ -D-glucopyranosyl derivatives (Marumoside B)	Leaves	Sahakitpichan et al. (2011)
56.	N, $\alpha$ -L-rhamnopyranosyl vincosamide	Leaves	Panda, Kar, Sharma, and Sharma (2013)
57.		Leaves	Sahakitpichan et al. (2011)

(Continues)

TABLE 1 (Continued)

S.N.	Compound name	Plant part	References
	Pyrrolemarumine-4''-O- $\alpha$ -L-rhamnopyranoside		
58.	Aurantiamide acetate	Roots	Sashidara et al. (2009)
59.	O-Ethyl-4-[( $\alpha$ -L-rhamnosyloxy)-benzyl] carbamate	Seeds	Guevara et al. (1999)
60.	Niazimicin	Leaves, seeds	Guevara et al. (1999); Jung (2014)
61.	N-benzyl, S-ethylthioformate	Root bark	Nikkon, Saud, Rahman, and Haque (2003)
62.	1, 3-Dibenzyl urea	Roots	Sashidara et al. (2009)
63.	Pterygosperrin	Seeds	Das, Kurup, Rao, and Ramaswamy (1957)
64.	Spirochin	Roots	Anonymous (2005)
65.	Niaziminin	Leaves	Murakami, Kitazono, Jiwajinda, Koshimizu, and Ohigashi (1998)
66.	$\beta$ -sitosterol	Leaves, seeds	Abd El Baky and El-Baroty (2013); Maiyo et al. (2016)
67.	$\beta$ -sitosterol-3-O- $\alpha$ -D-galactopyranoside	Stem bark	Tahany et al. (2010); Bargah and Das (2014)
Others			
68.	Oleic acid	Oil,	Nibret and Wink (2010); Gaikwad, Kale, Bhandare, Urunkar, and Rajmane (2011); Abd El Baky and El-Baroty (2013)
69.	Linoleic acid	Oil	Kleiman, Ashley, and Brown (2008); Abd El Baky and El-Baroty (2013)
70.	Myristic acid	Oil	Kleiman et al. (2008); Nibret and Wink (2010); Abd El Baky and El-Baroty (2013)
71.	Palmitic acid	Oil, roots	Kleiman et al. (2008); Nibret and Wink (2010); Abd El Baky and El-Baroty (2013)
72.	Palmitoleic acid	Oil	Abd El Baky and El-Baroty (2013)
73.	Stearic acid	Oil	Kleiman et al. (2008); Abd El Baky and El-Baroty (2013)
74.	Arachidic acid	Roots, oil	Kleiman et al. (2008); Abd El Baky and El-Baroty (2013)
75.	Linolenic acid	Oil	Abd El Baky and El-Baroty (2013)
76.	Behenic acid	Oil	Kleiman et al. (2008); Abd El Baky and El-Baroty (2013)
77.	Paullinic acid	Oil	Kleiman et al. (2008); Abd El Baky and El-Baroty (2013)
78.	Benzoic acid 4-O- $\beta$ -glucoside	Leaves	Manguro and Lemmen (2007)
79.	Benzoic acid 4-O- $\alpha$ -rhamnosyl-(1 $\rightarrow$ 2)- $\beta$ -glucoside		Manguro and Lemmen (2007)
80.	Benzaldehyde 4-O- $\beta$ -glucoside		Manguro and Lemmen (2007)
81.	Niazirin	Leaves, pod	Shanker et al. (2007); Sahakitpichan et al. (2011)
82.	Niaziridin	Pod, leaves	Shanker et al. (2007)
83.	Niazirin	Leaves	Faizi et al. (1994)
84.	Moringyne	Seeds	Memon, Memon, and Memon (1985); Nadkarni (1976)
85.	$\alpha$ -Phellandrene	Oil	Ogunbino, Flamini, Cioni, Adebayo, and Oguwande (2009)
86.	p-Cymene	Oil	Ogunbino et al. (2009); Dehshahri, Afsharypuor, Asghari, and Mohagheghzadeh (2012)
87.	Eugenol	Stem bark	Al-Asmari et al. (2015)
88.	Vanillin	Leaves, fruits, seeds	Singh et al. (2009)
89.	Benzylamine	Leaves	Iffiu-Soltesz et al. (2010)
90.	D-allose	Leaves	Al-Asmari et al. (2015)

(Continues)

**TABLE 1** (Continued)

S.N.	Compound name	Plant part	References
91.	L-arabinose, D-galactose, D-glucuronic acid, L-rhamnose	Gum	Nadkarni (1976); Shah et al. (2016)
92.	Aldotriouronic acid	Gum	Rastogi and Mehrotra (2006)
93.	D-mannose	Gum, flower	Nadkarni (1976)
94.	Leucoanthocyanin	Gum	Shah et al. (2016)
95.	Benzylglucosinolate	Seeds, roots	Nadkarni (1976)
96.	4-hydroxymellein	Stem	Nadkarni (1976)
97.	Carotene	Seed oil	Nadkarni (1976)
	Mono-palmitic	Seeds	Nadkarni (1976)
98.	Di-oleic triglyceride	Seeds	Nadkarni (1976)
99.	$\beta$ -sitosterone	Stem	Nadkarni (1976)
100.	Octacosanic	Stem	Nadkarni (1976)
101.	O-[2'-hydroxy-3'-(2''-heptenyloxy)]-propylundecanoate	Pods	Nadkarni (1976)
102.	O-ethyl,4-[( $\alpha$ -1-rhamnosyloxy)-benzyl] carbamate	Pods	Nadkarni (1976)
103.	Methyl-p-hydroxybenzoate	Pods	Nadkarni (1976)

**TABLE 2** Major phytochemical composition (% DW) of *Moringa oleifera* parts

Phytochemical	Bark	Leaf	Flour	Seed
Phenols	1.95 $\pm$ 1.63	2.65 $\pm$ 1.15	1.20 $\pm$ 0.70	1.10 $\pm$ 0.10
Flavonoids	3.70 $\pm$ 0.30	2.50 $\pm$ 0.14	2.73 $\pm$ 0.11	18.70 $\pm$ 0.50
Saponins	1.20 $\pm$ 0.70	3.20 $\pm$ 0.90	1.75 $\pm$ 0.53	13.65 $\pm$ 4.56
Anthocyanins	1.25 $\pm$ 0.11	1.60 $\pm$ 0.70	2.30 $\pm$ 0.20	2.15 $\pm$ 0.35
Alkaloids	13.83 $\pm$ 1.23	6.17 $\pm$ 3.84	6.47 $\pm$ 4.30	5.60 $\pm$ 0.60

Source: Marcel, Hubert, Bienvenu, and Pascal (2016)

which oleic acid shared 64.56% followed by behenic acid (6.98%), elaidic acid (6.39%), vaccenic acid (6.0%), palmitic acid (5.98%), and arachidic acid (4.40%), whereas cholesterol was not detected in *M. oleifera* seed oil. They reported higher crude protein and essential amino acid content in seeds than leaves. Unsaturated fatty acids shared 73.1% of MO seed oil (Stadtlander & Becker, 2017), and 28 minerals were reported in seed kernel. Aja et al. (2014) isolated five main constituents, that is, oleic acid (84%), L-(+)-ascorbic acid- 2,6-dihexadecanoate (9.80%), 9-octadecenoic acid (1.88%), methyl ester-hexadecanoic acid (1.31%), and 9-octadecenamide (0.78%) from methanolic extract of seeds. Oleic acid (73.22%) was the major fatty acid in seed, whereas 9-octadecenoic acid (20.8%) was highest pursued by palmitic, stearic, behenic, and arachidic acids 6.45%, 5.50%, 6.16%, and 4.08%, respectively, in leaf extricate (Mahmood et al., 2010). A study was conducted by Chelliah et al. (2017) to assess the chemical constituent's variability in the seeds of same genotype grown at different locations in south India. They identified 161 chemical compounds in MO seeds grown at Madurai locality and 185 compounds from Chennai locality. 6-octadecenoic acid (52.24%), n-hexadecanoic acid

(6.17%), and oleic acid (5.12%) from seeds of Madurai locality, whereas 13-docosenamide(Z)- (13.62%), propionamide (4.48%), and ethyl oleate (4.33%) from Chennai locality were major constituents.

### 3.2 | Nutritional assesment

All parts of *M. oleifera* are a storehouse of important nutrients, of which dry leaves are rich source, as presented in Table 3. Out of 13 species, *M. oleifera* is well known for high amount of nutrients such as proteins with a quality equal to that of meat, milk, and eggs (Fuglie, 2005), fat, fiber, carbohydrate, vitamins, and essential amino acids (Stadtlander & Becker, 2017). Due to less lipid content, MO leaves can be used in diet routine of the obese (Berkovich et al., 2013). Varying level of nutrient (Table 3) and mineral content (Table 4) in different plant parts was reported in previous studies. About 44.4% carbohydrate, 28.7% crude protein, 7.1% fat, and 10.9% ash content were reported from dried leaf powder. The protein profile showed 70.1% insoluble proteins, 3.5% glutelin, 3.1% albumin, 2.2% prolamin, and 0.3% globulin (Teixeira et al., 2014). Asiedu-Gyekye, Frimpong-Manso,

**TABLE 3** The nutrient compositions (100 g/sample) of different parts of *Moringa oleifera* Lam.

Nutrients	Fresh leaves	Yellow leaves	Dry leaves	Leaf powder	Pods	Seeds	Stem	Bark	Root
Protein (g)	6.70	12.00	29.40	27.10	2.50	35.97	18.66	23.95	16.87
Fats/lipids (g)	1.70	16.57	5.20	2.30	0.10	38.67	12.2	17.47	10.80
Carbohydrate (g)	12.50	26.93	41.20	38.20	3.70	8.67	20.4	12.33	14.92
Total fiber (g)	0.90	-	12.50	19.20	4.80	2.87	41.60	25.75	45.43
Vitamin B1 (mg)	0.06	-	2.02	2.64	0.05	0.05	-	-	-
Vitamin B2 (mg)	0.05	-	21.30	20.50	0.07	0.06	-	-	-
Vitamin B3 (mg)	0.80	-	7.60	8.20	0.20	0.20	-	-	-
Vitamin C (mg)	220.00	-	15.80	17.30	120.00	4.50	-	-	-
Vitamin E (mg)	448.00	-	10.80	113.00	-	-	-	-	-
Ca (mg)	440.00	1576.00	2185.00	2003.00	30.00	751.67	12.55	26.41	28.61
Mg (mg)	42.00	471.40	448.00	368.00	24.00	45.00	1.30	1.09	4.38
P (mg)	70.00	0.83	252.00	204.00	110.00	635	-	-	-
K (mg)	259.00	131.20	1236.00	1324.00	259.00	75.00	82.98	25.98	86.06
Fe (mg)	0.85	96.68	25.60	28.20	5.30	5.20	0.28	2.25	0.50
S (mg)	-	-	-	870.00	137.00	-	-	-	-

Source: Gopalakrishnan, Doriya, and Kumar (2016); Fokwen et al. (2018); Gopalakrishnan et al. (2016); Fuglie (2005); Olagbemide and Alikwe (2014); Verma and Nigam (2014)

**TABLE 4** Mineral profiling of leaf and kernel samples of different *Moringa* species

Mineral (per kg DM)	<i>M. oleifera</i>				<i>M. drouhardtii</i>		<i>M. hilderbrandtii</i>		<i>M. Stenoptala</i>
	Leaves			Kernel	Leaves	Kernel	Leaves	Kernel	Kernel
	India (Asia)	Malawai (Central Africa)	Nicaragua (Central America)	Ethiopia (East Africa)	Madagascar (East Africa)		Gran Canary (West Africa)		Israel (West Asia)
Ca (g)	11.00	22.20	20.80	2.53	11.90	0.75	15.70	0.87	1.45
P (g)	6.32	3.70	3.36	8.28	2.190	3.55	4.66	8.03	3.62
Mg (g)	3.73	8.79	3.69	4.15	9.06	1.83	4.90	2.79	2.37
K (g)	29.60	13.50	18.80	11.30	12.30	7.84	19.50	9.03	5.64
Na (g)	0.67	0.60	0.70	0.04	9.78	0.15	5.04	0.06	0.49
Fe (mg)	132.0	1239.0	229.0	123.0	138.0	39.70	350.0	24.10	21.20
Zn (mg)	30.10	21.70	20.70	34.20	12.20	14.70	21.80	27.00	21.70
Mn (mg)	16.70	76.90	71.40	17.70	79.00	5.34	80.50	5.11	9.16
Mo (mg)	0.518	4.53	1.30	<0.241	1.46	0.283	1.60	<0.243	<0.238
Co (mg)	0.080	0.860	0.083	0.081	<0.082	<0.081	0.248	<0.081	<0.08
Se (mg)	1.570	27.70	2.85	<0.644	<0.653	<0.645	<0.645	0.796	<0.636
Si (mg)	248.0	279.0	343.0	124.0	389.0	33.50	192.0	5.38	9.73
Cu (mg)	5.89	8.21	5.37	4.91	4.23	3.56	2.52	3.91	3.01
B (mg)	29.10	64.60	32.20	7.93	126.0	7.35	62.70	6.31	6.14
V (mg)	0.136	3.64	0.276	0.189	0.215	<0.081	0.733	<0.081	<0.08
Al (mg)	65.30	862.0	79.70	65.70	80.60	15.40	301.0	1.30	2.81
Ni (mg)	1.49	3.80	0.681	0.886	0.729	1.560	1.86	0.741	1.66

Note: Minerals are (g/kg or mg/kg) dry matter basis.

Source: Stadlander and Becker (2017).

Awortwe, Antwi, and Nyarko (2014) reported 35 elements (14 macroelements and 21 microelements) from leaf aqueous extract. The leaf powder have high contents (mg/100 g) of Ca (2787–3307),

K (2672–4892), Mg (553–1566), Cl (188–332), Na (14.78–17.76), Fe (14.11–16.57), and other minerals (Mulyaningsih & Yusuf, 2018). MO leaves can provide 1,000 mg Ca and 4,000 mg by dry leaf powder,



whereas 8 ounces of milk can give 300–400 mg only (Gopalakrishnan et al., 2016). In fact, MO fresh leaves contain two times more protein than milk (Thurber & Fahey, 2009), nine times of protein than yoghurt, four times of Ca than milk, 25 times of Fe than spinach (Rockwood, Anderson, & Casamatta, 2013) or 28 mg of Fe from 100 g of leaves (Fuglie, 2005), 63 times K than milk, and three times of banana, 36 times Mg than egg, four times vitamin A than carrots, and 13 times of spinach, seven times vitamin C than orange, four times of vitamin B than pork meat, 50 times of vitamin B2 than sarones, 50 times of vitamin B3 than peanut, six times of vitamin E than rapeseed oil, two times of amino acids than black vinegar, and 30 times more of brown rice (Ganatra, Umang, Payal, Tusharbindu, & Pravin, 2012).

MO leaves are rich source of essential amino acids, higher than adequate with the standards of WHO, FAO, and UNO recommendations for small children and approximately equal to soyabean seeds in terms of digestibility score (Fuglie, 2005). As per the study conducted by Al-Juhaimi, Ghafoor, Mohamed Ahmed, et al. (2017), amino acids profile in leaves and seeds (g/100 g, respectively) of MO is glutamic acid ( $2.660 \pm 0.13$ ;  $3.724 \pm 0.18$ ), aspartic acid ( $2.185 \pm 0.06$ ;  $3.059 \pm 0.02$ ), leucine ( $2.070 \pm 0.15$ ;  $2.898 \pm 0.22$ ), arginine ( $1.820 \pm 0.06$ ;  $2.548 \pm 0.08$ ), alanine ( $1.605 \pm 0.33$ ;  $2.247 \pm 0.46$ ), phenylalanine ( $1.595 \pm 0.06$ ;  $2.233 \pm 0.09$ ), lysine ( $1.540 \pm 0.14$ ;  $2.156 \pm 0.20$ ), glycine ( $1.450 \pm 0.11$ ;  $2.030 \pm 0.16$ ), valine ( $1.345 \pm 0.12$ ;  $1.883 \pm 0.17$ ), proline ( $1.280 \pm 0.11$ ;  $1.792 \pm 0.16$ ), threonine ( $1.265 \pm 0.13$ ;  $1.771 \pm 0.19$ ), isoleucine ( $1.155 \pm 0.03$ ;  $1.617 \pm 0.05$ ), serine ( $1.060 \pm 0.04$ ;  $1.484 \pm 0.06$ ), tyrosine ( $0.915 \pm 0.08$ ;  $1.281 \pm 0.11$ ), histidine ( $0.730 \pm 0.03$ ;  $1.022 \pm 0.04$ ), hydroxylysine ( $0.690 \pm 0.04$ ;  $0.966 \pm 0.06$ ), methionine ( $0.560 \pm 0.08$ ;  $0.784 \pm 0.12$ ), tryptophan ( $0.510 \pm 0.03$ ;  $0.714 \pm 0.04$ ), cysteine ( $0.280 \pm 0.04$ ;  $0.392 \pm 0.06$ ), taurine ( $0.105 \pm 0.02$ ;  $0.147 \pm 0.03$ ), hydroxyproline ( $0.100 \pm 0.01$ ;  $0.140 \pm 0.02$ ), and ornithine ( $0.060 \pm 0.01$ ;  $0.084 \pm 0.02$ ), whereas glutamic acid, aspartic acid, argentine, proline, and glycien are major amino acids in seed kernel (Stadtlander & Becker, 2017). A very recent study (Pasha et al., 2019) carried out for quantification of the important molecules in different parts of the tree and an understanding of the protein chain reaction that leads to creation of such molecules. Enzymes in the biosynthesis of vitamins and metabolites like quercetin and kaempferol are highly expressed in leaves, flowers, and seeds. The expression of Fe, Zn, Mg transporters, and Ca storage proteins were observed in root and leaves, respectively. In general, leaves retain the highest amount of small molecules of human interest to combat the malnutrition (Pasha et al., 2019).

The immature flowers and pods are the source of palmitic, linolenic, linoleic, and oleic acids (Sanchez-Machado, Nunez-Gastelum, Reyes-Moreno, Ramirez-Wong, & Lopez-Cervantes, 2010). The green pods are the source of vitamin C with range of 92 to 126 mg/100 g of pulp (Dogra, Singh, & Tandon, 1975). Immature pods contain approximately 46.78% fiber, 20.66% protein, 30% of amino acid content, whereas leaves and flowers have 44% and 31% amino acid content, respectively (Sohani, 2018). The seeds of Indian MO collected from Assam contain 40.34% protein with seven essential amino acids, 39.12% fats, K (2,357.71 mg/kg), Na (1074.09 mg/kg), Mg (972.06 mg/kg), Ca (121.14 mg/kg), and iron (36.2 mg/kg; Liang, Wang, Li,

Chu, & Sun, 2019). The root bark is rich in moringine alkaloid, stimulating drug component similar to ephedrine (Morton, 1991). Study carried out by Lalas and Tsaknis (2002a) demonstrated that seed oil contains around 76% PUFAs, that is, palmitic, linolenic, linoleic, and oleic acids. Refined seed oil of MO is the best substitute of olive oil (Morton, 1991). Seeds are the best substitute to legumes due to high content of essential amino acids (Ferreira, Farias, Oliveira, & Carvalho, 2008). Thus, MO is very important plant as it is exceptional to discover such a high nutrient profile in a solitary tree (Razis, Ibrahim, & Kntayya, 2014).

It is an established fact that mineral content change with the species and also with geographical area where the plant grows (Table 4). Apart from this, factors of locality of different geographical areas also affect the nutrients level in the same cultivar of MO when it grows at different localities (Table 5). Leone et al. (2015a) clearly recorded the huge differences in nutritional status in leaves of MO planted in three different countries viz., Chad, Algeria, and Haiti. Chelliah et al. (2017) reported wide variation in nutraceuticals and phytochemical profiling of MO grown at two different regions. Fuglie (2005) revealed that nutrient content is also influenced by seasons. Vitamin A was recorded higher in leaves in the summer–rainy season, whereas vitamin C and Fe were high in the winter season (Yang et al., 2006). Safaeian, Asghari, Javanmard, and Heidarinejad (2015) stated that MO is majorly used in summer season due to the higher concentration of bioactive compounds. The concentration of phenolic compound was higher in leaves harvested in winter season (Shih, Chang, Kang, & Tsai, 2011). Moyo et al. (2011) clearly mentioned that the variation in nutritional profile is significantly attributed due to locality and ecological factors. Phytochemical variations were also reported in different 13 MO cultivars collected from different parts of the globe (Ndhlala et al., 2014). These variations could be due to many factors including genetic variations, soil, climate, time of harvest, and storage conditions.

Phytochemicals, for example, alkaloids, flavonoids, anthocyanins, tannins, sterols, terpenoids, saponins, anthraquinones, glucosinolates, isothiocyanates, glycosides, and glycerol-1-9-octadecanoate isolated from different plant parts of MO considered as antinutritional factors are responsible for medicinal properties (Berkovich et al., 2013). The amounts of antinutrients, that is, alkaloids, flavonoids, oxalates, phytates, saponins, tannins are generally low and do not have any health risk to animal and human beings (Gidamis, Panga, Sarwatt, Chove, & Shayo, 2003; Nouman et al., 2013). Teixeira et al. (2014) documented the antinutritional factors, such as, tannins (20.7 mg/g), trypsin inhibitor (1.45 mg/g), nitrates (17 mg/g), and oxalic acids (10.5 mg/g) in leaves. However, these antinutrients can be removed through soaking, boiling, or even frying (Igwilo, Oloyode, & Enemor, 2007). Saponins for instance, confer bitter taste on moringa without any harmful effects (Makkar & Becker, 1996). Saponin is used as an antibiotic and high blood pressure (Ajayi, Akomolafe, & Adefioye, 2014). *Moringa* leaves have a high proportion of insoluble form of oxalates (Noonan & Savage, 1999). The comparatively higher amounts of antinutrients in the seeds rather than the leaves may explain why *Moringa* seeds are used more in ethno-medicine and are used to cure the diseases (Sallau, Mada, Ibrahim, & Ibrahim, 2012; Stevens, Ugese,

**TABLE 5** Nutritional characterization of leaf of *Moringa oleifera* due to geographical locations

Nutrients	India (Mean ± SD)	Algeria (Mean ± SD)	Ethiopia (Mean ± SD)	Nigeria (Mean ± SD)	South Africa (Mean ± SD)	Brazil (Mean ± SD)	Haiti (Mean ± SD)	Mexico (Mean ± SD)
Proteins (g/100 g)	28.40 ± 0.20	30.60 ± 0.80	10.71 ± 0.81	17.01 ± 0.1	30.29 ± 1.48	28.65 ± 0.04	20.80 ± 0.01	11.48 ± 1.4
Lipids (g/100 g)	1.90 ± 0.30	5.60 ± 0.30	10.31 ± 1.2	2.11 ± 0.11	6.5 ± 1.04	7.09 ± 0.43	7.05 ± 0.11	10.21 ± 1.83
Fibre (g/100 g)	19.20 ± 0.20	32.80 ± 0.20	8.05 ± 0.02	7.09 ± 0.11	11.40 ± 0.4	-	37.63 ± 1.00	9.46 ± 1.14
Ash (g/100 g)	3.00 ± 0.10	15.10 ± 0.30	7.29 ± 0.84	7.93 ± 0.12	7.64 ± 0.43	10.9 ± 0.8	9.62 ± 0.02	11.18 ± 0.19
Ca (mg/100 g)	2225 ± 0.1	2997 ± 27.00	2016.5 ± 22.6	1910 ± 0.08	3650 ± 0.04	2970 ± 0.0	2150.26 ± 56.07	2620.5 ± 5.6
Mg (mg/100 g)	3760.2 ± 0.30	-	322.5 ± 0.0	380 ± 0.01	500 ± 0.05	1900 ± 0.0	533.51 ± 23.87	340.6 ± 2.8
P (mg/100 g)	250 ± 0.3	-	-	301.5 ± 0.5	300 ± 0.01	-	-	-
K (mg/100 g)	1391 ± 0.1	1492 ± 13.00	1845 ± 7.0	970 ± 0.01	1500 ± 0.02	4160 ± 0.0	-	1817 ± 14.1
S (mg/100 g)	870 ± 0.1	-	-	-	630 ± 0.15	-	-	-
Na (mg/100 g)	-	502 ± 5.00	8.13 ± 0.6	192.95 ± 4.4	164 ± 0.02	-	262.50 ± 5.45	40.78 ± 0.7
Fe (mg/100 g)	29.3 ± 0.2	30.20 ± 0.30	19.37 ± 6.6	107.48 ± 8.2	49 ± 49.65	103.12 ± 0.0	11.91 ± 0.82	7.07 ± 0.4
Cu (mg/100 g)	0.6 ± 0.1	-	1.03 ± 0.47	6.10 ± 0.19	0.83 ± 0.14	3.38 ± 0.0	-	0.41 ± 0.0

Source: Chelliah et al. (2017); Leone et al. (2018); Amabye (2016); Ogbé and Affiku (2011); Moyo, Masika, Hugo, and Muchenje (2011); Teixeira et al. (2014); Leone et al. (2015); Valdez-Solana et al. (2015).

Otitoju, & Baiyeri, 2015). Roots are the rich source of antinutritional compounds; therefore, it is traditionally used in African, Asian, and Latin American countries (Iqwilo et al., 2014).

### 3.3 | *M. oleifera* as potential food

The different parts of MO are used in more than 80 countries to supplement mineral and vitamin deficiencies (Mahmood et al., 2010). The foliages, flowers, green pods, seeds, and seed oil contain low saturated fatty acid and high amount of monounsaturated unsaturated fatty acids and PUFAs; therefore, it can be used for value addition of food products (Saini et al., 2016). Fried flowers have taste like mushrooms (Owusu & Oduro, 2011). Immature pods are widely used in *sambhar*, *corma*, *dal*, *cutlet*, and *curry* dishes by mixing with coconut, poppy seeds, and mustard in Indian subcontinent (Ganatra et al., 2012). The immature seeds are used as peanuts with oil frying (Ramachandran, Peter, & Gopalakrishnan, 1980). Peeled dried roots mixed with acetic acid are consumed as a condiment (Martin & Ruperte, 1979). The leaf, seed, and flower powder are utilized in various food applications such as in fortifying amala (stiff dough), ogi (maize gruel), bread, biscuits, yoghurt, cheese, and soups (Oyeyinka & Oyeyinka, 2018). MO leaves alone or in combination with spinach, melon etc. can be used as ingredient in soups (Babayehu et al., 2014). Leaf powder was reported to increase the swelling and pasting properties along with protein and macronutrients in fortified plantain flour (Karim, Kayode, Oyeyinka, & Oyeyinka, 2015). Flower powder or leaf powder is known to increase nutritional value of weaning foods (Arise, Arise, Sanusi, Esan, & Oyeyinka, 2014). Paneer with extract of MO leaf of different concentration were investigated, and it was found to have high nutrient content than normal paneer (Sachan, Khan, Yadav, & Sonkar, 2010). The protein, crude fiber, and ash content increased appreciably with increasing concentration of *Moringa* leaf powder in chocolate and

halwa (Abou-Zaid & Nadir, 2014). Biscuits incorporated with 5% leaf powder were reported to increase protein content by 14%, and these are known as *Moringa* biscuits (Alam et al., 2014). Bread fortified with 5% of MO leaf powder was found to have 17% and 88% increased in protein and dietary fiber content (Chinma, Abu, & Akoma, 2014).

Moisture, protein, fiber, and ash content increased, whereas total fat and carbohydrate content decreased with increased amount of *Moringa* leaf (2, 4, 6, 8, and 10 g) in cakes (Chinma et al., 2014). Leaves increased the taste of yoghurt and did not negatively affect the growth of *Lactobacillus rhamnosus* GR-1 (Hekmat, Morgan, Soltani, & Gough, 2015). Incorporation of dried leaves (sun dried, shadow dried, and mechanically dried) in proportion of 0%, 2%, 4%, 6%, 8%, 10% increased the moisture, fat, ash, protein, carbohydrate, and antioxidant activity of khakhra (Maghu, Sharma, & Younis, 2017). Chin chin is a Nigerian snack product, which is made of wheat flour, butter, egg, and milk. Oven-dried leaves mixed with chin chin had high Ca (190.5 mg/100 g), high Zn with sun dried leaves (7.1 mg/100 g), and high Fe (51.3 mg/100 g) with shade dried leaves (Emelike & Eberé, 2016). Salama, Sayed, and Abdalla (2017) reported that 0.5% of MO dried leaves and 6% of oil used to prepare fortified ice milk to increase nutritional quality and suitable sensory aspects. MO dried leaf powder at 12% concentration (per 55 g of flour used) has been used in the production of muffin to increase ash content, protein, fat, beta carotene, vitamin C, and minerals (Srinivasamurthy, Yadav, Sahay, & Singh, 2017).

A very recent study showed the effect of the administration of dried leaf powder of MO on the nutritional status and body composition of a sample of malnourished children, aged between 4 and 18 years in Zambia (Barichella et al., 2019). It was found that daily dose of 14 g dried leaf powder was safe from any side effects and well accepted in terms of taste and visual attractiveness as dietary supplementation for 30 days, homogeneously distributed between lunch and dinner. Mild side effects were reported independently of age in three

out of 16 subjects within 3 days when 20 g daily MO leaf powder was supplemented. However, no significant effect was reported on body weight and body composition. These finding was not taken for granted as previous studies have mainly investigated the acceptability of fortified baked products like cookies, cereals, and crackers in adults (Owusu & Oduro, 2011).

### 3.4 | Therapeutic activities

*M. oleifera* has various medicinal uses, which have long been recognized and used in the Ayurvedic and Unani systems of medicine (Mughal, Ali, Srivastava, & Iqbal, 1999). The extracts of different part of MO offer a high level of safety without any adverse effects on human being (Saini et al., 2016). Pyridine-3-carboxamide and 2-myristoyl-glycinamide isolated from leaves and seeds have been used against approximately 80 diseases (Cohen & Grifo, 2007). The biological and medicinal practices attributed to different parts of *M. oleifera* are summarized here, of which some are outlined in Table 6.

#### 3.4.1 | Antibacterial activity

Numerous investigations have recommended that different extracts from various tissues of *M. oleifera* showed antibacterial properties against both gram-negative and gram-positive bacteria viz., fresh leaf juice against *Pseudomonas aeruginosa* and *Staphylococcus aureus* (Caceres, Cabrera, Morales, Mollinedo, & Mendia, 1991); aqueous leaf extract against *S. aureus*, *Vibrio cholerae*, and *Escherichia coli* (Vieira, Mourao, Angelo, Costa, & Vieira, 2010); methanol extract of leaves against 13 different bacterial strains tested, including *E. coli*, *Enterobacter aerogenes*, *Klebsiella pneumonia*, *P. aeruginosa*, and *Providencia stuartii* (Dzotam et al., 2016); ethanol extracts of foliages and seeds against *Trichophyton rubrum*, *T. mentagrophytes*, *Epidermophyton floccosum*, and *Microsporum canis* (Chuang et al., 2007); chloroform and ethanol extracts of seeds and foliage against *E. coli*, *P. aeruginosa*, *S. aureus*, and *E. aerogenes* (Bukar, Uba, & Oyeyi, 2010); chloroform extract of flowers and ethanol extract of pods against *V. cholera*, *V. vulnificus*, and *V. mimicus* (Brilhante et al. (2015); water-soluble seed lectin against *S. aureus* and *E. coli* (Ferreira et al., 2011); phenolics rich extract of seed flour against *Bacillus cereus*, *S. aureus*, *E. coli*, and *Yersinia enterocolitica* (Govardhan-Singh, Negi, & Radha, 2013); hexane, petroleum ether, butanol, chloroform, acetone, ethyl acetate, methanol, and water extracts of pod husk against *S. aureus*, *S. epidermidis*, *Salmonella typhimurium*, *E. coli*, and *K. pneumonia* (Arora & Onsare, 2014); acetone extract of foliage against *S. aureus*, *Enterococcus faecalis*, *E. coli*, *P. aeruginosa*, *Candida albicans*, *Aspergillus fumigates*, and *Cryptococcus neoformans* (Ratshilivha, Awouafack, du Toit, & Eloff, 2014); aqueous, methanol and ethyl acetate extracts of seeds against *E. coli*, *K. pneumoniae*, *Proteus mirabilis*, *P. aeruginosa*, and *S. aureus* (Emmanuel et al., 2014); ethyl acetate extract of bark against *S. aureus*, *Citrobacter freundii*, *B. megaterium*, and *P. fluorescens* (Zaffer et al., 2014); flavonoids extract of seeds against *S. aureus*, *P. aeruginosa*, and *C. albicans* (Onsare & Arora, 2015).

Antimicrobial properties are possibly attributed by 4-( $\alpha$ -L-rhamnopyranosyloxy)benzyl isothiocyanate, methyl N-4-( $\alpha$ -L-rhamnopyranosyloxy)benzyl carbamate and 4-( $\beta$ -D-glucopyranosyl-1  $\rightarrow$  4- $\alpha$ -L-rhamnopyranosyloxy)-benzyl thiocarboxamide (Oluduro, Aderiye, Connolly, Akintayo, & Famurewa, 2010; Padla et al., 2012); water soluble lectine protein (Coelho et al., 2009; Ferreira et al., 2011; Moura et al., 2015); flocculating cationic polypeptides (Shebek et al., 2015); niazimicin (Rim et al., 2014); glycosides 4-( $\alpha$ -L-rhamnosyloxy)-benzyl isothiocyanate, 4-( $\alpha$ -L-rhamnosyloxy)-phenylacetoneitrile and moringine (Jahn, Musnad, & Burgstaller, 1986); proanthocyanidins (Maldini et al., 2014) in seeds; niaziminin, niazinin (Wang, Chen, & Wu, 2016), and silver nanoparticles (Prasad & Elumalai, 2011) in leave extract; cardiac glycosides in pods (Arora & Onsare, 2014); kaempferol, rhamnetin, kaempferitin, isoquercitrin, and pterygospermin in flowers; spirochin and anthonine in roots (Farooq, Rai, Tiwari, Khan, & Farooq, 2012; Mehta, Shukla, Bukhariya, & Charde, 2011; Raj, Gopalakrishnan, Yadav, & Dorairaj, 2011); and aglycon of deoxy-niazimicine (N-benzyl, S-ethyl thioformate) in bark (Nikken et al., 2003). It is cleared from previous studies that seeds may also act directly upon microorganisms and showed broader antimicrobial spectrum than other parts of MO. Methods used to assess the antibacterial activities are inhibition halos test (De-Martino, De Feo, Fratianni, & Nazzaro, 2009) and total polyphenols test (Marrufo et al., 2013) for leaf essential oil; modified disk diffusion method for aqueous and ethanol extracts of leaves (Peixoto et al., 2011); thin layer chromatography bioassay for methanol crude extract of seeds (Oluduro et al., 2010); disc diffusion method for hexane and methonolic extracts of leaves and seeds (Chelliah et al., 2017); modified Kirby-Bauer disk diffusion method for aqueous and ethanolic extracts of seeds (Vieira et al., 2010); and pour plate method for seed flour (Singh, Negi, & Radha, 2013). Thus, the extracts of different parts of MO could be an effective source of natural antimicrobial agents with versatile potential applications in the pharmaceutical industries.

#### 3.4.2 | Antifungal activity

MO has been used for antifungal activities in several previous investigations. Extract of seeds (Chuang et al., 2007; Jabeen, Shahid, Jamil, & Ashraf, 2008; Oluduro et al., 2010), leaves (Kekuda et al., 2010; Oluduro, 2012), and leaf oil (Chuang et al., 2007) are reported to exhibit significant antifungal activity against *T. mentagrophyte*, *Pullarium* spp., *A. flavus*, *Penicillium* spp., *A. niger*, *A. oryzae*, *A. terreus*, *A. nidulans*, *F. solani*, *R. solani*, *C. cladosporioides*, *P. sclerotigenum*, and *Dermatophytes* (*T. rubrum*, *E. xoccosum*, *M. canis*). Growth of saprophytic fungi was inhibited by extract of MO leaves (Ayanbimpe et al., 2009). A metabolite griseofulvin isolated from endophytic fungus of MO demonstrated inhibitory effect on the growth of 8 plant pathogenic fungi in an in vitro antifungal test (Zhao, Zhang, Wang, Wang, & Zhang, 2012). The essential oil of MO was evaluated using different concentration of total polyphenols against *P. aurantiogriseum*, *P. expansum*, *P. citrinum*, *P. digitatum*, and *A. niger* using inhibition halo technique test (Marrufo et al., 2013). MO roots have an anti-infection compound, pterygospermin, which has incredible fungicidal effects

**TABLE 6** Biological and medicinal attributes of *Moringa oleifera* Lam.

Traditional uses/ effects	Mode of action	References
Biocidal activities		
Bacterial	<i>Aeromonas caviae</i> , <i>Bacillus cereus</i> , <i>B. megaterium</i> , <i>B. subtilis</i> , <i>Citrobacter freundii</i> , <i>E. coli</i> , <i>Enterobacter sp.</i> , <i>Enterobacter aerogenes</i> , <i>Enterococcus faecalis</i> , <i>K. pneumoniae</i> , <i>Microcystis aeruginosa</i> , <i>Mycobacterium phlei</i> , <i>Providencia stuartii</i> , <i>Proteus vulgaris</i> , <i>P. aeruginosa</i> , <i>P. fluorescens</i> , <i>Propionibacterium acnes</i> , <i>Serratia marcescens</i> , <i>S. boydii</i> , <i>S. dysenteriae</i> , <i>Sarcina lutea</i> , <i>S. epidermidis</i> , <i>S. pyogenes</i> , <i>S. mutans</i> , <i>S. aureus</i> , <i>Streptococcus mutans</i> , <i>S. typhii</i> , <i>Vibrio cholera</i> , <i>V. parahaemolyticus</i>	Nwosu and Okafor (1995); Nikkon et al. (2003); Chuang et al. (2007); Lurling and Beekman (2010); Rahman et al. (2010); Peixoto et al. (2011); Saadabi and Abu-Zaid (2011); Walter, Samuel, Peter, and Jospheh (2011); Padla, Solis, Levida, Shen, and Ragasa (2012); Rattanasena (2012); Galuppo et al. (2013); Marrufo et al. (2013); Patel, Patel, Patel, Desai, and Meshram (2014); Zaffer et al. (2014); Eyarefe, Idowu, and Afolabi (2015); Dzotam, Touani, and Kuete (2016); Elgamily et al. (2016)
Fungal	<i>Aspergillus flavin</i> , <i>A. niger</i> , <i>Basidiobolus ranarum</i> , <i>B. haptosporus</i> , <i>Candida albicans</i> , <i>Epidermophyton floccosum</i> , <i>Microsporium canis</i> , <i>Penicillium aurantiogriseum</i> , <i>P. citrinum</i> , <i>P. digitatum</i> , <i>P. expansum</i> , <i>Trichophyton mentagrophytes</i> , <i>T. rubrum</i>	
Dental caries	Filling cavities with gum; juice of root bark; ethyl acetate, acetone, and ethanol extracts of seeds, roots and leaves	Bhattacharya, Das, and Banerji (1982); Prakash, Tiwari, Shukla, Mathur, and Tiwari (1987); Elgamily et al. (2016)
Gastrointestinal disease	4-( $\beta$ -D-glucopyranosyl-1 $\rightarrow$ 4- $\alpha$ -L-rhamnopyranosyloxy)-benzyl thiocarbonyl amide from seed extract	Oluduro (2012)
Urinary tract infection	Phenolic compound of stem bark decoction	Maurya and Singh (2014)
Tuberculosis	Root bark	Prakash et al. (1987)
Syphilis	Gum	Bhattacharya et al. (1982)
Parasites		
Dracunculiasis (guinea-worm)	Leaf extract	Fabiya, Kela, Tal, and Istifanus (1993)
Lymphatic filarial nematode <i>Brugia malayi</i> (Round-worm)	Gum extract inhibited 100% motility and 56% MTT reduction potential of the adult female worms	Kushwaha et al. (2011)
Protozoa	Lectin protein from seed extract	Kohler et al. (2002)
Schistosomes	Seed flocculation	Olsen (1987)
Nonspecified pathogens		
Cholera	Pterospersmin isolated from flowers	Lizzy, Rao, and Puttaswamy (1968)
Conjunctivitis	Bark, leaf juice drops	Yabesh, Prabhu, and Vijayakumar (2014); Mbikay (2012)
Cataractogenesis	Leave ethanol extract	Kurmi, Ganeshpurkar, Bansal, Agnihotri, and Dubey (2014)
Cataracts	Consumption of leaves, pods	Anwar et al. (2007)
Muscle disease	Leaves	Armand-Stussi, Basocak, Pauly, and McCaulley, (2003)
Pyoderma (skin sores)	Seed extract	Caceres and Lopez (1991); Bharali, Tabassum, and Azad (2003)
Throat infection	Flower extract	Fuglie (1999)
Cancer/protection		

(Continues)

TABLE 6 (Continued)

Traditional uses/ effects	Mode of action	References
Antiproliferation through apoptosis	Cancer cell line inhibition against A549, B16F10, Hep-G2, Panc-1, p34, COLO 357, MDA-MB-231, HCT-8, MCF-7, HeLa, CACO-2, L929, HCT-16, HCT18, PC3, K562, THP-1, T47D, HL-60, Colo-205, SW480, HepG2, Caco-2, MCF-7 cells	Monera, Wolfe, Maponga, Benet, and Guglielmo (2008); Pamok, Vinitketkumuen, and Saenphet (2012); Waivaput, Payungporn, Issara-Amphorn, Nattanan, and Panjavorayan (2012); Berkovich et al. (2013); Gismondi et al. (2013); Tiloke, Phulukdaree, and Chuturgoon (2013); Al-Asmari et al. (2015); Diab, Guru, Bhushan, and Saxena (2015); Elsayed, Sharaf-Eldin, and Wadaan (2015); Jung, Lee, and Kang (2015); Madi, Dany, Abdoun, and Usta J. (2016)
Ovarian cancer, Prostate cancer, Breast cancer	Leaf extract	Zayas-Viera, Vivas-Mejia, and Reyes (2016)
Photo-oxidative damage	Chlorogenic acid from leaf	Ramabulana et al. (2016)
Radioprotective	Leaf extract	Rao, Devi, and Kamath (2001)
Circulatory/endocrine disorders	Leaf extract	Munim, Puteri, and Sari (2016)
Anti-anemic	Ethanollic leaf extract	Mehta, Balaraman, Amin, Bafna, and Gulati (2003)
Anti-atherosclerotic	Pod extract	Gupta and Arya (2011); Thakur, Soren, Pathapati, and Buchineni (2016)
Diuretic activity	Aqueous infusion of seeds, decoction of leaves, leaf juice	
Detoxification		Sutar et al. (2008)
Analgesic	Antagonism of NMDA receptors through leaf extract	Dhakar, Maurya, Pooniya, Bairwa, and Gupta (2011)
Anti-aging	Zeatin from fresh leaf juice	Bhattacharya et al. (1982); Hukkeri, Nagathan, Karadi, and Patil (2006); Ahmad, Ejaz, Anwar, and Ashraf (2014)
Antipyretic	Gum, ethanol and ethyl acetate extracts of seed	Ranjan, Swarup, Patra, and Chandra (2009)
Antitoxidant to fluoride toxicity	Seeds	Fuglie (1999); Rajendran, Balaji, and Basu (2008)
Antidote to snakebite and scorpion-bite	Oral doses of powdered stem bark	Sing et al. (2009)
Oxidative DNA damage protective	Aqueous extract of pods and seeds	
Digestive disorders		
Anticollitis	Hydro-alcoholic extract and its chloroform fraction of seeds	Minaiyan, Asghari, Taheri, Saeidi, and Nasr-Esfahani (2014)
Cholagogue	Leaves	Armand-Stussi et al. (2003)
Diarrhea and dysentery	Ethanol leaf extract, gum, root	Misra, Srivastava, and Srivastava (2014); Bhattacharya et al. (1982); Agrawal and Mehta (2008)
Purgative, laxative	Seed oil, Roots	Fuglie (1999); Agrawal and Mehta (2008)
Flatulence/Carminative	Roots	Agrawal and Mehta (2008)

(Continues)

TABLE 6 (Continued)

Traditional uses/ effects	Mode of action	References
Hepatic gluconeogenesis	Isothiocyanate extract of leaves	Waterman et al. (2015)
Liver fibrosis	Seed extract	Hamza (2010)
Trypsin inhibitor	Flower extract had antilarvicidal activity on <i>Aedes aegypti</i>	Pontual et al. (2014)
Nervous disorders		
Antinociceptive	Methanol extract of leaves, Chitin-binding protein from seeds	Adedapo, Falayi, and Oyagbemi (2014); Mirella et al. (2011)
Epilepsy and anxiety	Aqueous extract of leaves	Ingale and Gandhi (2016)
Paralytic affections	Roots	Agrawal and Mehta (2008)
Down-regulation of nuclear factor kappa-B	Aqueous extract of leaves	Berkovich et al. (2013)
Upregulation of TNF- $\alpha$	Ethanol extract of leaves	Akanni, Adedeji, and Oloke (2014)
Respiratory disorders		
Asthma	Gum, moringine and moringine from seed extract	Bhattacharya et al. (1982), Agrawal and Mehta (2008)
Bronchitis	Moringine and moringine from seed extract	Mahajan and Mehta (2008)
Reproductive health		
Abortifacient	Leaves extract, gum, aqueous root extract	Armand-Stussi et al. (2003); Bhattacharya et al. (1982); Agrawal and Mehta (2008)
Aphrodisiac	Leaves, methanolic extract of seeds	Armand-Stussi et al. (2003); Goswami et al. (2016)
Lactation enhancer	Dry leaf capsule	Raguindin, Dans, and King (2014)
Neural tube defects during pregnancy	Folate from leaf extract	Williams et al. (2015)
Prostate function	Zeatin from leaves, seed oil	Dhakar et al. (2011); Fuglie (1999)
Sexual hormonal balance	Aqueous extract of roots	Shukla, Mathur, and Prakash (1988)
Skin disorders		
Astringent	Root bark, gum	Nikkon et al. (2003); Bhattacharya et al. (1982)
Pyoderma	Chitin-binding protein from seeds	Mirella et al. (2011)
Rubefacient and vesicant	Gum, bark, root bark,	Bhattacharya et al. (1982); Prakash et al. (1987); Agrawal and Mehta (2008)
General disorders		
Antianaphylactic	Seeds	Mahajan and Mehta (2007)
Antiperoxidative	Hydroalcoholic extract of leaves	Nandave, Ojha, Joshi, Kumari, and Arya (2009)

(Continues)

TABLE 6 (Continued)

Traditional uses/ effects	Mode of action	References
Catarrh	Leaves and fruit extract	Fuglie (1999)
Low back/kidney pain	Roots	Agarwal and Mehta (2008)
Kidney stone cure	Root wood	Karadi, Gadge, Alagawadi, and Savadi (2006)
Kidney enlargement	Methanolic extract of leaves	Omodanisi, Aboua, and Oguntibeju (2017)
Splenomegaly	Roots, bark, leaves, flowers	Fuglie (1999); Prakash et al. (1987); Armand-Stussi et al. (2003); Yabesh et al., 2014
Anti-obese properties	Chlorogenic acid from leaves, ethanolic leaves extract	Panda et al. (2013); Metwally et al. (2017)
Weight gain	Benzylamine, Isothiocyanate extract of leaves	Iffiu-Soltesz et al. (2010); Waterman et al. (2015)
Earache	Gum, juice of root bark	Fuglie (1999); Prakash et al. (1987)
Migraine, headache	Alcoholic extract of leaf juice, gum with sesame oil	Upadhye, Rangari, and Mathur (2012); Bhattacharya et al. (1982)
Toothache	Roots decoction	Popoola and Obembe (2013)
Urinary bladder problems	Oral administration of seed powder	Gupta, Kannan, Sharma, and Flora (2005)

(Ruckmani, Kavimani, Anandan, & Jaykar B., 1998). The aglycone of deoxy-niazimicine isolated from root bark showed antifungal activity (Nikken et al., 2003). The stem bark possesses antifungal action (Bhatnagar, Santapau, Desai, Yellore, & Rao, 1961). Mo-CBP3 isolated from seeds showed in vitro antifungal effect on the phytopathogenic parasites *F. solani*, *F. oxysporum*, *Colletotrichum musae*, and *C. gloesporioides* at concentration of 0.05 mg/ml with 62% inhibition within 48 hr. They explained that Mo-CBP3 had antifungal action by reducing mycelia development and conidial feasibility. It is proposed that, through connections with the cell layer, Mo-CBP3 induced the creation of reactive oxygen species and caused cell death in the parasites (Batista et al., 2014). Nowadays, Mo-CBP3 is also considered as new antifungal drug in the development of transgenic crops for different traits viz., thermo-stability, broad antifungal spectrum, and low toxicity (Freire et al., 2015; Pinto et al., 2015).

### 3.4.3 | Antiviral activity

As a traditional medicinal plant, antiviral properties were observed in MO as reported in previous studies. Epstein-Barr infection activation in Raji cells could be inhibited by niaziminin isolated from foliages (Murakami et al., 1998; Sudha, Asdaq, Dhamingi, & Chandrakala, 2010). Ethanolic extract of leaves showed negative activity for herpes simplex virus (HSV; Lipipun et al., 2003) and seeds against HSV-1 (Ali, El-Taweel, & Ali, 2004). Ethanolic extract of leaves showed significant antiviral activity against infectious bursal disease virus (Ahmad et al., 2014), foot and mouth disease virus, equine herpes virus, hepatitis virus, and rhinovirus (Younus et al., 2016). Leaf aqueous extract activated the cellular immunity in infected mice with HSV-1 by decreasing the infection rate and reducing herpetic skin lesion growth (Kurokawa et al., 2016). Aqueous seed extract showed potential antiviral activity against Newcastle disease virus (Chollom et al., 2012). Any disease virus treatment with MO extract could defer skin injury development, extend the mean survival time, and reduce the mortality of HSV-1 infected mice (Khan, Ather, Thompson, & Gambari, 2005). Waiyaput et al. (2012) proposed that 80% ethanol extract of MO fruit demonstrated anti-Hepatitis B virus effect by reducing Hepatitis B virus replication with mellow cytotoxicity on HepG2 cells. MO was also used as a complement toward antiretroviral treatment in HIV infection (Monera & Maponga, 2010). MO leaves have generally been utilized in the treatment of HIV-related side effects, potentially in early strides in the infectivity of HIV-1 lentiviral particles in a viral vector-based screening (Nworu, Okoye, Ezeifeke, & Esimone, 2013). By assessing these researches discussed, it is concluded that MO could be used in the improvement of promising antiviral medications.

### 3.4.4 | Anticancer and antitumor activities

It is established fact that MO may be used as antineoproliferative mediator, by suppressing the development of cancer cells. Leaf extract exhibited chemoprotective (Anwar et al., 2007), cytotoxic (Berkovich et al., 2013; Nair & Varalakshmi, 2011), antihepatocarcinoma, antileukemia (Khalafalla et al., 2010), antimyelomic (Parvathy &

Umamaheshwari, 2007), and antiproliferative (Pamok et al., 2012) activities. These activities are because of 3-O-6x-oleoyl- $\beta$ -D-glucopyranosyl- $\beta$ -sitosterol,  $\beta$ -sitosterol-3-O- $\beta$ -D-glucopyranoside, 4-( $\alpha$ -L-rhamnosyloxy) phenylacetoneitrile, 4-hydroxyphenylacetoneitrile and 4-hydroxyphenyl-acetamide in seeds (Guevara et al., 1999); cis-9-hexadecenal, quinic acid, 3,5-dihydroxy-6-methyl-2,3-dihydro-4H-pyran-4-1, 9-octadecenamamide, methyl octadecenoate in flowers (Inbathamizh & Padmini, 2012) and quercetin, kaempferol (Krishnamurthy, Vardarajalu, Wadhvani, & Patel, 2015; Sreelatha, Jeyachitra, & Padma, 2011), 4-(4'-O-acetyl- $\alpha$ -L-rhamnosyloxy) benzyl isothiocyanate, O-ethyl-4-( $\alpha$ -L-rhamnosyloxy) benzyl carbamate, 4-(L-rhamnosyloxy) benzyl isothiocyanate, niaziminin, and niazimicin (Anwar et al., 2007; Murakami et al., 1998; Razis et al., 2014), D-allose (Sui et al., 2005) in leaves, eugenol (Al-Sharif, Remmal, & Aboussekhra, 2013), and isopropyl isothiocyanate (Matsuda, Ochi, Nagatomo, & Yoshikawa, 2007) in bark, palmitic acid in leaf, seed, and bark (Harada et al., 2001).

Previous studies showed that eugenol has a strong anticancer potential against melanoma (Pisano et al., 2007), osteosarcoma (Shin et al., 2007), acute lymphoblastic leukemia and hepatocellular carcinoma (Khalafalla et al., 2010), gastric cancer (Manikandan, Vinothini, Priyadarsini, Prathiba, & Nagini, 2011), skin tumor (Kaur, Athar, & Alam, 2010), mast cells (Park et al., 2005), and prostate cancer (Ghosh, Ganapathy, Alworth, Chan, & Kumar, 2009). Sharma, Paliwal, Janmeda, and Sharma (2012) reported that the enzyme, that is, glutathione and glutathione S-transferase (protection against carcinogens) activity loss was recovered by MO pod extract. Bharali et al. (2003) reported skin papillomagenesis subsequent to ingestion of seed and pod extracts. Previous studies had also shown the cytotoxicity of different extracts of MO on human cancer such as leukemia (Harada et al., 2001), pancreatic cancer (Berkovich et al., 2013), breast and colorectal cancer (Al-Asmari et al., 2015), colon cancer (Lea et al., 2012), KB tumor cell (Sreelatha et al., 2011), lung cancer (Jung et al., 2015), alveolar epithelial cancer (Tiloke et al., 2013), hepatocellular carcinoma (Khalafalla et al., 2010), and melanoma (Gismondi et al., 2013).

### 3.4.5 | Antidiabetic activity (Antihyperglycemic)

MO is outstanding for its therapeutic behavior and may utilized be for the traditional treatment of *Diabetes mellitus* (Konmy, Olounlade, Allou, Azando, & Hounzangbe-Adote, 2016; Leone et al., 2018). Antihyperglycemic (Divi, Bellamkonda, & Dasireddy, 2012; Jaiswal, Rai, Kumar, Mehta, & Watal, 2009) and hypoglycemic (Ajit, Choudhary, & Bandyopadhyay, 2002; Tende, Ezekiel, & Dikko, 2011) activities of leaves might be presumably because of the presence of terpenoids and flavonoids, which involved in stimulation of  $\beta$  cells and resulting discharge of insulin. Glucomoringin (glucosinolates), quercetin and kaempferol (flavonoids), and aschlorogenic acid showed hypoglycemic properties (Sayed, 2012). Ethanolic extract MO leaves possesses hypoglycemic activity in STZ-induced diabetic Wistar rats, which is due to lowering blood glucose level (Tende et al., 2011). Methanol extract of dried fruit powder accelerates the insulin release significantly due to benzyl, benzyl nitriles, and N-benzyl thiocarbamates (Francis, Jayaprakasam, Olson, & Nair, 2004). An

examination revealed that benzylamine isolated from MO leaves decreased the plasma cholesterol, body weight gain, hyperglycemic reactions, and fasting blood glucose dimensions of high-fat diet-induced mice (Iffiu-Soltesz et al., 2010). Leaf extract reduced blood glycosylated hemoglobin and cholesterol level in Type 2 diabetes (Ghiridhari, Malhati, & Geetha, 2011; Nambiar, Guin, Parnami, & Daniel, 2010). Leaf powder capsules (4 g) considerably improved the discharge of insulin in 10 healthy human volunteers (Anthanont, Lumlerdkij, Akarasereenont, Vannasaeng, & Sriwijitkamol, 2016). 4-[( $\alpha$ -L-rhamnosyloxy)benzyl] isothiocyanate and 4-[(4'-O-acetyl- $\alpha$ -L-rhamnosyloxy)benzyl] isothiocyanate showed antidiabetic activity (Tumer et al., 2015; Waterman et al., 2015). In alloxan-induced diabetic rats, the aqueous extract (300 mg/kg) exhibited 44.06% reduction respectively of blood glucose concentration within 6 hr of administration, whereas almost as effective as the standard drug tolbutamide (200 mg/kg.) caused 46.75% reduction (Edoga, Njoku, Amadi, & Okeke, 2013). A watery concentrate of leaves accelerated insulin level in diabetic rodents (Tuorkey, 2016). A protein had antigenic epitopes like insulin reported in seed coat and leaves of MO and showed hypoglycemic properties on oral administration. It also cross-reacted with anti-insulin antibodies, which proved that it might have antigenic epitopes similar to insulin (Paula et al., 2017). Leaf methanolic extract indicated defensive activity against diabetic-prompted renal damage, receptive oxygen species, and irritation (Omodanisi et al., 2017). Thus, majority of studies have used leaf extracts on animal subjects like rodents. However, as regards to the antidiabetic activity, a recent study showed that the administration of a dose of 20 g/day of *M. oleifera* leaves determined a reduction of the postprandial glycemic response in human subject (Leone et al., 2018) that was maintained for up to 3 hr from the beginning of the meal blended with MO leaf powder.

### 3.4.6 | Antioxidant activity

MO is well-known tree for its high antioxidant in leaves, fruits, and seeds among other fruits and vegetables (Yang et al., 2006). MO tree contains about 40 natural antioxidants (Mahmood et al., 2010). The antioxidant activity is attributed due to the presence of various types of antioxidant compounds (Omodanisi et al., 2017; Siddhuraju & Becker, 2003) such as ascorbic acid,  $\beta$ -carotene (Kumar, Pandey, Mohan, & Singh, 2012; Mahajan & Mehta, 2007), quercetin, kaempferol (Gupta et al., 2012), flavonoids, phenolics (Anwar, Ashraf, & Bhangar, 2005; Sreelatha et al., 2011), isothiocyanates, polyphenols, and rutin in leaves (Tumer et al., 2015); myricetin (Lalas & Tsaknis, 2002b), tocopherols and lectins in seeds (Singh et al., 2013); procyanidins in stem and root bark (Atawodi et al., 2010) and palmitic acid, phytosterols, and 9-octadecenamamide in flowers (Inbathamizh & Padmini, 2012), monopalmitic acid, oleic acid, tri-oleic triglycerides in seed oil (Lalas & Tsaknis, 2002a; Mahajan & Mehta, 2007). Methanol and acetone extracts of leaves (Atawodi et al., 2010; Charoensin, 2014), aqueous extracts of roots (Satish, Kumar, Rakshith, Satish, & Ahmed, 2013), methanolic extracts of stem and pods (Gupta et al., 2012; Kumbhare, Guleha, & Sivakumar, 2012) showed wide antioxidant profile. The



highest antioxidant activity is recorded for isoquercetin as it increased the mRNA articulation dimensions of catalase, oxygenase, and superoxide dismutase (Vongsak, Mangmool, & Gritsanapan, 2015).

Folate isolated from foliage is the notable among the most imperative water-soluble vitamins, assumes a basic role in different cell metabolisms (Scotti, Stella, Shearer, & Stover, 2013). Antioxidant benefits could be obtained for chicken sausages in cold storage through the application of dry leaf powder by reducing lipid oxidation (Jayawardana, Liyanage, Lalantha, Iddamalagoda, & Weththasinghe, 2015). Quercetin isolated from essential oil is quite compelling and may inhibit tumor necrosis factor by Kupffer cells (Baghel, Shrirastava, Baghel, Agrawal, & Rajput, 2012). Luteolin has additionally a solid antioxidant agent and may show evidence of defensive ability on DNA (Romanova, Vachalkova, Cipak, Ovesna, & Rauko, 2001). Further studies revealed that luteolin is fit for applying a variety of biological and pharmacological activities, namely, antioxidant, anti-inflammatory, antimicrobial, anticancer, antiallergic, antiplatelet, and other activities (El-Hawary, El-Sofany, Abdel-Monem, Ashour, & Sleem, 2012). The hydrocarbons of seed essential oil demonstrated radical scavenging effect (Yassa, Masoomi, Rankouhi, & Hadjiakhoondi, 2009). The application of leaf extract maintained plasma malondialdehyde (MDA) level and ferric reducing ability of plasma without hypoglycemia effect in human volunteers (Ngamukote et al., 2016).

### 3.4.7 | Anti-inflammatory activity

This is an established fact that leaf, fruit, seed, and root extracts have been used since long back in enhancement of inflammation related disorders, for example, asthma, allergic rhinitis, atopic dermatitis, and rheumatoid arthritis (Chumark et al., 2008; Hamza, 2010; Lee et al., 2013; Muangnoi, Chingsuwanrote, Praengamthanachoti, Svasti, & Tuntipopipat, 2012). Anti-inflammatory activities are possibly attributed due to the presence of 4-[( $\alpha$ -L-rhamnosyloxy) benzyl] isothiocyanate, 4-[(4'-O-acetyl- $\alpha$ -L-rhamnosyloxy) benzyl]isothiocyanate (Stohs & Hartman, 2015), quercetin, quercetin-3-O-glucoside, kaempferol glucosides, (Coppin et al., 2013), 4-(2-O-acetyl- $\alpha$ -L-rhamnosyloxy) benzyl isothiocyanate, 4-(3-O-acetyl- $\alpha$ -L-rhamnosyloxy)benzyl isothiocyanate (Cheenpracha et al., 2010), 3,5-dihydroxy-6-methyl-2,3-dihydro-4H-pyran-4-1, 9-octadecenamamide (Inbathamizh & Padmini, 2012), crypto chlorogenic acid (Vongsak, Gritsanapan, Wongkrajang, & Jantan, 2013), aurantiamide acetate, 1,3-dibenzyl urea (Maheshwari, Yadav, Malhotra, Dhawan, & Mohan, 2014; Pandey et al., 2012) and 4( $\alpha$ -L-rhamnosyloxy)-benzyl glucosinolate (Galuppo et al., 2014).

Leaves ethanolic extract demonstrated anti-inflammatory properties by inhibiting chemotactic oxidation of polymorphonuclear leukocytes (Vongsak et al., 2013), keratinocytes (Choi et al., 2016) and leaves, seeds, and roots extracts against multiple sclerosis cascades (Galuppo et al., 2014). The hydroalcoholic extract of seeds additionally reduced the action of TNF- $\alpha$ , IL-4, IL-6, and myeloperoxidase, a cause of inflammatory bowel diseases (Minaiyan et al., 2014). Hydroalcoholic leaf extract enhanced the cellular and humoral immunity in immunosuppressed rodents (Nfambi, Bbosa, Sembajwe, Gakunga, & Kasolo, 2015). Methanol extract decreased the edematogenic impact of

carrageenan- and histamine-induced foot edema in laboratory animals (Adedapo, Falayi, & Oyagbemi, 2015) and also pain relieving impacts by reducing mechanical allodynia and thermal hyperalgesia in Freund's adjuvant joint inflammation prompted rodents, whereas methanolic root decoction decreased thermal hyperalgesia (Manajeji, Jafari, Zaringhalam, Rezazadeh, & Taghizadfarid, 2011). The hydroethanolic extract of flowers also decreased the movement of incendiary mediators and proinflammatory cytokines, for example, PGE<sub>2</sub>, IL-1 $\alpha$ , IL-6, TNF- $\alpha$ , NF- $\kappa$ B, iNOS, NO, and COX2 in LPS prompted RAW264.7 macrophage cells (Tan, Arulselvan, Karthivashan, & F akurazi S., 2015). The fruit demonstrated the highest activity in reducing NO release induced by LPS in RAW264.7 cells (Lee et al., 2013). The fruit extract of MO blocked the atomic translocation of NF- $\kappa$ B and increased inhibitor  $\kappa$ B articulation (Arulselvan et al., 2016). MO roots showed evidence of restraint against IL-2 activity (Sashidara et al., 2009). Fermentation of MO decreased hepatic adiposity, endoplasmic reticulum anxiety and improved resistance to glucose in obese mice (Joung et al., 2017).

### 3.4.8 | Antiparasitic activity

MO possesses different activities such as antiulcerogenic by leaf aqueous extract (Dahiru, Onubiyi, & Umaru, 2006); schizonticidal by leaf acetone extract (Patel, Gami, & Patel, 2010), larvicidal by seed coagulant lectin protein (De Oliveira et al., 2011; Prabhu, Murugan, Nareshkumar, Ramasubramanian, & Bragadeeswaran, 2011), antiulcerative colitis by root extract (Gholap, Nirmal, Pattan, Pal, & Mandal, 2012), antiurolithiatic by root wood extract (Karadi et al., 2006), and antimacrofilaricidal activities by gum exudates (Kushwaha et al., 2011). The antilarvicidal activity for *Aedes aegypt*, a vector of dengue virus, was reported in flower extract through trypsin inhibitor activity (Pontual et al., 2014). The growth of fourth-stage *Aedes aegypti* larvae was interrupted by lectin protein isolated from seeds (Agra-Neto et al., 2014). The growth of *Biomphalari aglabrata* embryos was delayed in aqueous extract of flowers and increased also the mortality of young snails (Rocha-Filho et al., 2015). Antileishmanial activity for *Leishmania donovani* promastigotes was reported by Kaur, Kaur, Singh, and Singh (2014) with the application of alcoholic extracts of leaves and roots. Niazinin isolated from leaf ethyl acetate extract showed the most antileishmanial activity (Abd Rani, Husain, & Kumolosasi, 2018). Singh, Paul, De, and Chakraborti (2015) observed that various parts, that is, leaf, flower, bark, roots, stem wood exhibited antileishmanial activities, of which ethyl acetate extract of flower had the maximum negative impact on *L. donovani*. Sengupta et al. (2012) demonstrated that seed extract could be useful to check the growth of helminth eggs and to remove turbidity from irrigation water. The inhibitory effect on egg hatching of *Haemonchus contortus* was reported by aqueous extract of seeds (Salles et al., 2014). Similarly, seed aqueous extract showed larvicidal, pupicidal, and in addition adult mosquito killing activities for *Culex quinquefasciatus* (Ashfaq & Ashfaq, 2012). Antitrypanosomal activity against *Trypanosoma brucei* was reported in petroleum ether, chloroform, methanol, and aqueous extracts of leaves, stem bark, and roots (Ibrahim, Mohammed, Isah, & Aliyu, 2014).

### 3.4.9 | Cardio protective activities

Every part of MO is used for cardiovascular, circulatory stimulant, and lowering cholesterol and blood pressure level. Leaf extract showed hypocholesterolemic activities (Atsukwei, Eze, Adams, Adinoyi, & Ukpabi, 2014; Ghasi, Nwobodo, & Ofili, 2000; Marcela, Almatrafi, & Fernandez, 2017; Okwari, Dasofunjo, Asuk, Alagwu, & Mokwe, 2013). Moringinine is utilized as a cardiac tonic, which acts on sympathetic nervous system (Anwar et al., 2007). N- $\alpha$ -L-rhamnopyranosyl vincosamide in bark is showed cardioprotective effects (Morton, 1991). Leaf extract decreased mortality and morbidity of cells in coronary heart disease, and this may be due to the gossypetin, quercetagenin, and proanthocyanidins (Kumar et al., 2012; Seshadri & Nambiar, 2003).  $\beta$ -sitosterol had significant cholesterol reducing property (Ghasi et al., 2000). Cardioprotective activity of leaf extract is assessed in rats (Nandave et al., 2009). Lipid profile of liver, heart, and aorta in hypercholesteremic rabbits was decreased by MO fruits and increased the discharge of fecal cholesterol because of the having lower cholesterol, phospholipids, triglycerides, lipoprotein (Mehta et al., 2003). Leaf ingredients used to check blood glycated hemoglobin, total cholesterol, non-HDL-C, HDL-C, VLDL-C, and LDL-C in Type 2 diabetic patients (Nambiar et al., 2010).

Various leaf ingredients such as niazinin, niazimicin, niaziminin, niazimin, niazirin, niazicin, niazirinin, 4-(4'-O-acetyl- $\alpha$ -rhamnosyloxy) benzyl isothiocyanate are responsible for hypotensive and bradycardiac activities (Bose, 2007), and glucomoringine is reported to have hypertensive effect (Dangi, Jolly, & Narayana, 2002; Maheshwari et al., 2014). Thiocarbamate and isothiocyanate glycosides (Faizi et al., 1995), methyl phydroxybenzoate and  $\beta$ -sitosterol (Faizi et al., 1998) isolated from pods are known for hypotensive activity. Seeds demonstrated vascular oxidative and nitrosative stresses in spontaneously hypertensive rats (Randriamboavonjy, Rio, Pacaud, Loirand, & Tesse, 2017). Niazinin-A, niazinin-B, and niazimicin found in ethanolic extract of leaves are effective in controlling the blood pressure level possibly through a calcium antagonist effect (Anwar et al., 2007; Dubey, Dora, Kumar, & Gulsan, 2013). Similarly, blood pressure lowering effect was reported by ethanol and aqueous extract of MO pods (Faizi et al., 1998). Blood pressure is also controlled by the application of MO leaves due to the presence of nitrile, glycosides, and thiocarbamate glycosides (Dangi et al., 2002; Marcela et al., 2017). Extracts of seeds and leaves had blood pressure reducing effect because of alkaloids and flavonoids through the inhibitory impact on angiotensin converting enzymes (Abdulazeez, Ajiboye, Wudil, & Abubakar, 2016). Dried seeds had the evidence of cardioprotective effect, induced cardiovascular diastolic function, and reduced nighttime pulse rate without adjusting the circulatory strain in hypertensive rats (Randriamboavonjy et al., 2016). Butanolic extract of leaf decreased the possibility of cardiac necrosis and oxidative stress in isoproterenol-induced rodents (Panda, 2015).

### 3.4.10 | Metabolic disorders

Different leaf extracts showed antihyperlipidemic effect (Chumark et al., 2008; Divi et al., 2012; Irfan, Asmawi, Khan, & Sadikun, 2016;

Jain, Patil, Haswani, Girase, & Surana, 2010) and hepatoprotective activities (Efiong, Igile, Mgbeje, Out, & Ebong, 2013; Pari & Kumar, 2002). Ethanolic extract of foliages, leaf oil, and seeds had hepatoprotective and kidney defensive properties against  $\gamma$ -radiation, HgCl<sub>2</sub>, acetaminophen, and arsenic (Abarikwu, Benjamin, Ebah, Obilor, & Agbam, 2017; Gupta, Dubey, Kannan, & Flora, 2007; Sinha, Das, Bhattacharjee, Majumdar, & Dey, 2011). Afterward, Tahiliani and Kar (2000), Anwar et al. (2007), and Chumark et al. (2008) confirmed the antihyperthyroidism and atherosclerotic properties for leaf extracts. Additionally, aqueous extract of roots have been found to have estrogenic and progestational activity (Shukla et al., 1988) along with liver and renal defensive property (Mazumder, Gupta, Chakrabarti, & Pal, 1999). Roots extract has been reported to possess antispasmodic activity (Caceres et al., 1992; Dangi et al., 2002). Hydroethanolic extract of leaves reduced mRNA articulation, which is liable for keeping up lipid homeostasis (Sangkitikomol, Rocejanasaroj, & Tencomnao, 2014). Leaf aqueous extract inhibited formation of nonfluorescent and fluorescent advanced glycation end products (Nunthanawanich et al., 2016). The pods with high fiber content used to treat stomach-related issues and intestinal problems (Oduro, Ellis, & Owusu, 2008). Seed powder is used to reduce the arsenic from blood, liver, and kidneys (Gupta et al., 2005).

Moringine and moringinine had antihypoglycemic properties (Maheshwari et al., 2014; Mehta et al., 2011). Phytochemical examination demonstrated that  $\beta$ -sitosterol and 4-[ $\alpha$ -(L-rhamnosyloxy) benzyl]-o-methyl thiocarbamate (trans) are powerful agent to reduce the cholesterol level, antidiarrheal, antispasmodic, and hypolipidemic activities (Anwar et al., 2007; Jain et al., 2010; Maheshwari et al., 2014), whereas quercetin-3-glycoside is responsible for antidiyslipidemic (Mbikay, 2012) and hypoglycemic (Maheshwari et al., 2014) properties. The quercetin and kaempferol in flowers and pods showed hepatoprotective exercises (Farooq et al., 2012; Gupta et al., 2012). The extract of leaves and fruits has ability to heal chronic gastric ulcers induced via acetic acid (Devaraj & Gopala-Krishna, 2013). Niaziridin improved the gastrointestinal retention of nutrients and supplements (Stohs & Hartman, 2015). Leaf powder capsules (4 g) induced the discharge of insulin (Anthanont et al., 2016). The action of CD69, INF-g, and CD44 protein is increased by oral administration of leaf aqueous extract, which are responsible for hypoglycemic activity and furthermore decreased creatinine and urea levels from damaged kidneys (Paula et al., 2017). The reduction in lipid peroxidation and immunoglobulin IgG and IgA was observed for seed extract (Al-Malki & El Rabey, 2015). Antihypoglacemia (antidiabetic) effect was exhibited by vacuum dried 95% ethanolic extract of stem bark (Kar, Choudhary, & Bandyopadhyay, 2003) and ethanolic extract (Irfan et al., 2016).

### 3.4.11 | Nervous system disorders

Immunosuppressive and immunostimulatory activities were reported for MO leaf extract (Rachmawati & Rifa, 2014). The immunomodulatory impact of leaves extract is mediated through reduction in cyclophosphamide incited immunosuppression by invigorating cell and humoral immunity (Gupta et al., 2010), which is attributed to the presence of compounds like isothiocyanates and glycoside

cyanides (Sudha et al., 2010). Leaves had neuroprotective impact by progressing neuronal survival and neurite outgrowth (Hannan et al., 2014). Leaf extract had defensive impact against Alzheimer's disease by adjusting cerebrum monoamine levels and electrical impulses (Ganguly & Guha, 2008). Further, it was discovered that the CNS depressant action is mediated through its pain relieving and anticonvulsive properties because of triterpenoid, saponins, and flavonoids in leaves (Bakre, Aderibigbe, & Ademowo, 2013; Gupta, Mazumder, & Chakrabarti, 1999; Ray, Hazra, & Guha, 2003). MO has used to improve memory by nootropics movement and protect against the oxidative pressure in Alzheimer's illness (Ganguly, Hazra, Ray, & Guha, 2005). A model was proposed by Ganguly and Guha (2008) for Alzheimer's disease including the infusion of colchicine into the brain of rodents. They showed that MO induced the modification of cerebrum monoamines and electrical patterns. Protective effect against neurodegeneration has been reported by Giacoppo et al. (2015) due to the glucosinolates (R,S-Sulforaphane-SFN), which offer protection to mesencephalic dopaminergic neurons from cytotoxicity and oxidative stress through the prevention of reactive oxygen, DNA crumbling, and membrane rupture (Han et al., 2007).

### 3.4.12 | Wound healing activity

Aqueous leaves extract decreased scar regions and accelerated the healing rates of wounds in addition to increase the granuloma and skin breaking strength in albino rats (Lambale & Kumar, 2012; Rathi, Bodhankar, & Baheti, 2006). MO phytochemical ingredients mediate the wound healing activity by suppressing antihealing agents. Wound healing constituents, for example, quercetin, kaempferol, phytosterols (Hukkeri et al., 2006), and vicenin-2 (Muhammad, Pauzi, Arulselvan, Abas, & Fakurazi, 2013) have been reported in ethyl acetate extract of leaves and dexamethasone in bark (Lambale and Kumar et al., 2012). Proteolytic, fibrinolytic, and fibrinogenolytic actions on blood coagulating process by protease activity have been assessed for leaf and root aqueous extracts (Satish, Sairam, Ahmed, & Urooj, 2012). The protease also demonstrated similar activity to plasmin and thrombin coagulating protein and also attenuated apoptosis in addition to suppressing the peroxisome proliferator-activated receptor gamma activation and shortened the time taken to activate prothrombin and thromboplastin. The wound healing activity has also been reported in the seeds of MO (Bhatnagar, Parwani, Sharma, Ganguli, & Bhatnagar, 2013; Parwani, Bhatnagar, Bhatnagar, Sharma, & Sharma, 2016).

## 3.5 | Miscellaneous uses

### 3.5.1 | *M. oleifera* for agriculture and animal husbandry

Extracts of MO leaves in 80% ethanol having cytokinin type plant growth regulator, which may be used as foliar spray to quicken growth in peanut, soya bean, black bean, maize, onion, sorghum, tomato, coffee, and sugarcane crops. This makes crop plant resistant to insect-

pest and increase the biological yield (Foidl, Makkar, & Becker, 2001). The sole crop of maize and sweet potato were compared with the maize with moringa and sweet potato with moringa. Intercropping with moringa produced the highest crop growth than the monoculture cropping pattern. The results also indicated a decrease in soil acidity from 1.86 to 1.60 (Abusuwar & Abohassan, 2017). Yield can be increased by 25–30% for nearly any group of plants (maize, black pepper, tomato, soya bean, onion, sorghum, tea, coffee, melon, and chili) with application of juice of fresh MO leaves. One of the dynamic constituent, that is, zeatin, acts as a characteristic plant development enhancer and helps to increase crop yield (Leone, Spada, et al., 2015). Leaf extract may be used to boost up the *Rhizobium* root nodulation and nitrogenase activity in *Vigna mungo* when applied over seeds or as a root dressing (Bandana, Srivastava, & Mathur, 1987). The oilcake of MO is used as a fertilizer (Dastur, 1964). The flowers are a unique source of pollens for honey bees (Rajan, 1986; Sohani, 2018). The wood is little utilized outside of its local range with the exception of as a fuelwood and also for light construction (Little & Wadsworth, 1964). Its wood provides a fairly good fuel for cooking even being light in weight (Zheng, Zhang, & Wu, 2016).

The tender twigs are being utilized as feed for cows, sheep, goats, and camels worldwide (Mahatab, Ali, & Asaduzzaman A.H.M., 1987). Presence of high protein and low content of antinutritional compounds makes it substitute to regular cattle feeds, that is, alfalfa and oil seed cakes (Babiker, Al-Juhaimi, Ghafoor, & Abdoun, 2016). Feeding of MO leaf helps to improve the milk quality and quantity (Mendieta, Sporndly, Reyes-Sanchez, & Sporndly, 2011), meat production (Babiker et al., 2016), beef production, and dietetic nature of roughages (Roy, Bashar, Hossain, Huque, & Makkar, 2016). However, the gas production increased with a normal feed of MO leaves and methane content of the gas was 81% (Essien, Essien, & Eluagu, 2016). The MO has also been reported as an effective feedstock for biofuels, for example, biogas and biodiesel (Kafuku & Mbarawa, 2010; Rashid, Anwar, Moser, & Knothe, 2008). Pontual et al. (2012) assessed the caseinolytic and milk coagulating properties of proteases in flowers; further, milk clotting enzymes were isolated and purified from MO seeds by Ahmed (2016). It was effectively used in cheese production, therefore recommending an option in contrast to conventional animal rennets.

### 3.5.2 | *M. oleifera* as coagulant

There are several genuine drawbacks of using synthetics and polymers due to the harmful consequences for human well-being and nature. In this way, a restoration of enthusiasm for natural coagulants has risen as a result of the high cost, health issues, and nonbiodegradable nature (Choy, Prasad, Wu, Raghunandan, & Ramanan, 2014). Seeds of MO have used to treat the waste water as they contain cationic polyelectrolyte, which neutralizes the negatively charged colloids present in dirty and muddy water (Makkar & Becker, 1996). Approximately, 300 mg of MO seed powder is sufficient to purified 1 L of water. In this condition, microorganisms can be removed by settling with the pattern of removal of colloids from coagulated and

flocculated water (Casey, 1997). Various coagulating and flocculating components were separated from various plant parts. Okuda, Baes, Nishijima, and Okada (2001) identified a coagulating agent from the salt solution of seeds extract, whereas Santos et al. (2009) purified hemagglutinating proteins and lectin coagulant from MO seeds and other tissue extracts. These coagulants were evaluated with aluminum sulphate to check its efficacy for turbid water treatment (Pritchard, Craven, Mkandawire, Edmondson, & O'Neill, 2010) along with the flower extract for purification of the waste water (Moura et al., 2011). Further, Broin et al. (2002) cloned cDNA encoding flocculating proteins from MO seeds and used in purification of water against clay particles, gram positive, and negative bacteria. Seeds are outstanding coagulant agents among other characteristic coagulants (Ferreira et al., 2008). Cationic proteins (MO2.1 and MO2.2 also referred to as Flo; P24303, Swissprot, *Moringa oleifera* cationic protein (MOCP)) and chitin-binding protein isoform (Mo-CBP3) isolated from seeds are being used for flocculating activities (Freire et al., 2015; Ullah et al., 2015). The coagulation efficacy of MOCP is observed to be increased with an increase in the initial turbidity of water (Katayon et al., 2006). Owing to amphiphilic properties, MOCP integrate into the bacterial membranes and are specifically kill several microorganisms, including waterborne pathogens (Shebek et al., 2015). The seed contain polypeptides, a coagulant, used to treat river water with suspended solid particles and groundwater (Lijesh & Malhotra, 2016).

The dried or powdered seeds have been exploited directly or in petroleum ether extract form (Johri, Reeta, & Johri, 2004), as an efficient and low-cost coagulant to remove turbidity and decrease the microbial contamination from drinking water from village communities in the Sudan, Malawi, India, Myanmar, and Indonesia (Mandloi, Chaudhari, & Folkard, 2004; Nyein et al., 1997). Crude extract of MO seeds was used by rural women to treat the very turbid Nile stream water instead of alum causing gastrointestinal disturbances and Alzheimer's illness in Sudan (Ravikumar & Sheeja, 2013). Turbidity removal effectiveness was resulted 80–99% at a pH of 7–7.5 by using MOCP. Turbidity reduction of 85–98% paralleled by a primary *E. coli* decrease of 99.2–99.97% was attained within the first 1–2 hr of treatment (Bina et al., 2010). Thus, MOCP as coagulant may be used for drinking water treatment without any risk.

### 3.5.3 | *M. oleifera* seeds as biosorbent

A billion people across the Asia, Africa, and Latin America are estimated to depend on untreated surface water sources for their day to day water needs. Of these, approximately two million are thought to die by using polluted water each year, with the larger part of these deaths happening among infants (Mahamood et al., 2010). There are several studies on MO for biosorption properties and removal of different heavy metals from water. The row husk as biosorbent does not require any kind of handling, and it permits valorization of the waste produced after coagulant extraction from the seed. The biosorption of all metals on the adsorbent was rapid as over 80% of the metals were removed within the first 20 min to 1 hr of application (Adhiambo, Lusweti, & Morang, 2015). Effective removal of azo and

anthroquinonic dyes, and organic compounds viz., benzene, toluene, ethylbenzene, and cumene using seed extracts and fruits, respectively has also been demonstrated (Akhtar, Hasany, Bhangar, & Iqbal, 2007; Beltran & Sanchez, 2008). Different parts of *M. oleifera* viz., seed powder for  $\text{Cd}^{+2}$  (Kituyi et al., 2013; Mataka, Sajidu, Masamba, & Mwatseteza, 2010; Sharma, Kumari, Srivastava, & Srivastava, 2006),  $\text{Pb}^{+2}$  and  $\text{Cr}^{+2}$  (Adhiambo et al., 2015);  $\text{Zn}^{+2}$  (Kituyi et al., 2013),  $\text{As}^{+3}$  and  $\text{As}^{+5}$  (Kumari, Sharma, Srivastava, & Srivastava, 2006); seeds for  $\text{Ag}^{+1}$  and  $\text{Co}^{+2}$  (Araujo et al., 2010); activated carbon from husk and pods for  $\text{Pb}^{+2}$  (Nadeem et al., 2006); activated carbon from wood for  $\text{Cu}^{+2}$ ,  $\text{Ni}^{+2}$ , and  $\text{Zn}^{+2}$  (Kalavathy & Miranda, 2010); activated carbon from leaves for  $\text{Cd}^{+2}$ ,  $\text{Cu}^{+2}$ , and  $\text{Ni}^{+2}$  (Reddy, Seshaiyah, Reddy, & Lee, 2012); biomass for  $\text{Zn}^{+2}$  (Bhatti, Mumtaz, Hanif, & Nadeem, 2007); bark for  $\text{Pb}^{+2}$  (Reddy, Seshaiyah, Reddy, Rao, & Wang, 2010); leaves for  $\text{Pb}^{+2}$  (Reddy, Harinath, Seshaiyah, & Reddy, 2010); husk  $\text{Cu}^{+2}$  and  $\text{Cd}^{+2}$  (Garcia-Fayos, Arnal, Piris, & Sancho, 2016) have been tested as biosorbents agents. However, seeds of different *Moringa* species could also be used as a low-cost biosorbent for removal of heavy metals, that is,  $\text{Cd}^{+2}$ ,  $\text{Cr}^{+2}$ , and  $\text{Ni}^{+2}$  from aqueous media (Mataka et al., 2010; Sharma, Kumari, Srivastava, & Srivastava, 2007).

### 3.5.4 | *M. oleifera* as biodiesel

In the last decade, biodiesel production from unconventional oils yielding plants, such as, *Jatropha*, *Moringa*, *Pongamia*, and tobacco have acknowledged more attention (Kivevele & Huan, 2015; Rashid et al., 2008). Azam, Waris, and Nahar (2005) concluded in their study led on 75 Indian plant species, which have more than 30% fixed oil in their seed/kernel and unsaturated fatty acid methyl esters of MO seed oil meet all the principle determinations of biodiesel standard of Germany, Europe, and the United States. MO seeds contain 33–41% (w/w) oil on account of the behenic acid (docosanoic acid and 7% w/w), which has significant protection from oxidative degradation (Rashid et al., 2008). On account of the significant of high amount of monounsaturated unsaturated fats as oleic acid (72.2%), MO seed oil is a potential substitute for biodiesel production (Sohani, 2018). The fuel properties of MO are predominantly dependent on the unsaturated fat composition, wherein  $\text{C}_{16}$  and  $\text{C}_{18}$  monounsaturated fatty acid methyl esters are ideal components to accomplish an adequate balance between oxidative stability and cold flow properties of biodiesel (Knothe, 2008). Azad, Rasul, Khan, Sharma, and Islam (2015) proposed the possibility of MO seed oil as a supportable biodiesel fuel and concluded, MO is one of the forthcoming industrial crops for biodiesel generation in calm atmosphere of Austria.

### 3.5.5 | Industrial uses

*M. oleifera* seed oil known as "ben oil" is commonly used for the preparation of cosmetics and blending in perfumery materials (Arise et al., 2014). The oil is commonly consumed by perfumers for its capacity of engrossing and holding smells (Ramachandran et al., 1980). Seed oil is used for different purposes, that is, lighting, hair dressing, and watches oil (Morton, 1991). Seed oil is also used by soaps making industries

(Rashid et al., 2008). Seed oil was used as grease oil for hardwires and watches (Rashid et al., 2008). Cognis Laboratories Serobiol Ogiques group formulated Puricare TM and Purisoft TM, two dynamic ingredients based on natural peptides of MO seeds, which offer protection against pollution effects on hairs and skin (Shah et al., 2016). The salt extract of seeds has also been used with povidone (polyvinylpyrrolidone) to enhance solubility of the partially soluble Non-Steroidal Anti-Inflammatory Drugs (NSAIDS) (Noolkar, Jadhav, Bhende, & Killedar, 2013). A study conducted by Torondel, Opare, Brandberg, Cobb, and Cairncross (2014) concluded the viability of dried and wet leaf powder as a hand-washing item and indicated dimensions of inhibition of *E. coli* similar to a nonmedicated liquid soap because of saponin content. Recently, MO leaf extract in blended with gelatin isolated from puffer fish skin was used for production of antimicrobial, antioxidative, and biodegradable packaging materials (Lee, Yang, & Song, 2016). The gum exudate of MO is used for blue dye in Jamaica (Verma, Banerji, Misra, & Nigam, 1976). Its adhesive gum is utilized in leather tanning and calico printing (Morton, 1991; Sohani, 2018). Gum also be utilized for production of mucoadhesive polymer, disintegrant, binder (Patel, Patel, & Upadhyay, 2012), and a competent amorphous state stabilizer in drug formulations, that is, ibuprofen, meloxicam, and felodipine (Bhende & Jadhav, 2012). Apart from gum, its viscose resin is used textile industries (Fuglie, 1999). A coarse fiber is produced by its corky bark, which is used in paper making mats and cordage. In India, wood is utilized to a limited extend in the textile industry for transports and picking-sticks and is suitable for pulpwood alternative for newsprint (Singh, Wadhvani, & Johri, 1983), wrapping, textiles (Ganatra et al., 2012), cellophane, and materials (Nautiyal & Venhataraman, 1987).

#### 4 | CONCLUSIONS AND FUTURE PERSPECTIVES

The *M. oleifera* is the most inexpensive and credible alternative to provide good nutrition and curing and prevention of several disorders. The secondary metabolites and higher amount of phytochemicals with other essential components could be valuable for food fortification and to ensure the practice of traditional medicines. Therefore, the poor nations should promote planting and uses of *M. oleifera* instead of bounties for food relief from the rich nations. Previous studies have evaluated the traditional uses of *Moringa* species, and the majority of these examinations supported the traditional claims. However, there are lots of uses that have not been assessed till date, particularly in species other than MO. Because MO normally grows in different environments, a variation in phytochemical ingredients and nutraceutical status is reported from different parts. In any case, the degree to which the chemical composition fluctuates in seed sources adapted to different habitats is still unknown. In spite of the fact that *Moringa* leaves are viewed as great protein source, despite everything, it must be studied that which protein contend with the more common protein in milk producing ruminants. Few studies are carried out on human subjects, whereas the dominant parts of studies have used aqueous

and alcoholic extractions basically in rodents and other animals. Keeping in view the importance of MO in humans as a "Nutritional Dynamite," clinical trials to assess the efficacy of *Moringa* with human subjects and positive effects on human immune system in fighting diseases should be taken to explore before promoting it and using it for treating malnutrition and various disease in low income countries. Likewise, few bioactivity-based extraction techniques have been used to decide the connections between extraction systems and solvents. It is not clear about the degree of different ingredients of MO interrelate through additive, synergistic, and/or inhibitory impacts in therapeutic treatment. Assessment of the safety and toxicity of antimicrobial agents from MO is important before implementing the uses of these compounds. Numerous studies have also been demonstrated the *Moringa* seeds as coagulant agent for treating waste water. Subsequently, it is essential to recognize the dynamic constituents of MO seed to understand its coagulation mechanism.

#### CONFLICT OF INTEREST

The authors have declared that no competing interests exist.

#### ORCID

Ashok K. Dhakad  <https://orcid.org/0000-0001-5241-8036>

#### REFERENCES

- Abarikwu, S. O., Benjamin, S., Ebah, S. G., Obilor, G., & Agbam, G. (2017). Protective effect of *Moringa oleifera* oil against HgCl<sub>2</sub> induced hepatoand nephro-toxicity in rats. *Journal of Basic Clinical Physiology and Pharmacology*, 28, 337–345.
- Abd El Baky, H. H., & El-Baroty, G. S. (2013). Characterization of Egyptian *Moringa peregrina* seed oil and its bioactivities. *International Journal of Management Science and Business Research*, 2, 98–108.
- Abd Rani, N. Z., Husain, K., & Kumolosasi, E. (2018). *Moringa* Genus: A review of phytochemistry and pharmacology. *Frontiers in Pharmacology*, 9, 1–26.
- Abdulazeez, A. M., Ajiboye, O. S., Wudil, A. M., & Abubakar, H. (2016). Partial purification and characterization of angiotensin converting enzyme inhibitory alkaloids and flavonoids from the leaves and seeds of *Moringa oleifera*. *Journal of Advance Biology and Biotechnology*, 5, 1–11.
- Abou-Zaid, A. A., & Nadir, A. S. (2014). Quality evaluation of nutritious chocolate and halawa tahinia produced with *Moringa (Moringa oleifera)* leaves powder. *Middle East Journal of Applied Science*, 4(4), 1007–1015.
- Abusuwar, A. O., & Abohassan, R. A. (2017). Effect of *Moringa olifera* leaf extract on growth and productivity of three cereal forages. *Journal of Agricultural Science*, 9(7), 236–243.
- Adedapo, A., Falayi, O., & Oyagbemi, A. (2014). The anti-oxidant, anti-inflammatory, and antinociceptive activities of the methanol leaf extract of *Moringa oleifera* in some laboratory animals. *FASEB Journal*, 28(1). Abstract no 657.19
- Adedapo, A. A., Falayi, O. O., & Oyagbemi, A. A. (2015). Evaluation of the analgesic, anti-inflammatory, anti-oxidant, phytochemical and toxicological properties of the methanolic leaf extract of commercially processed *Moringa oleifera* in some laboratory animals. *Journal Basic and Clinical Physiology and Pharmacology*, 26, 491–499.

- Adhiambo, O. R., Lusweti, K. J., & Morang, G. Z. (2015). Biosorption of Pb<sup>2+</sup> and Cr<sup>2+</sup> using *Moringa oleifera* and their adsorption isotherms. *Science Journal of Analytical Chemistry*, 3(6), 100–108.
- Agra-Neto, A. C., Napoleao, T. H., Pontual, E. V., Santos, N. D., Luz-Lde, A., de Oliveira, C. M., ... Paiva, P. M. (2014). Effect of *Moringa oleifera* lectins on survival and enzyme activities of *Aedes aegypti* larvae susceptible and resistant to organophosphate. *Parasitology Research*, 113, 175–184.
- Agrawal, B., & Mehta, A. (2008). Antiasthmatic activity of *Moringa oleifera* Lam: a clinical study. *Indian Journal of Pharmacology*, 40, 28–31.
- Ahmad, W., Ejaz, S., Anwar, K., & Ashraf, M. (2014). Exploration of the in vitro cytotoxic and antiviral activities of different medicinal plants against Infectious Bursal Disease (IBD) virus. *Central European Journal of Biology*, 9(5), 531–542.
- Aja, P. M., Nwachukwu, N., Ibiam, U. A., Igwenyi, I. O., Offor, C. E., & Orji, U. O. (2014). Chemical constituents of *Moringa oleifera* leaves and seeds from Abakaliki, Nigeria. *American Journal of Phytomedicine and Clinical Therapeutics*, 2(3), 310–321.
- Ajayi, O. B., Akomolafe, S. F., & Adefioye, A. (2014). Proximate analysis, mineral contents and phytochemical screening of *Brachystegia eurycoma* Harms and *Piper guinense* schm and Thonn. *American Journal of Food and Nutrition*, 2(1), 11–17.
- Ajit, K., Choudhary, B. K., & Bandyopadhyay, N. G. (2002). Comparative evaluation of hypoglycaemic activity of some Indian medicinal plants in alloxan diabetic rats. *Journal Ethnopharmacol*, 84, 105–108.
- Akanni, O. E., Adedeji, A. L., & Oloke, K. J. (2014). Upregulation of TNF- $\alpha$  by ethanol extract of *Moringa oleifera* leaves in benzene-induced leukemic Wistar rat: A possible mechanism of anticancer property. *Cancer Research*, 74(19), 3792–3792.
- Akhtar, M., Hasany, S. M., Bhangar, M. I., & Iqbal, S. (2007). Sorption potential of *Moringa oleifera* pods for the removal of organic pollutants from aqueous solutions. *Journal of Hazardous Material*, 141, 546–556.
- Alam, M., Alam, M., Hakim, M., Huq, A., Obidul, A., & Moktadir, S. (2014). Development of fiber enriched herbal biscuits: A preliminary study on sensory evaluation and chemical composition. *International Journal of Nutrition and Food Science*, 3, 246–250.
- Al-Asmari, A. K., Albalawi, S. M., Athar, M. T., Khan, A. Q., Al-Shahrani, H., & Islam, M. (2015). *Moringa oleifera* as an anti-cancer agent against breast and colorectal cancer cell lines. *Plos one*, 10(8), e0135814.
- Ali, G. H., El-Taweel, G. E., & Ali, M. A. (2004). The cytotoxicity and antimicrobial efficiency of *Moringa oleifera* seed extracts. *International Journal of Environmental Studies*, 61, 699–708.
- Al-Juhaimi, F., Babiker, E. E., Ghafoor, K., & Ozcan, M. M. (2016). Fatty acid composition of three different *Moringa* leave oils. *Rivista Italiana Delle Sostanze Grasse Aprile/Giugno*, 93, 111–113.
- Al-Juhaimi, F., Ghafoor, K., Babiker, E. E., Matthauss, B., & Ozcan, M. M. (2017). The biochemical composition of the leaves and seeds meals of moringa species as nonconventional sources of nutrients. *Journal of Food Biochemistry*, 41, e12322.
- Al-Juhaimi, F., Ghafoor, K., Mohamed Ahmed, I. A., Babiker, E. E., & Ozcan, M. M. (2017). Comparative study of mineral and oxidative status of *Sonchus oleraceus*, *Moringa oleifera* and *Moringa peregrina* leaves. *Journal of Food Measurement and Characterization*, 11(4), 1745–1751.
- Al-Malki, A. L., & El Rabey, H. A. (2015). The Antidiabetic effect of low doses of *Moringa oleifera* seeds on streptozotocin induced diabetes and diabetic nephropathy in male rats. *BioMed Research International*, 2015(131040), 1–13.
- Al-Sharif, I., Remmal, A., & Aboussekhra, A. (2013). Eugenol triggers apoptosis in breast cancer cells through E2F1/survivin down-regulation. *BMC Cancer*, 13(600), 1–10.
- Amabye, T. G. (2016). Chemical compositions and nutritional value of *Moringa oleifera* available in the market of Mekelle. *Journal of Food and Nutrition Sciences*, 3(5), 187–190.
- Amaglo, N. K., Deng, J., Bennett, R. N., Lo Curtod, R. B., Rosac, E. A. S., Dominguez-Perlesc, R., ... Crea, F. (2010). Profiling selected phytochemicals and nutrients in different tissues of the multipurpose tree *Moringa oleifera* Lam. grown in Ghana. *Food Chemistry*, 122(4), 1047–1054.
- Anonymous (2005). *The Wealth of India—Raw Materials*. India: Council of Scientific and Industrial Research, New Delhi.
- Anthanont, P., Lumlerkij, N., Akaraseenont, P., Vannasaeng, S., & Sriwijitkamol, A. (2016). *Moringa oleifera* leaf increases insulin secretion after single dose administration: A preliminary study in healthy subjects. *Journal of the Medical Association of Thailand*, 99, 308–313.
- Anwar, F., Ashraf, M., & Bhangar, M. I. (2005). Interprovenance variation in the composition of *Moringa oleifera* oilseeds from Pakistan. *Journal of the American Oil Chemists Society*, 82, 45–51.
- Anwar, F., Latif, S., Ashraf, M., & Gilani, A. H. (2007). *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytotherapy Research*, 21, 17–25.
- Anwar, F., & Rashid, U. (2007). Physico-chemical characteristics of *Moringa oleifera* seeds and seed oil from a wild provenance of Pakistan. *Pakistan Journal of Botany*, 39, 1443–1453.
- Araujo, C. S. T., Alves, V. N., Rezende, H. C., Almeida, I. L. S., de Assunc, R. M. N., Tarley, C. R. T., ... Coelho, N. M. M. (2010). Characterization and use of *Moringa oleifera* seeds as biosorbent for removing metal ions from aqueous effluents. *Water Science and Technology*, 62(9), 2198–2203.
- Arise, A. K., Arise, R. O., Sanusi, M. O., Esan, O. T., & Oyeyinka, S. A. (2014). Effect of *Moringa oleifera* flower fortification on the nutritional quality and sensory properties of weaning food. *Croatia Journal of Food Science and Technology*, 6(2), 65–71.
- Armand-Stussi, I., Basocak, V., Pauly, G., & McCaulley, J. (2003). *Moringa Oleifera*: An interesting source of active ingredients for skin and hair care. *SOFW Journal*, 129(9), 45–52.
- Arora, D. S., & Onsare, J. G. (2014). In vitro antimicrobial evaluation and phytoconstituents of *Moringa oleifera* pod husks. *Industrial Crops and Products*, 52, 125–135.
- Arulselvan, P., Tan, W. S., Gothai, S., Muniandy, K., Fakurazi, S., Esa, N. M., ... Kumar, S. S. (2016). Anti-inflammatory potential of ethyl acetate fraction of *Moringa oleifera* in downregulating the NF- $\kappa$ B signaling pathway in lipopolysaccharide-stimulated macrophage. *Molecules*, 21, 1452–1465.
- Ashfaq, M., & Ashfaq, U. (2012). Evaluation of mosquitocidal activity of water extract of *Moringa oleifera* seeds against culex quinquefasciatus (Diptera: culicidae) in Pakistan. *Pakistan Entomology*, 34, 21–26.
- Asiedu-Gyekye, I. J., Frimpong-Manso, S., Awortwe, C., Antwi, D. A., & Nyarko, A. K. (2014). Micro- and macroelemental composition and safety evaluation of the nutraceutical *Moringa oleifera* leaves. *Journal of Toxicology*, 2014(786979), 1–13.
- Atawodi, S. E., Atawodi, J. C., Idakwo, G. A., Pfundstein, B., Haubner, R., Wurtele, G., ... Owen, R. W. (2010). Evaluation of the polyphenol content and antioxidant properties of methanol extracts of the leaves, stem, and root barks of *Moringa oleifera* Lam. *Journal of Medicinal Food*, 13, 710–716.
- Atsukwei, D., Eze, E. D., Adams, M. D., Adinoyi, S. S., & Ukpabi, C. N. (2014). Hypolipidaemic effect of ethanol leaf extract of *Moringa oleifera* Lam. in experimentally induced hypercholesterolemic wistar rats. *International Journal of Nutrition and Food Science*, 3(4), 355–360.

- Ayanbimpe, G. M., Ojo, T. K., Afolabi, E., Opara, F., Orsaah, S., & Ojerinde, O. S. (2009). Evaluation of extracts of *Jatropha curcas* and *Moringa oleifera* in culture media for selective inhibition of saprophytic fungal contaminants. *Journal of Clinical Laboratory Analysis*, 23, 161–164.
- Azad, A. K., Rasul, M. G., Khan, M. M. K., Sharma, S. C., & Islam, R. (2015). Prospect of *Moringa* seed oil as a sustainable biodiesel fuel in Australia: A review. *Procedia Engineering*, 105, 601–606.
- Azam, M. M., Waris, A., & Nahar, N. M. (2005). Prospects and potential of fatty acid methyl esters of some non-traditional seed oils for use as biodiesel in India. *Biomass Bioenergy*, 29, 293–302.
- Babayehu, A. A., Gbadebo, C. T., Obalolu, M. A., Otunola, G. A., Nmom, I. O., Kayode, R. M. O., ... Ojo, F. E. (2014). Comparison of organoleptic properties of egusi and eforio soup blends produced with *Moringa* and spinach leaves. *Food Science and Quality Management*, 28, 15–18.
- Babiker, E. E., Al-Juhaimi, F., Ghafoor, K., & Abdoun, K. A. (2016). Comparative study on feeding value of *Moringa* leaves as a partial replacement for alfalfa hay in ewes and goats. *Livestock Science*, 195, 21–26.
- Baghel, S. S., Shirastava, N., Baghel, R. S., Agrawal, P., & Rajput, S. (2012). A review of quercetin: Antioxidant and anticancer properties. *World Journal of Pharmacy and Pharmaceutical Science*, 1, 146–160.
- Bakre, A. G., Aderibigbe, A. O., & Ademowo, O. G. (2013). Studies on neuropharmacological profile of ethanol extract of *Moringa oleifera* leaves in mice. *Journal of Ethnopharmacology*, 149, 783–789.
- Bandana, B., Srivastava, R. C., & Mathur, S. N. (1987). Nodulation and nitrogenase activity in *Vigna mungo* in response to seed-soaking and root-dressing treatments of *Moringa* leaf extracts. *Indian Journal of Plant Physiology*, 30(4), 362–367.
- Bargah, R. K., & Das, C. (2014). Isolation and characterization of steroidal glycoside from chloroform extract of the stem bark of *Moringa pterygosperma* Gaertn. *International Journal of Innovative Research Science, Engineering and Technology*, 3, 18319–18322.
- Barichella, M., Pezzoli, G., Faierman, S. A., Raspini, B., Rimoldi, M., Cassani, E., ... Cereda, E. (2019). Nutritional characterisation of Zambian *Moringa oleifera*: Acceptability and safety of short-term daily supplementation in a group of malnourished girls. *International Journal of Food Sciences and Nutrition*, 70(1), 107–115.
- Barreto, M. B., Freitas, J. V. B. D., Silveira, E. R., Bezerra, A. M. E., Nunes, E. P., & Gramosa, N. V. (2009). Volatile and non-volatile chemical constituents of *Moringa oleifera* Lam., Moringaceae. *Revista Brasileira de Farmacognosia*, 19, 893–897.
- Batista, A. B., Oliveira, J. T., Gifoni, J. M., Pereira, M. L., Almeida, M. G., Gomes, V. M., ... Vasconcelos, I. M. (2014). New insights into the structure and mode of action of Mo-CBP 3, an antifungal chitin-binding protein of *Moringa oleifera* seeds. *PLoS one*, 9, e111427.
- Beltran, H. J., & Sanchez, M. J. (2008). Azo dye removal by *Moringa oleifera* seed extracts coagulation. *Coloration Technology*, 124, 310–317.
- Berkovich, L., Earon, G., Ron, I., Rimmon, A., Vexler, A., & Lev-Ari, S. (2013). *Moringa oleifera* aqueous leaf extract down-regulates nuclear factor-kappa-B and increases cytotoxic effect of chemotherapy in pancreatic cancer cells. *BMC Complementary and Alternative Medicine*, 13, 212–219.
- Bharali, R., Tabassum, J., & Azad, M. R. H. (2003). Chemomodulatory effect of *Moringa oleifera* Lam, on hepatic carcinogen metabolizing enzymes, antioxidant parameters and skin papillomagenesis in mice. *Asian Pacific Journal of Cancer Prevention*, 4, 131–139.
- Bhatnagar, M., Parwani, L., Sharma, V., Ganguli, J., & Bhatnagar, A. (2013). Hemostatic: Antibacterial biopolymers from *Acacia arabica* (Lam.) Willd. and *Moringa oleifera* (Lam.) as potential wound dressing materials. *Indian Journal Experimental Biology*, 51, 804–810.
- Bhatnagar, S. S., Santapau, H., Desai, J. D. H., Yellore, S., & Rao, T. N. S. (1961). Biological activity of Indian medicinal plants. Part 1. Antibacterial, antitubercular and antifungal action. *Indian Journal of Medical Research*, 49, 799–805.
- Bhattacharya, S. B., Das, A. K., & Banerji, N. (1982). Chemical investigations on the gum exudates from Sonja (*Moringa oleifera*). *Carbohydrate Research*, 102, 253–262.
- Bhatti, H. N., Mumtaz, B., Hanif, M. A., & Nadeem, R. (2007). Removal of Zn (II) ions from aqueous solution using *Moringa oleifera* Lam. (horse-radish tree) biomass. *Process Biochemistry*, 42, 547–553.
- Bhende, S., & Jadhav, N. (2012). *Moringa* coagulant as a stabilizer for amorphous solids: part I. *AAPS PharmSciTech*, 13, 400–410.
- Bina, B., Mehdinejad, M. H., Dalhammer, G., Rajarao, G., Nikaeen, M., & Attar, M. H. (2010). Effectiveness of *Moringa oleifera* coagulant protein as natural coagulant aid in removal of turbidity and bacteria from turbid waters. *International Scholarly and Scientific Research and Innovation*, 4(7), 261–263.
- Bose, C. K. (2007). Possible role of *Moringa oleifera* Lam. root in epithelial ovarian cancer. *Medscape General Medicine*, 9, 26–34.
- Brilhante, R. S., Sales, J. A., de Souza, S. C. M., Barbosa, F. G., Paiva, M. D., Guedes, G. M. M., ... Rocha, M. F. G. (2015). *Vibrio* spp. from *Macrobrachium amazonicum* prawn farming is inhibited by *Moringa oleifera* extracts. *Asian Pacific journal of tropical medicine*, 8, 919–922.
- Broin, M., Santaella, C., Cuine, S., Kokou, K., Peltier, G., & Joet, T. (2002). Flocculent activity of a recombinant protein from *Moringa oleifera* Lam. seeds. *Applied Microbiology and Biotechnology*, 60, 114–119.
- Bukar, A., Uba, A., & Oyeyi, T. I. (2010). Antimicrobial profile of *Moringa oleifera* Lam. extracts against some food-borne microorganisms. *Bayero Journal of Pure and Applied Sciences*, 3, 43–48.
- Caceres, A., Cabrera, O., Morales, O., Mollinedo, P., & Mendia, P. (1991). Pharmacological properties of *Moringa oleifera*. 1: Preliminary screening for antimicrobial activity. *Journal of Ethnopharmacology*, 33, 213–216.
- Caceres, A., & Lopez, S. (1991). Pharmacological properties of *Moringa oleifera*: 3. Effect of seed extracts in the treatment of experimental pyoderma. *Fitoterapia*, 62(5), 449–450.
- Caceres, A., Saravia, A., Rizzo, S., Zabala, L., De-Leon, E., & Nave, F. (1992). Pharmacologic properties of *Moringa oleifera*. 2: Screening for antispasmodic, antiinflammatory and diuretic activity. *Journal of Ethnopharmacology*, 36, 233–237.
- Casey, T. J. (1997). *Unit treatment processes in water and wastewater engineering*. London, UK: John Wiley & Sons.
- Charoensin, S. (2014). Antioxidant and anticancer activities of *Moringa oleifera* leaves. *Journal of Medicinal Plant Research*, 8(7), 318–325.
- Cheenpracha, S., Park, E. J., Yoshida, W. Y., Barit, C., Wall, M., Pezzuto, J. M., & Chang, L. C. (2010). Potential anti-inflammatory phenolic glycosides from the medicinal plant *Moringa oleifera* fruits. *Bioorganic & Medicinal Chemistry*, 18, 6598–6602.
- Chelliah, R., Ramakrishnan, S., & Antony, U. (2017). Nutritional quality of *Moringa oleifera* for its bioactivity and antibacterial properties. *International Food Research Journal*, 24(2), 825–833.
- Chinma, C., Abu, J., & Akoma, S. (2014). Effect of germinated tigernut and *Moringa* flour blends on the quality of wheat-based bread Food Process. *Preservation*, 38, 721–727.
- Choi, E. J., Debnath, T., Tang, Y., Ryu, Y. B., Moon, S. H., & Kim, E. K. (2016). Topical application of *Moringa oleifera* leaf extract ameliorates experimentally induced atopic dermatitis by the regulation of Th1/Th2/Th17 balance. *Biomedicine and Pharmacotherapy*, 84, 870–877.
- Chollom, S. C., Agada, G. O. A., Gotep, J. G., Mwankon, S. E., Dus, P. C., Bot, Y. S., ... Okwori, A. E. J. (2012). Investigation of aqueous extract of

- Moringa oleifera* lam seed for antiviral activity against Newcastle disease virus in ovo. *Journal of Medicinal Plant Research*, 6(22), 3870–3875.
- Choy, S. Y., Prasad, K. M. N., Wu, T. Y., Raghunandan, M. E., & Ramanan, R. N. (2014). Utilization of plant based natural coagulants as future alternatives towards sustainable water clarification. *Journal of Environmental Science*, 26, 2178–2189.
- Chuang, P. H., Lee, C. W., Chou, J. Y., Murugan, M., Shieh, B. J., & Chen, H. M. (2007). Anti-fungal activity of crude extracts and essential oil of *Moringa oleifera* Lam. *Bioresource Technology*, 98, 232–236.
- Chumark, P., Khunawat, P., Sanvarinda, Y., Phornchirasilp, S., Morales, N. P., Phivthong-Ngam, L., ... Pongrapeeporn, K. U. (2008). The *in vitro* and *ex vivo* antioxidant properties, hypolipidaemic and antiatherosclerotic activities of water extract of *Moringa oleifera* Lam. leaves. *Journal of Ethnopharmacology*, 116, 439–446.
- Coelho, J. S., Santos, N. D., Napoleao, T. H., Gomes, F. S., Ferreira, R. S., Zingali, R. B., ... Paiva, P. M. (2009). Effect of *Moringa oleifera* lectin on development and mortality of *Aedes aegypti* larvae. *Chemosphere*, 77, 934–938.
- Cohen, J., & Grifo, J. A. (2007). Multicentre trial of preimplantation genetic screening reported in the New England journal of medicine: an in-depth look at the findings. *Reproductive Biomedicine Online*, 15(4), 365–366.
- Coppin, J. P., Xu, Y., Chen, H., Pan, M. H., Ho, C. T., Juliania, R., ... Wu, Q. (2013). Determination of flavonoids by LC/MS and anti-inflammatory activity in *Moringa oleifera*. *Journal of Functional Foods*, 5, 1892–1899.
- Dahiru, D., Onubiyi, J. A., & Umaru, H. A. (2006). Phytochemical screening and antiulcerogenic effect of *Moringa oleifera* aqueous leaf extract. *African Journal of Traditional Complementary and Alternative Medicine*, 3, 70–75.
- Dangi, S. Y., Jolly, C. I., & Narayana, S. (2002). Antihypertensive activity of the total alkaloids from the leaves of *Moringa oleifera*. *Pharmaceutical Biology*, 40, 144–148.
- Das, B. R., Kurup, P. A., Rao, N. P. L., & Ramaswamy, A. S. (1957). Antibiotic principle from *Moringa pterygosperma*. Part VIII. Some pharmacological properties and *in vivo* action of pterygospermin and related compounds. *Indian Journal of Medical Research*, 45, 197–206.
- Dastur, J. F. (1964). *Useful Plants of India and Pakistan*. Bombay, India: D.B. Taraporevala Sons and Co.
- Dehshahri, S., Afsharypuor, S., Asghari, G., & Mohagheghzadeh, A. (2012). Determination of volatile glucosinolate degradation products in seed coat, stem and *in vitro* cultures of *Moringa peregrina* (Forssk.) Fiori. *Research in Pharmaceutical Sciences*, 7, 51–56.
- De-Martino, L., De Feo, V., Fratianni, F., & Nazzaro, F. (2009). Chemistry, antioxidant, antibacterial and antifungal activities of volatile oils and their components. *Natural Product Communications*, 4, 1741–1750.
- Devaraj, V. C., & Gopala-Krishna, B. (2013). Antiulcer activity of a polyherbal formulation (PHP) from Indian medicinal plants. *Chinese Journal of Natural Medicines*, 11(2), 145–148.
- Devaraj, V. C., Krishna, B. G., & Viswanatha, G. L. (2011). Simultaneous determination of quercetin, rutin and kaempferol in the leaf extracts of *Moringa oleifera* Lam. and *Raphinus sativus* Linn. by liquid chromatography tandem mass spectrometry. *Dastur of Chinal Integrative Medicine*, 9, 1022–1030.
- Dhakar, R. C., Maurya, S. D., Pooniya, B. K., Bairwa, N., & Gupta, M. S. (2011). *Moringa*: The herbal gold to combat malnutrition. *Chronicles of Young Scientists*, 2, 119–125.
- Diab, K. A., Guru, S. K., Bhushan, S., & Saxena, A. K. (2015). *In vitro* anticancer activities of *Anogeissus latifolia*, *Terminalia bellerica*, *Acacia catechu* and *Moringa oleifera* Indian plants. *Asian Pacific Journal of Cancer Prevention*, 16, 6423–6428.
- Divi, S. M., Bellamkonda, R., & Dasireddy, S. K. (2012). Evaluation of antidiabetic and antihyperlipidemic potential of aqueous extract of *Moringa oleifera* in fructose fed insulin resistant and STZ induced diabetic wistar rats: a comparative study. *Asian Journal of Pharmaceutical and Clinical Research*, 5, 67–72.
- Dogra, P. D., Singh, B. P., & Tandon, S. (1975). Vitamin C content in *Moringa* pod vegetable. *Current Science*, 44(1), 31.
- Dubey, D. K., Dora, J., Kumar, A., & Gulsan, R. K. (2013). A Multipurpose Tree—*Moringa oleifera*. *International Journal of Pharmaceutical and Chemical Science*, 2(1), 415–423.
- Dzotam, J. K., Touani, F. K., & Kuete, V. (2016). Antibacterial and antibiotic-modifying activities of three food plants (*Xanthosoma mafaffa* Lam., *Moringa oleifera* (L.) Schott and *Passiflora edulis* Sims) against multidrug-resistant (MDR) Gram-negative bacteria. *BMC complementary and alternative medicine*, 16(9), 1–8.
- Edoga, C. O., Njoku, O. O., Amadi, E. N., & Okeke, J. J. (2013). Blood sugar lowering effect of *Moringa oleifera* Lam in albino rats. *International Journal of Science and Technology*, 3(1), 88–90.
- Efiong, E. E., Igile, G. O., Mgbeje, B. I. A., Out, E. A., & Ebong, P. E. (2013). Hepatoprotective and anti-diabetic effect of combined extracts of *Moringa oleifera* and *Vernonia amygdalina* in streptozotocin-induced diabetic albino Wistar rats. *Journal of Diabetes Endocrinology*, 4, 45–50.
- Ekor, M. (2014). The growing use of herbal medicines: Issues relating to adverse reactions and challenges in monitoring safety. *Frontiers in Pharmacology*, 4, 1–10.
- El-Alfy, T. S., Ezzat, S. M., Hegazy, A. K., Amer, A. M. M., & Kamel, G. M. (2011). Isolation of biologically active constituents from *Moringa peregrina* (Forssk.) Fiori. (family: Moringaceae) growing in Egypt. *Pharmacognosy Magazine*, 7, 109–115.
- Elbatran, S. A., Abdel-Salam, O. M., Abdelshfeek, K. A., Nazif, N. M., Ismail, S. I., & Hammouda, F. M. (2005). Phytochemical and pharmacological investigations on *Moringa peregrina* (Forssk.). *Natural Product Science*, 11, 199–206.
- Elgamily, H., Moussa, A., Elborae, A., El-Sayed, H., Al-Moghazy, M., & Abdalla, A. (2016). Microbiological assessment of *Moringa oleifera* extracts and its incorporation in novel dental remedies against some oral pathogens. *Open Access Macedonian Journal of Medical Sciences*, 4, 585–590.
- El-Hawary, S. S., El-Sofany, R. H., Abdel-Monem, A. R., Ashour, R. S., & Sleem, A. A. (2012). Polyphenolics content and biological activity of *Plectranthus amboinicus* (Lour.) Spreng growing in Egypt (Lamiaceae). *Pharmacognosy Journal*, 4, 45–54.
- Elsayed, E. A., Sharaf-Eldin, M. A., & Wadaan, M. (2015). *In vitro* evaluation of cytotoxic activities of essential oil from *Moringa oleifera* seeds on HeLa, HepG2, MCF-7, CACO-2 and L929 cell lines. *Asian Pacific Journal of Cancer Prevention*, 16, 4671–4675.
- Emelike, N. J. T., & Ebere, C. O. (2016). Effect of drying techniques of *Moringa* leaf on the quality of chin-chin enriched with *Moringa* leaf powder. *IOSR Journal of Environmental Science, Toxicology and Food Technology*, 10(4), 65–70.
- Emmanuel, S. A., Olajide, O. O., Abubakar, S., Idowu, I. D., Orishadipe, A. T., & Thomas, S. A. (2014). Phytochemical and antimicrobial studies of methanol, ethyl acetate, and aqueous extracts of *Moringa oleifera* seeds. *American Journal of Ethnomedicine*, 1, 346–354.
- Essien, B. A., Essien, J. B., & Eluagu, C. J. (2016). Contributions of *Moringa oleifera* in intercropping systems of food security in the derived Savanna zone of southeastern Nigerai. *Nigeria Agricultural Journal*, 46, 2.
- Eyarefe, O. D., Idowu, A., & Afolabi, J. M. (2015). Healing potentials of oral *Moringa oleifera* leaves extract and tetracycline on methicillin resistant *Staphylococcus aureus* infected wounds of Wistar rats. *Niger Journal of Physiological Science*, 30, 73–78.



- Fabiyyi, J. P., Kela, S. L., Tal, K. M., & Istifanus, W. A. (1993). Traditional therapy of dracunculiasis in the state of Bauchi-Nigeria. *Dakar Medical*, 38, 193–195.
- Fahey, J. W. (2005). *Moringa oleifera*: A review of the medical evidence for its nutritional, therapeutic, and prophylactic properties. Part 1. *Trees for Life Journal*, 1, 5.
- Faizi, S., Siddiqui, B. S., Saleem, R., Aftab, K., Shaheen, F., & Gilani, A. H. (1998). Hypotensive constituents from the pods of *Moringa oleifera*. *Planta Medica*, 64, 225–228.
- Faizi, S., Siddiqui, B. S., Saleem, R., Siddiqui, S., Aftab, K., & Gilani, A. H. (1995). Fully acetylated carbamate and hypotensive thiocarbamate glycosides from *Moringa oleifera*. *Phytochemistry*, 38, 957–963.
- Faizi, S., Siddiqui, S. B., Saleem, R., Siddiqui, S., Aftab, K., & Gilan, A. H. (1994). Novel hypotensive agents, niazimin A, niazimin B, niazicin A and niazicin B from *Moringa oleifera*: Isolation of first naturally occurring carbamates. *Journal of the Chemical Society, Perkin Transactions-1*, 20, 3035–3040.
- Farooq, F., Rai, M., Tiwari, A., Khan, A. A., & Farooq, S. (2012). Medicinal properties of *Moringa oleifera*: an overview of promising healer. *Journal of Medicinal Plants Research*, 6, 4368–4374.
- Ferreira, P. M. P., Farias, D. F., Oliveira, J. T. D. A., & Carvalho, A. D. F. U. (2008). *Moringa oleifera*: Bioactive compounds and nutritional potential. *Brazilian Journal of Nutrition*, 21, 431–437.
- Ferreira, R. S., Napoleao, T. H., Santos, A. F. S., Sa, R. A., Carneiro-da-Cunha, M. G., Morais, M. M. C., ... Paiva, P. M. G. (2011). Coagulant and antibacterial activities of the water-soluble seed lectin from *Moringa oleifera*. *Letters of Applied Microbiology*, 53, 186–192.
- Foidl, N., Makkar, H. P. S., & Becker, K. (2001). The potential of *Moringa oleifera* for agricultural and industrial uses. In L. J. Fuglie (Ed.), *The Miracle Tree- The Multiple Attributes of Moringa*. USA: CTA.
- Fokwen, V. F., Tsafack, H. D., Touko, B. A. H., Djopnang, J. D., Afeanyi, T. A., Kong, A. T., ... Womeni, H. M. (2018). Nutrients composition, phenolic content and antioxidant activity of green and yellow *Moringa oleifera* leaves. *Journal of Food Stability*, 1(1), 46–56.
- Forster, N., Ulrichs, C., Schreiner, M., Arndt, N., Schmidt, R., & Mewis, I. (2015). Ecotype variability in growth and secondary metabolite profile in *Moringa oleifera*: Impact of sulfur and water availability. *Journal of Agricultural and Food Chemistry*, 63, 2852–2861.
- Francis, J. A., Jayaprakasam, B., Olson, L. K., & Nair, M. G. (2004). Insulin secretagogues from *Moringa oleifera* with cyclooxygenase enzyme and lipid peroxidation inhibition activities. *Helvetica Chimica Acta*, 87, 317–326.
- Freire, J. E., Vasconcelos, I. M., Moreno, F. B., Batista, A. B., Lobo, M. P. D., Pereira, M. L., ... Oliveira, J. T. A. (2015). Mo-CBP 3, an antifungal chitin-binding protein from *Moringa oleifera* seeds, is a member of the 2S albumin family. *PLoS one*, 10, e0119871.
- Fuglie, L. J. (1999). *The Miracle Tree, Moringa oleifera: Natural Nutrition for the Tropics*. New York, USA: Church World Service.
- Fuglie, L. J. (2005). *The Moringa tree: A local solution to malnutrition*. USA: Church World Service in Senegal.
- Gaikwad, M., Kale, S., Bhandare, S., Urunkar, V., & Rajmane, A. (2011). Extraction, characterization and comparison of fixed oil of *Moringa oleifera* Lam. & *Moringa concanensis* Nimmo Fam. Moringaceae. *International Journal of PharmTech Research*, 3, 1567–1575.
- Galuppo, M., Giacoppo, S., De Nicola, G. R., Iori, R., Navarra, M., Lombardo, G. E., ... Mazzon, E. (2014). Antiinflammatory activity of glucomoringin isothiocyanate in a mouse model of experimental autoimmune encephalomyelitis. *Fitoterapia*, 95, 160–174.
- Galuppo, M., Nicola, G. R., Iori, R., Utri, P. D., Bramanti, P., & Mazzon, E. (2013). Antibacterial activity of glucomoringin bioactivated with myrosinase against two important pathogens affecting the health of long-term patients in hospitals. *Molecules*, 18(11), 14340–14348.
- Ganatra, T. H., Umang, H. J., Payal, N. B., Tusharbindu, R. D., & Pravin, R. T. (2012). A panoramic view on pharmacognostic, pharmacological, nutritional, therapeutic and prophylactic values of *Moringa oleifera* Lam. *International Research Journal of Pharmacy*, 3(6), 1–7.
- Ganguly, R., & Guha, D. (2008). Alteration of brain monoamines and EEG wave pattern in rat model of Alzheimer's disease and protection by *Moringa oleifera*. *Indian Journal of Medical Research*, 128, 744–755.
- Ganguly, R., Hazra, R., Ray, K., & Guha, D. (2005). Effect of *Moringa oleifera* in experimental model of Alzheimer's disease: Role of antioxidants. *Annals of Neuroscience*, 12, 36–39.
- Garcia-Fayos, B., Arnal, J. M., Piris, J., & Sancho, M. (2016). Valorization of *Moringa oleifera* seed husk as biosorbent: isotherm and kinetics studies to remove cadmium and copper from aqueous solutions. *Desalination and Water Treatment*, 57(48–49), 23382–23396.
- Ghasi, S., Nwobodo, E., & Ofili, J. (2000). Hypocholesterolemic effects of crude extract of leaf of *Moringa oleifera* Lam. in high-fat diet fed Wistar rats. *Journal of Ethnopharmacology*, 69, 21–25.
- Ghiridhari, V. V. A., Malhati, D., & Geetha, K. (2011). Anti-diabetic properties of drumstick (*Moringa oleifera*) leaf tablets. *International Journal of Health Nutrition*, 2, 1–5.
- Gholap, P. A., Nirmal, S. A., Pattan, S. R., Pal, S. C., & Mandal, S. C. (2012). Potential of *Moringa oleifera* root and *Citrus sinensis* fruit extracts in the treatment of ulcerative colitis in mice. *Pharmaceutical Biology*, 50, 1297–1302.
- Ghosh, R., Ganapathy, M., Alworth, W. L., Chan, D. C., & Kumar, A. P. (2009). Combination of 2-methoxyestradiol (2-ME 2) and eugenol for apoptosis induction synergistically in androgen independent prostate cancer cells. *The Journal of steroid biochemistry and molecular biology*, 113(1), 25–35.
- Giacoppo, S., Galuppo, M., Montaut, S., Iori, R., Rollin, P., Bramanti, P., & Mazzon, E. (2015). An overview on neuroprotective effects of isothiocyanates for the treatment of neurodegenerative diseases. *Fitoterapia*, 106, 12–21.
- Gidamis, A. B., Panga, J. T., Sarwatt, S. V., Chove, B. E., & Shayo, N. B. (2003). Nutrient and antinutrient contents in raw and cooked young leaves and immature pods of *Moringa oleifera* Lam. *Ecology of Food and Nutrition*, 42(6), 399–411.
- Gismondi, A., Canuti, L., Impei, S., Marco, G. D., Kenzo, M., Colizzi, V., & Canini, A. (2013). Antioxidant extracts of African medicinal plants induce cell cycle arrest and differentiation in B16F10 melanoma cells. *International Journal of Oncology*, 43, 956–964.
- Gopalakrishnan, L., Doriya, K., & Kumar, D. S. (2016). *Moringa oleifera*: A review on nutritive importance and its medicinal application. *Food Science and Human Wellness*, 5, 49–56.
- Goswami, S. K., Inamdar, M. N., Dethle, S. M., Gururaj, G. M., Jamwal, R., Bhaskar, A., ... Agarwal, A. (2016). Erectogenic and aphrodisiac property of *Moringa oleifera*: Involvement of soluble epoxide hydrolase enzyme. *Phytotherapy Research*, 30(7), 1119–1127.
- Govardhan-Singh, R. S., Negi, P. S., & Radha, C. (2013). Phenolic composition, antioxidant and antimicrobial activities of free and bound phenolic extracts of *Moringa oleifera* seed flour. *Journal of Functional Foods*, 5, 1883–1891.
- Guevara, A. P., Vargas, C., Sakurai, H., Fujiwara, Y., Hashimoto, K., Maoka, T., ... Nishino, H. (1999). An antitumor promoter from *Moringa oleifera* Lam. *Mutation Research—Genetic Toxicology and Environmental Mutagenesis*, 440, 181–188.

- Gupta, A., Gautam, M. K., Singh, R. K., Kumar, M. V., Rao, C. V., Goel, R. K., & Anupurba, S. (2010). Immunomodulatory effect of *Moringa oleifera* Lam. extract on cyclophosphamide induced toxicity in mice. *Indian Journal Experimental Biology*, 48, 1157–1160.
- Gupta, M., Mazumder, U. K., & Chakrabarti, S. (1999). CNS activities of methanolic extract of *Moringa oleifera* root in mice. *Fitoterapia*, 70, 244–250.
- Gupta, R., Dubey, D. K., Kannan, G. M., & Flora, S. J. (2007). Concomitant administration of *Moringa oleifera* seed powder in the remediation of arsenic-induced oxidative stress in mouse. *Cell Biology International*, 31, 44–56.
- Gupta, R., Kannan, G. M., Sharma, M., & Flora, S. J. S. (2005). Therapeutic effects of *Moringa oleifera* on arsenic-induced toxicity in rats. *Environmental Toxicology and Pharmacology*, 20, 456–464.
- Gupta, R., Mathur, M., Bajaj, V. K., Kataria, P., Yadav, S., Kamal, R., & Gupta, R. S. (2012). Evaluation of antidiabetic and antioxidant activity of *Moringa oleifera* in experimental diabetes. *Journal of Diabetes*, 4, 164–171.
- Gupta, V. K., & Arya, V. (2011). A review on potential diuretics of Indian medicinal plants. *Journal of Chemical and Pharmaceutical Research*, 3(1), 613–620.
- Habtemariam, S., & Varghese, G. K. (2015). Extractability of rutin in herbal tea preparations of *Moringa stenopetala* leaves. *Beverages*, 1, 169–182.
- Rim J. S., Ha-Lee K., Ha-Shin D., Kwon S.S., & Hwang J.S. (2014). Synergistic antimicrobial efficacy of mesoporous ZnO loaded with 4-( $\alpha$ -L-rhamnosyloxy)-benzyl isothiocyanate isolated from the *Moringa oleifera* seed. *The Journal of General and Applied Microbiology*, 60: 251–255.
- Hamza, A. A. (2010). Ameliorative effects of *Moringa oleifera* Lam. seed extract on liver fibrosis in rats. *Food Chemistry and Toxicology*, 48, 345–355.
- Han, J. M., Lee, Y. J., Lee, S. Y., Kim, E. M., Moon, Y., Kim, H. W., & Hwang, O. (2007). Protective effect of sulforaphane against dopaminergic cell death. *Journal of Pharmacology and Experimental Therapeutics*, 321, 249–256.
- Hannan, M. A., Kang, J. Y., Mohibullah, M., Hong, Y. K., Lee, H., Choi, J. S., ... Moon, I. S. (2014). *Moringa oleifera* with promising neuronal survival and neurite outgrowth promoting potentials. *Journal of Ethnopharmacology*, 152, 142–150.
- Harada, H., Yamashita, U., Kurihara, H., Fukushi, E., Kawabata, J., & Kamei, Y. (2001). Antitumor activity of palmitic acid found as a selective cytotoxic substance in a marine red alga. *Anticancer research*, 22(5), 2587–2590.
- Hekmat, S., Morgan, K., Soltani, M., & Gough, R. (2015). Sensory evaluation of locally grown fruit purees and inulin fibre on probiotic yogurt in Mwanza, Tanzania and the microbial analysis of probiotic yogurt fortified with *Moringa oleifera*. *Journal of Health, Population and Nutrition*, 33(1), 60–67.
- Hukkeri, V. I., Nagathan, C. V., Karadi, R. V., & Patil, B. S. (2006). Antipyretic and wound healing activities of *Moringa oleifera* Lam. in rats. *Indian Journal of Pharmacy Science*, 68, 124–126.
- Ibrahim, M. A., Mohammed, A., Isah, M. B., & Aliyu, A. B. (2014). Antitrypanosomal activity of African medicinal plants: a review update. *Journal of Ethnopharmacology*, 154, 26–54.
- Iffiu-Soltesz, Z., Wanecq, E., Lomba, A., Portillo, M. P., Pellati, F., Szoko, E., ... Valet, P. (2010). Chronic benzylamine administration in the drinking water improves glucose tolerance, reduces body weight gain and circulating cholesterol in high fat diet-fed mice. *Pharmacology Research*, 61, 355–363.
- Igado, O. O., & Olopade, J. O. (2016). A review on the possible neuroprotective effects of *Moringa oleifera* leaf extract. *Niger Journal of Physiological Science*, 31, 183–187.
- Igwilo, I. O., Oloyode, O. B., & Enemor, V. H. A. (2007). Nutrient composition and the effect of processing on *Canavalia ensiformis* seed. *International Journal of Agriculture and Food Systems*, 1(1), 48–50.
- Inbathamizh, L., & Padmini, E. (2012). Gas chromatography-mass spectrometric analyses of methanol extract of *Moringa oleifera* flowers. *International Journal of Chemical and Analytical Science*, 2, 1394–1397.
- Ingale, S. P., & Gandhi, F. P. (2016). Effect of aqueous extract of *Moringa oleifera* leaves on pharmacological models of epilepsy and anxiety in mice. *International Journal of Epilepsy*, 3(1), 12–19.
- Irfan, H. M., Asmawi, M. Z., Khan, N. A. K., & Sadikun, A. (2016). Effect of ethanolic extract of *Moringa oleifera* Lam. leaves on body weight and hyperglycemia of diabetic rats. *Pakistan Journal of Nutrition*, 15(2), 112–117.
- Jabeen, R., Shahid, M., Jamil, A., & Ashraf, M. (2008). Microscopic evaluation of the antimicrobial activity of seed extracts of *Moringa oleifera*. *Pakistan Journal of Botany*, 40, 1349–1358.
- Jahn, S. A., Musnad, H. A., & Burgstaller, H. (1986). The tree that purifies water: Cultivating multipurpose Moringaceae in the Sudan. *Unasylva*, 38, 23–28.
- Jain, P. G., Patil, S. D., Haswani, N. G., Girase, M. V., & Surana, S. J. (2010). Hypolipidemic activity of *Moringa oleifera* Lam. Moringaceae, on high fat diet induced hyperlipidemia in albino rats. *Revista Brasileira de Farmacognosia*, 20, 969–973.
- Jaiswal, D., Rai, P. K., Kumar, A., Mehta, S., & Watal, G. (2009). Effect of *Moringa oleifera* Lam. leaves aqueous extract therapy on hyperglycemic rats. *Journal of Ethnopharmacology*, 123, 392–396.
- Jayawardana, B. C., Liyanage, R., Lalantha, N., Iddamalagoda, S., & Weththasinghe, P. (2015). Antioxidant and antimicrobial activity of drumstick (*Moringa oleifera*) leaves in herbal chicken sausages. *LWT—Food Science and Technology*, 64, 1204–1208.
- Johri, P. K., Reeta, J., & Johri, R. (2004). Dissipation of water pollutants through extract of drumstick, *Moringa oleifera*. *Journal of Applied Zoological Research*, 15(1), 103–105.
- Joung, H., Kim, B., Park, H., Lee, K., Kim, H. H., Sim, H. C., ... Do, M. S. (2017). Fermented *Moringa oleifera* decreases hepatic adiposity and ameliorates glucose intolerance in high-fat diet-induced obese mice. *Journal of Medicinal Food*, 20, 439–447.
- Jung, I. L. (2014). Soluble extract from *Moringa oleifera* leaves with a new anticancer activity. *Plos One*, 9, e95492.
- Jung, I. L., Lee, J. H., & Kang, S. C. (2015). A potential oral anticancer drug candidate, *Moringa oleifera* leaf extract, induces the apoptosis of human hepatocellular carcinoma cells. *Oncology Letter*, 10, 1597–1604.
- Kafuku, G., & Mbarawa, M. (2010). Alkaline catalyzed biodiesel production from *Moringa oleifera* oil with optimized production parameters. *Applied Energy*, 87, 2561–2565.
- Kalavathy, M. H., & Miranda, L. R. (2010). *Moringa oleifera* - A solid phase extractant for the removal of copper, nickel and zinc from aqueous solutions. *Chemical Engineering Journal*, 188–199.
- Kar, A., Choudhary, B. K., & Bandyopadhyay, N. G. (2003). Comparative evaluation of hypoglycemic activity of some Indian medicinal plants in alloxan diabetic rats. *Journal of Ethnopharmacology*, 84, 105–108.
- Karadi, R. V., Gadge, N. B., Alagawadi, K. R., & Savadi, R. V. (2006). Effect of *Moringa oleifera* Lam. root-wood on ethylene glycol induced urolithiasis in rats. *Journal of Ethnopharmacology*, 105, 306–311.

- Karim, O., Kayode, R., Oyeyinka, S., & Oyeyinka, A. (2015). Physicochemical properties of stiff dough 'amala' prepared from plantain (*Musa Paradisica*) flour and Moringa (*Moringa oleifera*) leaf powder. *Food Health Disparities*, 4, 48–58.
- Karthivashan, G., Fard, M. T., Arulselvan, P., Abas, F., & Fakurazi, S. (2013). Identification of bioactive candidate compounds responsible for oxidative challenge from hydro-ethanolic extract of *Moringa oleifera* leaves. *Journal of Food Science*, 78(9), C1368–C1375.
- Katayon, S., Noor, M. J. M., Asma, M., Abdul-Ghani, L. A., Thamer, A. M., Azni, I., ... Suleyman, A. M. (2006). Effects of storage conditions of *Moringa oleifera* seeds on its performance in coagulation. *Bioresource Technology*, 97, 1455–1460.
- Kaur, A., Kaur, P. K., Singh, S., & Singh, I. P. (2014). Antileishmanial compounds from *Moringa oleifera* Lam. *Z Naturforsch C*, 69, 110–116.
- Kaur, G., Athar, M., & Alam, M. S. (2010). Eugenol precludes cutaneous chemical carcinogenesis in mouse by preventing oxidative stress and inflammation and by inducing apoptosis. *Molecular Carcinogenesis*, 49(3), 290–301.
- Kekuda, T. R. P., Mallikarjun, N., Swathi, D., Nayana, K. V., Aiyar, M. B., & Rohini, T. R. (2010). Antibacterial and antifungal efficacy of steam distillate of *Moringa oleifera* Lam. *Journal of Pharmaceutical Sciences and Research*, 2, 34–37.
- Khalafalla, M. M., Abdellatef, E., Dafalla, H. M., Nassrallah, A. A., Aboul-Enain, K. M., Lightfoot, D. A., ... El-Shemy, H. A. (2010). Active principle from *Moringa oleifera* lam leaves effective against two leukemias and a hepatocarcinoma. *African Journal of Biotechnology*, 9, 8467–8471.
- Khan, M. T., Ather, A., Thompson, K. D., & Gambari, R. (2005). Extracts and molecules from medicinal plants against herpes simplex viruses. *Antiviral Research*, 67, 107–119.
- Kituyi, J. L., Foulkes, M., Worsfold, P., Ongulu, R. A., Kiplagat, A., & Gachanja, A. (2013). Efficiency of pre-treated *Moringa oleifera* for the removal of Cd<sup>2+</sup> and Zn<sup>2+</sup> ions from wastewaters. *Ecohydrology and Hydrobiology*, 13, 267–271.
- Kivevele T., & Huan Z. (2015). Review of the stability of biodiesel produced from less common vegetable oils of African origin. *South African Journal of Science*, 111, 1–7.
- Kleiman, R., Ashley, D. A., & Brown, J. H. (2008). Comparison of two seed oils used in cosmetics, moringa and marula. *Indian Crops Production*, 28, 361–364.
- Knothe, G. (2008). "Designer" biodiesel: Optimizing fatty ester composition to improve fuel properties. *Energy Fuels*, 22, 1358–1364.
- Kohler, I., Jenett-Siems, K., Siems, K., Hernandez, M. A., Ibarra, R. A., Berendsohn, W. G., ... Eich, E. (2002). In vitro antiplasmodial investigation of medicinal plants from El Salvador. *Z Naturforsch C*, 57(3–4), 277–281.
- Konmy, B. B. S., Olounlade, P. A., Allou, S. D., Azando, E. V. B., & Hounzangbe-Adote, M. S. (2016). A review on phytochemistry and pharmacology of *Moringa oleifera* leaves (Moringaceae). *Journal of Pharmacognosy and Phytochemistry*, 5(5), 325–330.
- Krishnamurthy, P. T., Vardarajulu, A., Wadhvani, A., & Patel, V. (2015). Identification and characterization of a potent anticancer fraction from the leaf extracts of *Moringa oleifera* L. *Indian Journal of Experimental Biology*, 53, 98–103.
- Kumar, V., Pandey, N., Mohan, N., & Singh, R. P. (2012). Antibacterial & antioxidant activity of different extract of *Moringa oleifera* leaves: An in vitro study. *International Journal of Pharmaceutical Sciences Review and Research*, 12, 89–94.
- Kumari, P., Sharma, P., Srivastava, S., & Srivastava, M. M. (2006). Biosorption studies on shelled *Moringa oleifera* Lamarck seed powder: Removal and recovery of arsenic from aqueous system. *International Journal of Mineral Processing*, 78, 131–139.
- Kumbhare, M. R., Guleha, V., & Sivakumar, T. (2012). Estimation of total phenolic content, cytotoxicity and in vitro antioxidant activity of stem bark of *Moringa oleifera*. *Asian Pacific Journal of Tropical Disease*, 2, 144–150.
- Kurmi, R., Ganeshpurkar, A., Bansal, D., Agnihotri, A., & Dubey, N. (2014). Ethanol extract of *Moringa oleifera* prevents in vitro glucose induced cataract on isolated goat eye lens. *Indian Journal of Ophthalmology*, 62(2), 154–157.
- Kurokawa, M., Wadhvani, A., Kai, H., Hidaka, M., Yoshida, H., Sugita, C., ... Hagiwara, A. (2016). Activation of cellular immunity in herpes simplex virus type 1-infected mice by the oral administration of aqueous extract of *Moringa oleifera* Lam. leaves. *Phytotherapy Research*, 30, 797–804.
- Kushwaha, V., Saxena, K., Verma, S. K., Lakshmi, V., Sharma, R. K., & Murthy, P. K. (2011). Antifilarial activity of gum from *Moringa oleifera* Lam. on human lymphatic filarial *Brugia malayi*. *Chronicals of Young Scientists*, 2, 201–206.
- Lalas, S., Gortzi, O., Athanasiadis, V., Tsaknis, J., & Chinou, I. (2012). Determination of antimicrobial activity and resistance to oxidation of *Moringa peregrina* seed oil. *Molecules*, 17, 2330–2334.
- Lalas, S., & Tsaknis, J. (2002a). Characterization of *Moringa oleifera* seed oil variety Periyakulam-1. *Journal of Food Composition and Analysis*, 15, 65–77.
- Lalas, S., & Tsaknis, J. (2002b). Extraction and identification of natural antioxidant from the seeds of the *Moringa oleifera* tree variety of Malawi. *Journal of the American Oil Chemists' Society*, 79, 677–683.
- Lambole, V., & Kumar, U. (2012). Effect of *Moringa oleifera* Lam. on normal and dexamethasone suppressed wound healing. *Asian Pacific Journal of Tropical Biomedicine*, 2, S219–S223.
- Lea M.A., Akinpelu T., Amin R., Yarlagadda K., Enugala R., Dayal B., & Des Bordes C. (2012). Abstract 5439: inhibition of growth and induction of differentiation of colon cancer cells by extracts from okra (*Abelmoschus esculentus*) and drumstick (*Moringa oleifera*). In: Proceedings of the 103rd Annual Meeting of the American Association for Cancer Research; 2012 Mar 31-Apr 4; Chicago, IL. Philadelphia (PA): AACR Cancer Research 2012, 72(8 Suppl): Abstract no. 5439.
- Lee, H. J., Jeong, Y. J., Lee, T. S., Park, Y. Y., Chae, W. G., Chung, I. K., ... Chang, Y. C. (2013). *Moringa* fruit inhibits LPS induced NO/iNOS expression through suppressing the NFκB activation in RAW264.7 cells. *The American Journal of Chinese Medicine*, 41, 1109–1123.
- Lee, K. Y., Yang, H. J., & Song, K. B. (2016). Application of a puffer fish skin gelatin film containing *Moringa oleifera* Lam. leaf extract to the packaging of Gouda cheese. *Journal of Food Science Technology*, 53(11), 3876–3783.
- Leone, A., Bertoli, S., Di Lello, S., Bassoli, A., Ravasenghi, S., Borgonovo, G., ... Battezzati, A. (2018). Effect of *Moringa oleifera* leaf powder on post-prandial blood glucose response: In vivo study on Saharawi people living in refugee camps. *Nutrients*, 10(1491), 1–14.
- Leone, A., Fiorillo, G., Criscullo, F., Ravasenghi, S., Santagostini, L., Fico, G., ... Bertoli, S. (2015). Nutritional characterization and phenolic profiling of *Moringa oleifera* leaves grown in Chad, Sahrawi refugee camps, and Haiti. *International Journal of Molecular Science*, 16, 18923–18937.
- Leone, A., Spada, A., Battezzati, A., Schiraldi, A., Aristil, J., & Bertoli, S. (2015). Cultivation, genetic, ethnopharmacology, phytochemistry and pharmacology of *Moringa oleifera* leaves: an overview. *International Journal of Molecular Science*, 16, 12791–12835.
- Liang, L., Wang, C., Li, S., Chu, X., & Sun, K. (2019, In press). Nutritional compositions of Indian *Moringa oleifera* seed and antioxidant activity

- of its polypeptides. *Food Science and Nutrition*. <https://doi.org/10.1002/fsn3.1015>
- Lijesh, K. P., & Malhotra, R. (2016). Reduction of turbidity of water using *Moringa oleifera*. *International Journal of Applied Engineering Research*, 11(2), 1414–1423.
- Lipipun, V., Kurokawa, M., Suttisri, R., Taweechotipatr, P., Pramyothin, P., Hattori, M., & Shiraki, K. (2003). Efficacy of Thai medicinal plant extracts against herpes simplex virus type 1 infection *in vitro* and *in vivo*. *Antiviral Research*, 60, 175–180.
- Little, E. L. J. R., & Wadsworth, F. H. (1964). *Common trees of Puerto Rico and the Virgin Islands. Agriculture Handbook*. 249. Washington, D.C.: U.S. Dep. Agric.
- Lizzy, K. S., Rao, P. L. N., & Puttaswamy, T. L. (1968). Chemotherapy of bacterial infections. Part 4: potential anticholera agents. *Indian Journal of Experimental Biology*, 6(3), 168–169.
- Lurling, M., & Beekman, W. (2010). Anti-cyanobacterial activity of *Moringa oleifera* seeds. *Journal of Applied Phycology*, 22, 503–510.
- Madi, N., Dany, M., Abdoun, S., & Usta, J. (2016). *Moringa oleifera*'s nutritious aqueous leaf extract has anticancerous effects by compromising mitochondrial viability and ROS-dependent manner. *Journal of American College of Nutrition*, 35, 604–613.
- Maghu, T. K., Sharma, A., & Younis, K. (2017). Effect of drumstick leaves incorporation on Khakhra. In S. Ul Islam (Ed.), *Plant based natural products: Derivatives and Applications* (1st ed.) (pp. 129–142). Beverly, MA: Scrivener Publishing LLC, Wiley. ISBN: 978-1-119-42383-6. <https://doi.org/10.1002/9781119423898.ch7>
- Mahajan, S. G., & Mehta, A. A. (2007). Inhibitory action of ethanolic extract of seeds of *Moringa oleifera* Lam. on systemic and local anaphylaxis. *Journal of Immunotoxicology*, 4, 287–294.
- Mahajan, S. G., & Mehta, A. A. (2008). Effect of *Moringa oleifera* Lam. seed extract on ovalbumin-induced airway inflammation in guinea pigs. *Inhal Toxicology*, 20, 897–909.
- Mahatab, S. N., Ali, A., & Asaduzzaman, A.H.M. (1987). Nutritional potential of sajna leaves in goats. *Livestock Advisor*, 12, 9–12.
- Maheshwari, K., Yadav, R. K., Malhotra, J., Dhawan, N. G., & Mohan, L. (2014). Fascinating nutritional, prophylactic, therapeutic and socio-economic reconcile attributable to drum stick tree (*Moringa oleifera* Lam.). *Global Journal of Medical research: B Pharma, Drug Discovery, Toxicology and Medicine*, 14(1), 11–22.
- Mahmood, K. T., Mugall, T., & Haq, U. (2010). *Moringa oleifera*: A natural gift-a review. *Journal of Pharmaceutical Sciences and Research*, 2(11), 775–781.
- Maiyo, F. C., Moodley, R., & Singh, M. (2016). Cytotoxicity, antioxidant and apoptosis studies of quercetin-3-O-glucoside and 4-(beta-Dglucopyranosyl-1- > 4-alpha-L-rhamnopyranosyloxy)-benzyl isothiocyanate from *Moringa oleifera*. *Anti-Cancer Agents in Medicinal Chemistry*, 16, 648–656.
- Makkar, H. P. S., & Becker, K. (1996). Nutritional value and antinutritional components of whole and ethanol extracted *Moringa oleifera* leaves. *Animal Feed Science and Technology*, 63, 211–228.
- Maldini, M., Maksoud, S. A., Natella, F., Montoro, P., Petretto, G. L., Foddai, M., ... Pintorea, G. (2014). *Moringa oleifera*: Study of phenolics and glucosinolates by mass spectrometry. *Journal of Mass Spectrometry*, 49(9), 900–910.
- Manajeji, H., Jafari, S., Zaringhalam, J., Rezazadeh, S., & Taghizadfarid, R. (2011). Analgesic effects of methanolic extracts of the leaf or root of *Moringa oleifera* on complete Freund's adjuvant-induced arthritis in rats. *Chinese Journal of Integrative Medicine*, 9, 216–222.
- Mandloi, M., Chaudhari, S., & Folkard, G. K. (2004). Evaluation of natural coagulants for direct filtration. *Environmental Technology*, 25, 481–489.
- Manguro, L. O., & Lemmen, P. (2007). Phenolics of *Moringa oleifera* leaves. *Natural Product Research*, 21, 56–68.
- Manikandan, P., Vinothini, G., Priyadarsini, R. V., Prathiba, D., & Nagini, S. (2011). Eugenol inhibits cell proliferation via NF- $\kappa$ B suppression in a rat model of gastric carcinogenesis induced by MNNG. *Investigational New Drugs*, 29(1), 110–117.
- Marcel, A., Hubert, M., Bienvenu, M. J., & Pascal, O. (2016). Physico-chemical characteristics and biochemical potential of *Moringa oleifera* Lam. (Moringaceae). *Der Pharmacia Lettre*, 8(18), 43–47.
- Marcela, V. J., Almatrafi, M. M., & Fernandez, M. J. (2017). Bioactive components in *Moringa oleifera* leaves protect against chronic disease. *Antioxidants*, 6, 1–13.
- Mark, E. O., & Sylvain, G. R. (2006). *Moringa hildebrandtii* (Moringaceae): A tree extinct in the wild but preserved by indigenous horticultural practices in Madagascar, Adansonia. *Journal of Food and Nutrition*, 22, 217–221.
- Marrufo, T., Nazzaro, F., Mancini, E., Fratianni, F., Coppola, R., De Martino, L., ... De Feo, V. (2013). Chemical composition and biological activity of the essential oil from leaves of *Moringa oleifera* Lam. cultivated in Mozambique. *Molecules*, 18, 10989–11000.
- Martin, F. W., & Ruperte, R. M. (1979). *Edible Leaves of the Tropics* (2nd ed.). Southern Region, Mayaguez, PR: U.S. Department of Agriculture Science and Education Admin., Agric. Res.
- Mataka, L. M., Sajidu, S. M. I., Masamba, W. R. L., & Mwatseteza, J. F. (2010). Cadmium sorption by *Moringa stenopetala* and *Moringa oleifera* seed powders: Batch, time, temperature, pH and adsorption isotherm studies. *International Journal of Water Resources and Environmental Engineering*, 2, 50–59.
- Matsuda, H., Ochi, M., Nagatomo, A., & Yoshikawa, M. (2007). Effects of allyl isothiocyanate from horseradish on several experimental gastric lesions in rats. *European journal of pharmacology*, 561(1), 172–181.
- Maurya, S. K., & Singh, A. K. (2014). Clinical efficacy of *Moringa oleifera* Lam. Stems bark in urinary tract infections. *International Scholarly Research Notices*, 2014, 1–7.
- Mazumder, U. K., Gupta, M., Chakrabarti, S., & Pal, D. (1999). Evaluation of hematological and hepatorenal functions of methanolic extract of *Moringa oleifera* Lam. root treated mice. *Indian Journal of Experimental Biology*, 37, 612–614.
- Mbikay, M. (2012). Therapeutic potential of *Moringa oleifera* leaves in chronic hyperglycemia and dyslipidemia: a review. *Frontiers in Pharmacology*, 3, 1–12.
- Mehta, J., Shukla, A., Bukhariya, V., & Charde, R. (2011). The magic remedy of *Moringa oleifera*: an overview. *International Journal of Biomedical and Advance Research*, 2, 215–227.
- Mehta, L. K., Balaraman, R., Amin, A. H., Bafna, P. A., & Gulati, O. D. (2003). Effect of fruits of *Moringa oleifera* on the lipid profile of normal and hypercholesterolaemic rabbits. *Journal of Ethnopharmacology*, 86, 191–195.
- Mekonen, A., & Gebreyesus, T. (2000). Chemical investigation of the leaves of *Moringa stenopetala*. *Bulletin of the Chemical Society of Ethiopia*, 14, 51–55.
- Mekonnen, Y., & Drager, B. (2003). Glucosinolates in *Moringa stenopetala*. *Planta Medica*, 69, 380–382.
- Memon, G. M., Memon, S. A., & Memon, A. R. (1985). Isolation and structure elucidation of moringyne - a new glycoside from seeds of *Moringa oleifera* Lam. *Pakistan Journal of Scientific and Industrial Research*, 28, 7–9.
- Mendieta, A. B., Spornly, E., Reyes-Sanchez, N., & Spornly, R. (2011). Feeding *Moringa oleifera* fresh or ensiled to dairy cows effects on milk

- yield and milk flavor. *Tropical Animal Health and Production*, 43, 1039–1047.
- Metwally, F., Metwally, M., Rashad, M., Ahmed, H. H., Mahmoud, A. A., & Raouf, E. R. A. (2017). Molecular mechanisms of the anti-obesity potential effect of *Moringa oleifera* in the experimental model. *Asian Pacific Journal of tropical biomedicine*, 7(3), 214–221.
- Minaiyan, M., Asghari, G., Taheri, D., Saeidi, M., & Nasr-Esfahani, S. (2014). Anti-inflammatory effect of *Moringa oleifera* Lam. seeds on acetic acid-induced acute colitis in rats. *Avicenna Journal of Phytomedicine*, 4(2), 127–136.
- Mirella, L. P., Jose, T. A. O., Juliana, M. G., Raquel, O. R., Daniele, O. V. S., & Ilka, M. V. (2011). Purification of a chitin-binding protein from *Moringa oleifera* seeds with potential to relieve pain and inflammation. *Protein and Peptide Letters*, 18(11), 1078–1085.
- Misra, A., Srivastava, S., & Srivastava, M. (2014). Evaluation of anti diarrheal potential of *Moringa oleifera* (Lam.) leaves. *Journal of Pharmacognosy and Phytochemistry*, 2(5), 43–46.
- Monera, G. T., & Maponga, C. C. (2010). *Moringa oleifera* supplementation by patients on antiretroviral therapy. *Journal of International AIDS Society*, 13(4), 188.
- Monera, T. G., Wolfe, A. R., Maponga, C. C., Benet, L. Z., & Guglielmo, J. (2008). *Moringa oleifera* leaf extracts inhibit 6 $\beta$ -hydroxylation of testosterone by CYP3A4. *The Journal of Infection in Developing Countries*, 2, 379–383.
- Morton, J. F. (1991). The horseradish tree, *Moringa pterygosperma* (Moringaceae) - a boon to arid lands? *Economic Botany*, 45(3), 318–333.
- Moura, M. C., Napoleao, T. H., Coriolano, M. C., Paiva, P. M., Figueiredo, R. C., & Coelho, L. C. (2015). Water-soluble *Moringa oleifera* lectin interferes with growth, survival and cell permeability of corrosive and pathogenic bacteria. *Journal of applied microbiology*, 119, 666–676.
- Moura, M. C., Pontual, E. V., Gomes, F. S., Napoleao, T. H., Xavier, H. S., Paiva, P. M. G., & Coelho, L. C. B. B. (2011). Preparations of *Moringa oleifera* flowers to treat contaminated water. In J. A. Daniels (Ed.), *Advances in Environmental Research* (pp. 269–275). Inc.: Nova Science Publishers.
- Moyo, B., Masika, P., Hugo, A., & Muchenje, V. (2011). Nutritional characterization of *Moringa (Moringa oleifera)* Lam. leaves. *African Journal of Biotechnology*, 10, 12925–12933.
- Muangnoi, C., Chingsuwanrote, P., Praengamthanachoti, P., Svasti, S., & Tuntipipat, S. (2012). *Moringa oleifera* pod inhibits inflammatory mediator production by lipopolysaccharide-stimulated RAW 264.7 murine macrophage cell lines. *Inflammation*, 35, 445–455.
- Mughal, M. H., Ali, G., Srivastava, P. S., & Iqbal, M. (1999). Improvement of drumstick (*Moringa pterygosperma* Gaertn.)—A unique source of food and medicine through tissue culture. *Hamdard Medicus*, 42, 37–42.
- Muhammad, A. A., Arulselvan, P., Cheah, P. S., Abas, F., & Fakurazi, S. (2016). Evaluation of wound healing properties of bioactive aqueous fraction from *Moringa oleifera* Lam. on experimentally induced diabetic animal model. *Drug Design. Development and Therapy*, 10, 1715–1730.
- Muhammad, A. A., Pauzi, N. A. S., Arulselvan, P., Abas, F., & Fakurazi, S. (2013). *In vitro* wound healing potential and identification of bioactive compounds from *Moringa oleifera* Lam. *BioMed Research International*, 1D, 974580, 1–10.
- Mukunzi, D., Nsor-Atindana, J., Xiaoming, Z., Gahungu, A., Karangwa, E., & Mukamurezi, G. (2011). Comparison of volatile profile of *Moringa oleifera* leaves from Rwanda and China using HS-SPME. *Pakistan Journal of Nutrition*, 10, 602–608.
- Mulyaningsih, T. R., & Yusuf, S. (2018). Determination of minerals content in leaves of *Moringa oleifera* by neutron activation analysis. *Ganendra Journal of Nuclear Science and Technology*, 21(1), 11–16.
- Munim A., Puteri M. U., Sari S. P., & Azizahwati (2016). Anti-anemia effect of standardized extract of *Moringa oleifera* Lam. leaves on aniline induced rats. *Pharmacognosy Journal*, 8(3), 255–258.
- Murakami, A., Kitazono, Y., Jiwajinda, S., Koshimizu, K., & Ohigashi, H. (1998). Niaziminin, a thiocarbamate from the leaves of *Moringa oleifera*, holds a strict structural requirement for inhibition of tumor-promoter-induced Epstein-Barr virus activation. *Planta Medica*, 64, 319–323.
- Nadeem, A. M., Mahmood, S. A., Shahid, S. A., Shah, S. S., Khalid, A. M., & McKay, G. (2006). Sorption of lead from aqueous solution by chemically modified carbon adsorbents. *Journal of Hazardous Material*, 138, 604–613.
- Nadkarni, A. K. (1976). *Indian Materia Medica* (pp. 810–816). Bombay, India: Popular Prakashan.
- Nair, S., & Varalakshmi, K. N. (2011). Anticancer, cytotoxic potential of *Moringa oleifera* extracts on HeLa cell line. *Journal of Natural Pharmaceuticals*, 2, 138–142.
- Nambiar, V. S., Guin, P., Parnami, S., & Daniel, M. (2010). Impact of antioxidants from drumstick leaves on the lipid profile of hyperlipidemics. *Journal of Herbal Medicine and Toxicology*, 4, 165–172.
- Nandave, M., Ojha, S. K., Joshi, S., Kumari, S., & Arya, D. S. (2009). *Moringa oleifera* leaf extract prevents isoproterenol-induced myocardial damage in rats: evidence for an antioxidant, antiperoxidative, and cardioprotective intervention. *Journal of Medicinal Food*, 12, 47–55.
- Nautiyal, B. P., & Venhataraman, K. G. (1987). *Moringa* (Drumstick) - An ideal tree for social forestry. *Myforest*, 23(1), 53–58.
- Ndhkala, A., Mulauudzi, R., Ncube, B., Abdelgadir, H. A., du Plooy, C. P., & Van-Staden, J. (2014). Antioxidant, antimicrobial and phytochemical variations in thirteen *Moringa oleifera* Lam. Cultivars. *Molecules*, 19(7), 10480–10494.
- Nfambi, J., Bbosa, G. S., Sembajwe, L. F., Gakunga, J., & Kasolo, J. N. (2015). Immunomodulatory activity of methanolic leaf extract of *Moringa oleifera* in Wistar albino rats. *Journal of Basic and Clinical Physiology and Pharmacology*, 26, 603–611.
- Ngamukote, S., Khannongpho, T., Siriwatanapaiboon, M., Sirikwanpong, S., Dahlan, W., & Adisakwattana, S. (2016). *Moringa oleifera* leaf extract increases plasma antioxidant status associated with reduced plasma malondialdehyde concentration without hypoglycemia in fasting healthy volunteers. *Chinese Journal of Integrative Medicine*, 22, 1–6.
- Nibret, E., & Wink, M. (2010). Trypanocidal and antileukaemic effects of the essential oils of *Hagenia abyssinica*, *Leonotisocymifolia*, *Moringa stenopetala*, and their main individual constituents. *Phytomedicine*, 17, 911–920.
- Nikken, F., Saud, Z. A., Rahman, M. H., & Haque, M. E. (2003). *In vitro* antimicrobial activity of the compound isolated from chloroform extract of *Moringa oleifera* Lam. *Pakistan Journal of Biological Science*, 6, 1888–1890.
- Noolkar, S. B., Jadhav, N. R., Bhende, S. A., & Killedar, S. G. (2013). Solid-state characterization and dissolution properties of Meloxicam-Moringa Coagulant-PVP ternary solid dispersions. *AAPS PharmSciTech*, 14, 569–577.
- Noonan, S., & Savage, G. (1999). Oxalate content of food and its effect on humans. *Asia Pacific Journal of Clinical Nutrition*, 8, 64–74.
- Nouman, W., Anwar, F., Gull, T., Amaglo-Newton, A., Rosa, E., & Dominguez-Perles, R. (2016). Profiling of polyphenolics, nutrients and antioxidant potential of germplasm's leaves from seven cultivars of *Moringa oleifera* Lam. *Industrial Crops and Products*, 83, 166–176.
- Nouman, W., Basra, S. M. A., Siddiqui, M. T., Yasmeen, A., Gull, T., & Alcaide, M. A. C. (2013). Potential of *Moringa oleifera* Lam. as livestock fodder crop: A review. *Turkish Journal of Agriculture and Forestry*, 37, 1211–1266.

- Nunthanawanich, P., Sompong, W., Sirikwanpong, S., Makynen, K., Adisakwattana, S., Dahlan, W., & Ngamukote, S. (2016). *Moringa oleifera* aqueous leaf extract inhibits reducing monosaccharide-induced protein glycation and oxidation of bovine serum albumin. *SpringerPlus*, 5, 1098.
- Nworu, C. S., Okoye, E. L., Ezeifeke, G. O., & Esimone, C. O. (2013). Extracts of *Moringa oleifera* Lam. showing inhibitory activity against early steps in the infectivity of HIV-1 lentiviral particles in a viral vector-based screening. *African Journal of Biotechnology*, 12, 4866–4873.
- Nwosu, M. O., & Okafor, J. L. (1995). Preliminary studies of antifungal activities of some medicinal plants against *Basidiobolus* and some other pathogenic fungi. *Mycoses*, 38, 191–195.
- Nyein, M. M., Aye, T., Khine, W. W., Wai, K. T., Tun, S., Htwe, S. S., ... Swe, T. (1997). The use of *Moringa oleifera* (dan-da-lun) seed for the sedimentation and decontamination of household water. Part II: community-based study. *Myanmar Health Sciences Research Journal*, 9(3), 163–166.
- Oduro, I., Ellis, W. O., & Owusu, D. (2008). Nutritional potential of two leafy vegetables: *Moringa oleifera* and *Ipomoea batatas* leaves. *Scientific Research and Essays*, 3, 5–60.
- Ogbe, A. O., & Affiku, J. P. (2011). Proximate study, mineral and anti-nutrient composition of *Moringa oleifera* leaves harvested from Lafia, Nigeria: Potential benefits in poultry nutrition and health. *Journal of Microbiology, Biotechnology and Food Sciences*, 1(3), 296–308.
- Ogunbino, A. O., Flamini, G., Cioni, P. L., Adebayo, M. A., & Oguwande, I. A. (2009). Constituents of *Cajanus cajan* (L.) Millsp., *Moringa oleifera* Lam., *Heliotropium indicum* L. and *Bidens pilosa* L. from Nigeria. *Natural Product Communications*, 4, 573–578.
- Okuda, T., Baes, A. U., Nishijima, W., & Okada, M. (2001). Isolation and characterization of coagulant extracted from *Moringa oleifera* seed by salt solution. *Water Research*, 35, 40–410.
- Okwari, O., Dasofunjo, K., Asuk, A., Alagwu, E., & Mokwe, C. (2013). Anti-hypercholesterolemic and hepatoprotective effect of aqueous leaf extract of *Moringa oleifera* in rats fed with thermoxidized palm oil diet. *Journal of Pharmaceutical and Biological Sciences*, 8, 57–62.
- Olagbemide, P. T., & Alikwe, P. C. (2014). Proximate analysis and chemical composition of raw and defatted *Moringa oleifera* kernel. *Life Sciences Advanced Technologies*, 24, 92–99.
- Olsen, A. (1987). Low technology water purification by bentonite clay and *Moringa oleifera* seed flocculation as performed in Sudanese villages: Effects on *Schistosoma mansoni* cercariae. *Water Research*, 21, 517–522.
- Olsen M.E. (2010). Moringaceae Martinor; Drumstick Tree family; In: *Flora of North America, North of Mexico, Vol 7: Magnoliophyta: Dilleniidae, Part 2*. Oxford University Press, UK; 168.
- Oluduro, A. O. (2012). Evaluation of antimicrobial properties and nutritional potentials of *Moringa oleifera* Lam. leaf in South-Western Nigeria. *Malaysian Journal of Microbiology*, 8, 59–67.
- Oluduro, O. A., Aderiyi, B. I., Connolly, J. D., Akintayo, E. T., & Famurewa, O. (2010). Characterization and antimicrobial activity of 4-( $\beta$ -D-glucopyranosyl-1  $\rightarrow$  4- $\alpha$ -L-rhamnopyranosyloxy)-benzyl thiocarboxamide; a novel bioactive compound from *Moringa oleifera* seed extract. *Folia microbiologica*, 55, 422–426.
- Omodanisi, E. I., Aboua, Y. G., & Oguntibeju, O. O. (2017). Assessment of the anti-hyperglycaemic, anti-inflammatory and antioxidant activities of the methanol extract of *Moringa oleifera* in diabetes-induced nephrotoxic male wistar rats. *Molecules*, 22(4), 1–16.
- Onsare, J. G., & Arora, D. S. (2015). Antibiofilm potential of flavonoids extracted from *Moringa oleifera* seed coat against *Staphylococcus aureus*, *Pseudomonas aeruginosa* and *Candida albicans*. *Journal of Applied Microbiology*, 118, 313–325.
- Owusu, D., & Oduro, I. (2011). Development of crackers from cassava and sweet-potato flours using *Moringa oleifera* and *Ipomoea batatas* leaves as fortificant. *American Journal of Food Nutrition*, 1, 114–122.
- Oyeyinka, A. T., & Oyeyinka, S. A. (2018). *Moringa oleifera* as a food fortificant: recent trends and prospects. *Journal of Saudi Society of Agriculture Science*, 17, 127–136.
- Padla, E. P., Solis, L. T., Levida, R. M., Shen, C. C., & Ragasa, C. Y. (2012). Antimicrobial isothiocyanates from the seeds of *Moringa oleifera* Lam. *Z Naturforsch C*, 67, 557–564.
- Pamok, S., Vinitketkumnuen, S. S. U., & Saenphet, K. (2012). Antiproliferative effect of *Moringa oleifera* Lam. and *Pseuderanthemum palatiferum* (Nees) Radlk extracts on the colon cancer cells. *Journal of Medicinal Plants Research*, 6, 139–145.
- Pan, S. Y., Zhou, S. F., Gao, S. H., Yu, Z. L., Zhang, S. F., Tang, M. K., ... Ko, K. M. (2013). New perspectives on how to discover drugs from herbal medicines: CAM's outstanding contribution to modern therapeutics, new perspectives on how to discover drugs from herbal medicines: CAM's outstanding contribution to modern therapeutics. *Evidence-Based Complementary and Alternative Medicine*, 2013, 627375, 1–25.
- Panda, S. (2015). Butanolic fraction of *Moringa oleifera* Lam. (Moringaceae) attenuates isoproterenol-induced cardiac necrosis and oxidative stress in rats: an EPR study. *EXCLI Journal*, 14, 64–74.
- Panda, S., Kar, A., Sharma, P., & Sharma, A. (2013). Cardioprotective potential of N,  $\alpha$ -L-rhamnopyranosylvincosamide, an indole alkaloid, isolated from the leaves of *Moringa oleifera* in isoproterenol induced cardiotoxic rats: In vivo and in vitro studies. *Bioorganic and Medicinal Chemistry Letters*, 23, 959–962.
- Pandey, A., Pandey, R. D., Tripathi, P., Gupta, P. P., Haider, J., Bhatt, S., & Singh, A. V. (2012). *Moringa oleifera* Lam. (Sahijan) - a plant with a plethora of diverse therapeutic benefits: an updated retrospection. *Medicinal and Aromatic Plants*, 1, 101–108.
- Pari, L., & Kumar, N. A. (2002). Hepatoprotective activity of *Moringa oleifera* on antitubercular drug-induced liver damage in rats. *Journal of Medicinal Food*, 5, 171–177.
- Park, B., Song, Y., Yee, S. B., Lee, B., Seo, S., Park, Y., ... Yoo, Y. H. (2005). Phosphorus 15-p53 translocates into mitochondria and interacts with Bcl-2 and Bcl-xL in eugenol-induced apoptosis. *Apoptosis*, 10(1), 193–200.
- Parvathy, M. V. S., & Umamaheshwari, A. (2007). Cytotoxic effect of *Moringa oleifera* leaf extracts on human multiple myeloma cell lines. *Trends in Medical Research*, 2, 44–50.
- Parwani, L., Bhatnagar, M., Bhatnagar, A., Sharma, V., & Sharma, V. (2016). Evaluation of *Moringa oleifera* seed biopolymer-PVA composite hydrogel in wound healing dressing. *Iranian Polymer Journal*, 25(11), 919–931.
- Pasha S.N., Shafi K.M., Joshi A.G., Meenakshi I., Harini K., Mahita J., Sajeevan R.S., Karpe S.D., Ghosh P., Nitish S., Gandhimathi A., Mathew O.K., Prasanna S.H., Malini M., Mutt E., Naika M., Ravooru N., Rao R. M., Shingate P.N., Sukhwal A., Sunitha M.S., Upadhyay A.K., Vinekar R. S., & Sowdhamini R. (2019). The transcriptome enables the identification of candidate genes behind medicinal value of Drumstick tree (*Moringa oleifera*). *Genomics*, <https://doi.org/10.1016/j.ygeno.2019.04.014> (Published online on dated April 29, 2019)
- Patel, J. P., Gami, B., & Patel, K. (2010). Evaluation of in vitro schizonticidal properties of acetone extract of some Indian medicinal plants. *Advances in Biological Research*, 4, 253–258.
- Patel, M. T., Patel, J. K., & Upadhyay, U. M. (2012). Assessment of various pharmaceutical excipient properties of natural *Moringa oleifera* gum

- [Mucoadhesion, disintegration, binder]. *International Journal of Pharmacy and Life Sciences*, 3, 1833–1847.
- Patel, N., Patel, P., Patel, D., Desai, S., & Meshram, D. (2014). Phytochemical analysis and antibacterial activity of *Moringa oleifera*. *International Journal of Medicine and Pharmaceutical Sciences*, 4, 27–34.
- Paula, P. C., Sousa, D. O., Oliveira, J. T., Carvalho, A. F., Alves, B. G., Pereira, M. L., ... Vasconcelos, I. M. (2017). A protein isolate from *Moringa oleifera* leaves has hypoglycemic and antioxidant effects in alloxan-induced diabetic mice. *Molecules*, 22, 1–15.
- Peixoto, J. R., Silva, G. C., Costa, R. A., Vieira, G. H., Fonteles Filho, A. A., & dos Fernandes-Vieira, R. H. (2011). In vitro antibacterial effect of aqueous and ethanolic *Moringa oleifera* leaf extracts. *Asian Pacific Journal of Tropical Medicine*, 4, 201–204.
- Pinto, C. E., Farias, D. F., Carvalho, A. F., Oliveira, J. T., Pereira, M. L., Grangeiro, T. B., ... Vasconcelos, I. M. (2015). Food safety assessment of an antifungal protein from *Moringa oleifera* seeds in an agricultural biotechnology perspective. *Food and Chemical Toxicology*, 83, 1–9.
- Pisano, M., Pagnan, G., Loi, M., Mura, M. E., Tilocca, M. G., Palmieri, G., ... Rozzo, C. (2007). Antiproliferative and pro-apoptotic activity of eugenol-related biphenyls on malignant melanoma cells. *Molecular Cancer*, 6(8), 1–12.
- Pontual, E. V., Carvalho, B. E., Bezerra, R. S., Coelho, L. C. B. B., Napoleao, T. H., & Paiva, P. M. G. (2012). Caseinolytic and milk-clotting activities from *Moringa oleifera* flowers. *Food Chemistry*, 135, 1848–1854.
- Pontual, E. V., de Lima-Santos, N. D., de Moura, M. C., Coelho, L. C., Navarro, D. M. A. F., Napoleao, T. H., & Paiva, P. M. (2014). Trypsin inhibitor from *Moringa oleifera* flowers interferes with survival and development of *Aedes aegypti* larvae and kills bacteria inhabitant of larvae midgut. *Parasitology Research*, 113, 727–733.
- Popoola, J. O., & Obembe, O. O. (2013). Local knowledge, use pattern and geographical distribution of *Moringa oleifera* Lam. (Moringaceae) in Nigeria. *Journal of Ethnopharmacology*, 150, 682–691.
- Prabhu, K., Murugan, K., Nareshkumar, A., Ramasubramanian, N., & Bragadeeswaran, S. (2011). Larvicidal and repellent potential of *Moringa oleifera* against malarial vector, *Anopheles stephensi* Liston (Insecta: Diptera: Culicidae). *Asian Pacific Journal of Tropical Biomedicine*, 1, 124–129.
- Prakash, A. O., Tiwari, R. K., Shukla, S., Mathur, R., & Tiwari, K. K. (1987). Postcoital antifertility effect of some medicinal plants in rats. *Indian Drugs*, 25(2), 40–44.
- Prasad, T. N., & Elumalai, E. K. (2011). Biofabrication of Ag nanoparticles using *Moringa oleifera* leaf extract and their antimicrobial activity. *Asian Pacific Journal of Tropical Biomedicine*, 1, 439–442.
- Pritchard, M., Craven, T., Mkandawire, T., Edmondson, A. S., & O'Neill, J. G. (2010). A comparison between *Moringa oleifera* and chemical coagulants in the purification of drinking water—an alternative sustainable solution for developing countries. *Physics and Chemistry of the Earth*, 35, 798–805.
- Rachmawati, I., & Rifa, M. (2014). In vitro immunomodulatory activity of aqueous extract of *Moringa oleifera* Lam. leaf to the CD4+, CD8+ and B220+ cells in *Mus musculus*. *The Journal of Experimental Life Science*, 4, 15–20.
- Raguindin, P. F. N., Dans, L. F., & King, J. F. (2014). *Moringa oleifera* as a Galactagogue. *Breastfeeding Medicine*, 9(6), 323–234.
- Rahman, M. M., Rahman, M. M., Akhter, S., Jamal, M. A., Pandeya, D. R., Haque, M. A., ... Rahman, A. (2010). Control of coliform bacteria detected from diarrhea associated patients by extracts of *Moringa oleifera*. *Nepal Medical College Journal*, 12, 12–19.
- Raj, A. J., Gopalakrishnan, V. K., Yadav, S. A., & Dorairaj, S. (2011). Antimicrobial activity of *Moringa oleifera* (Lam.) root extract. *Journal of Pharmacy Research*, 4, 1426–1430.
- Rajan, B. K. C. (1986). Agriculture and farm forestry in semi-arid tracts of Karnataka. *Myforest*, 22(1), 41–49.
- Rajendran, K., Balaji, P., & Basu, M. J. (2008). Medicinal plants and their utilization by villagers in southern districts of Tamil Nadu. *Indian Journal of Traditional Knowledge*, 7(3), 417–420.
- Ramabulana, T., Mavunda, R. D., Steenkamp, P. A., Piater, L. A., Dubery, I. A., & Madala, N. E. (2016). Perturbation of pharmacologically relevant polyphenolic compounds in *Moringa oleifera* against photo-oxidative damages imposed by gamma radiation. *Journal of Photochemistry and Photobiology B: Biology*, 156, 79–86.
- Ramachandran, C., Peter, K. V., & Gopalakrishnan, P. K. (1980). Drumstick (*Moringa oleifera*) a multipurpose Indian vegetable. *Economic Botany*, 34(3), 276–283.
- Randriamboavonjy, J. I., Loirand, G., Vaillant, N., Lauzier, B., Derbre, S., Michalet, S., ... Tesse, A. (2016). Cardiac protective effects of *Moringa oleifera* seeds in spontaneous hypertensive rats. *American Journal of Hypertension*, 29, 873–881.
- Randriamboavonjy, J. I., Rio, M., Pacaud, P., Loirand, G., & Tesse, A. (2017). *Moringa oleifera* seeds attenuate vascular oxidative and nitrosative stresses in spontaneously hypertensive rats. *Oxidative Medicine and Cellular Longevity*, 2017(4129459), 1–10.
- Ranjan, R., Swarup, D., Patra, R. C., & Chandra, V. (2009). *Tamarindus indica* L. and *Moringa oleifera* M. extract administration ameliorates fluoride toxicity in rabbits. *Indian Journal of Experimental Biology*, 47, 900–905.
- Rao, A. V., Devi, P. U., & Kamath, R. (2001). In vivo radioprotective effect of *Moringa oleifera* leaves. *Indian Journal of Experimental Biology*, 39, 858–863.
- Rashid, U., Anwar, F., Moser, B. R., & Knothe, G. (2008). *Moringa oleifera* oil: a possible source of biodiesel. *Bioresource Technology*, 99, 8175–8179.
- Rastogi, R. P., & Mehrotra, B. N. (2006). *Compendium of Indian Medicinal plants. Vol II* (p. 468). New Delhi, India: Central Drug Research Institute, Lucknow and National Institute of Science Communication and Information Resources.
- Rathi, B. S., Bodhankar, S. L., & Baheti, A. M. (2006). Evaluation of aqueous leaves extract of *Moringa oleifera* Lam. for wound healing in albino rats. *Indian Journal of Experimental Biology*, 44, 898–901.
- Ratshilivha, N., Awouafack, M. D., du Toit, E. S., & Eloff, J. N. (2014). The variation in antimicrobial and antioxidant activities of acetone leaf extracts of 12 *Moringa oleifera* (Moringaceae) trees enables the selection of trees with additional uses. *South African Journal of Botany*, 92, 59–64.
- Rattanasena, P. (2012). Antioxidant and antibacterial activities of vegetables and fruits commonly consumed in Thailand. *Pakistan Journal of Biological Science*, 15, 877–882.
- Ravikumar, K., & Sheeja, A. K. (2013). Heavy metal removal from water using *Moringa oleifera* seed coagulant and double filtration. *International Journal of Scientific and Engineering Research*, 4, 10–13.
- Ray, K., Hazra, R., & Guha, D. (2003). Central inhibitory effect of *Moringa oleifera* root extract: Possible role of neurotransmitters. *Indian Journal of Experimental Biology*, 41, 1279–1284.
- Razis, A. F. A., Ibrahim, M. D., & Kntayya, S. B. (2014). Health benefits of *Moringa oleifera*. *Asian Pacific Journal of Cancer Prevention*, 15, 8571–8576.
- Reddy, D. H. K., Harinath, Y., Seshiah, K., & Reddy, A. V. R. (2010). Biosorption of Pb (II) from aqueous solutions using chemically modified *Moringa oleifera* tree leaves. *Chemical Engineering Journal*, 162, 626–634.
- Reddy, D. H. K., Seshiah, K., Reddy, A. V. R., & Lee, S. M. (2012). Optimization of Cd (II), Cu (II) and Ni (II) biosorption by chemically modified

- Moringa oleifera* leaves podwer. *Carbohydrate Polymers*, 88, 1077–1086.
- Reddy, D. H. K., Seshaiyah, K., Reddy, A. V. R., Rao, M. M., & Wang, M. C. (2010). Biosorption of Pb<sup>2+</sup> from aqueous solutions by *Moringa oleifera* bark: Equilibrium and kinetic studies. *Journal of Hazardous Material*, 174, 831–838.
- Rocha-Filho, C. A., Albuquerque, L. P., Silva, L. R., Silva, P. C., Coelho, L. C., Navarro, D. M. A. F., ... Paiva, P. M. G. (2015). Assessment of toxicity of *Moringa oleifera* flower extract to *Biomphalaria glabrata*, *Schistosoma mansoni* and *Artemia salina*. *Chemosphere*, 132, 188–192.
- Rockwood, J. L., Anderson, B. G., & Casamatta, D. A. (2013). Potential uses of *Moringa oleifera* and an examination of antibiotic efficacy conferred by *M. oleifera* seed and leaf extracts using crude extraction techniques available to under-served indigenous populations. *International Journal of Phototherapy Research*, 3, 61–71.
- Romanova, D., Vachalkova, A., Cipak, L., Ovesna, Z., & Rauko, P. (2001). Study of antioxidant effect of apigenin, luteolin and quercetin by DNA protective method. *Neoplasma*, 48, 104–107.
- Roy, B. K., Bashar, M. K., Hossain, S. M. J., Huque, K. S., & Makkar, H. P. S. (2016). Performance evaluation of *Moringa oleifera* and available Roughages (Maize and Australian Sweet Jumbo) on feeding values of growing BLRI cattle breed-1 (BCB-1) bulls. *American Journal of Experimental Agriculture*, 14, 1–9.
- Ruckmani, K., Kavimani, S., Anandan, R., & Jaykar B. (1998). Effect of *Moringa oleifera* Lam on paracetamol-induced hepatotoxicity. *Indian Journal of Pharmaceutical Sciences*, 60, 33–35.
- Saadabi, A. M., & Abu-Zaid, I. E. (2011). An in vitro antimicrobial activity of *Moringa oleifera* L. seed extracts against different groups of microorganisms. *Australian Journal of Basic and Applied Sciences*, 5, 129–134.
- Sachan, P., Khan, B. L., Yadav, M. P. S., & Sonkar, S. (2010). Effect of drumstick leaves extraction the nutritional quality of *Moringa* paneer prepared from different blend of cow milk and leaves extract. *Progressive Agriculture*, 10(1), 98–101.
- Safaeian, L., Asghari, G., Javanmard, S. H., & Heidarinejad, A. (2015). The effect of hydroalcoholic extract from the leaves of *Moringa peregrina* (Forssk.) Fiori. on blood pressure and oxidative status in dexamethasone-induced hypertensive rats. *Advanced Biomedical Research*, 4, 1–11.
- Sahakitpichan, P., Mahidol, C., Disadee, W., Ruchirawat, S., & Kanchanapoom, T. (2011). Unusual glycosides of pyrrole alkaloid and 4'-hydroxyphenylethanamide from leaves of *Moringa oleifera*. *Phytochemistry*, 72, 791–795.
- Saini, R. K., Shetty, N. P., & Giridhar, P. (2014). GC-FID/MS analysis of fatty acids in Indian cultivars of *Moringa oleifera*: potential sources of PUFA. *Journal of the American Oil Chemists Society*, 91, 1029–1034.
- Saini, R. K., Sivanesan, I., & Keum, Y. S. (2016). Phytochemicals of *Moringa oleifera*: A review of their nutritional, therapeutic and industrial significance. 3. *Biotech*, 6, 1–14.
- Salama, H. H., Sayed, S. M. E. L., & Abdalla, A. M. (2017). Enhancing the nutritive values of ice milk based on dry leaves and oil of *Moringa oleifera*. *American Journal of Food Technology*, 12(2), 86–95.
- Sallau, A. B., Mada, S. B., Ibrahim, S., & Ibrahim, U. (2012). Effect of boiling, simmering and blanching on the antinutritional content of *Moringa oleifera* leaves. *International Journal of Food Nutrition and Safety*, 2, 1–6.
- Salles, H. O., Braga, A. C., do Nascimento, M. T., Sousa, A. M., Lima, A. R., Vieira-Lda, S., ... Andrade, L. B. (2014). Lectin, hemolysin and protease inhibitors in seed fractions with ovicidal activity against *Haemonchus contortus*. *Revista Brasileira de Parasitologia Veterinaria*, 23, 136–143.
- Sanchez-Machado, D. I., Nunez-Gastelum, J. A., Reyes-Moreno, C., Ramirez-Wong, B., & Lopez-Cervantes, J. (2010). Nutritional quality of edible parts of *Moringa oleifera*. *Food Analytical Methods*, 3, 175–180.
- Sangkitikomol, W., Rocejanasaroj, A., & Tencomnao, T. (2014). Effect of *Moringa oleifera* on advanced glycation end-product formation and lipid metabolism gene expression in HepG2 cells. *Genetics and Molecular Research*, 13, 723–735.
- Santos, A. F. S., Luz, L. A., Argolo, A. C. C., Teixeira, J. A., Paiva, P. M. G., & Coelho, L. C. B. B. (2009). Isolation of a seed coagulant *Moringa oleifera* lectin. *Process Biochemistry*, 44, 504–508.
- Sashidara, K. V., Rosaiah, J. N., Tyagi, E., Shukla, R., Raghurib, R., & Rajendran, S. M. (2009). Rare dipeptide and urea derivatives from roots of *Moringa oleifera* as potential antiinflammatory and antinociceptive agents. *European Journal of Medicinal Chemistry*, 44, 432–436.
- Satish, A., Kumar, R. P., Rakshith, D., Satish, S., & Ahmed, F. (2013). Antimutagenic and antioxidant activity of *Ficus benghalensis* stem bark and *Moringa oleifera* root extract. *International Journal of Chemical and Analytical Science*, 4, 45–48.
- Satish, A., Sairam, S., Ahmed, F., & Urooj, A. (2012). *Moringa oleifera* Lam.: Protease activity against blood coagulation cascade. *Pharmacognosy Research*, 4, 44–49.
- Sayed, A. A. R. (2012). Ferulsinic acid modulates SOD, GSH, and antioxidant enzymes in diabetic kidney. *Evidence-Based Complementary and Alternative Medicine*, 2012(580104), 1–9.
- Scotti, M., Stella, L., Shearer, E. J., & Stover, P. J. (2013). Modeling cellular compartmentation in one-carbon metabolism. *Wiley Interdisciplinary Reviews: Systems Biology and Medicine*, 5, 343–365.
- Sengupta, A., & Gupta, M. P. (1970). Studies on the seed fat composition of Moringaceae family. *European Journal of Lipid Science and Technology*, 72, 6–10.
- Sengupta, M. E., Keraita, B., Olsen, A., Boateng, O. K., Thamsborg, S. M., Palsdottir, G. R., & Dalsgaard, A. (2012). Use of *Moringa oleifera* seed extracts to reduce helminth egg numbers and turbidity in irrigation water. *Water Research*, 46, 3646–3656.
- Seshadri, S., & Nambiar, V. S. (2003). Kanjero (*Digera arvensis*) and Drumstick leaves (*Moringa oleifera*): nutrient profile and potential for human consumption. In A. P. Simopoulos, & C. Gopalan (Eds.), *Plants in Human Health and Nutrition Policy* (pp. 41–59). Basel: Karger Medical and Scientific Publishers.
- Shah, S. K., Jhade, D., & Chouksey, R. (2016). *Moringa oleifera* Lam.: A study of ethnobotany, nutrients and pharmacological profile. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, 7(5), 2158–2165.
- Shanker, K., Gupta, M. M., Srivastava, S. K., Bawankule, D. U., Pal, A., & Khanuja, S. P. S. (2007). Determination of bioactive nitrile glycoside(s) in drumstick (*Moringa oleifera*) by reverse phase HPLC. *Food Chemistry*, 105, 376–382.
- Sharma, P., Kumari, P., Srivastava, M., & Srivastava, S. (2006). Removal of cadmium from aqueous system by shelled *Moringa oleifera* Lam. Seed powder. *Bioresource Technology*, 97, 290–305.
- Sharma, P., Kumari, P., Srivastava, M., & Srivastava, S. (2007). Ternary biosorption studies of Cd (II), Cr (II) and Ni (II) on shelled *Moringa oleifera* seeds. *Bioresource Technology*, 98, 474–477.
- Sharma, V., Paliwal, R., Janmeda, P., & Sharma, S. (2012). Chemopreventive efficacy of *Moringa oleifera* pods against 7, 12-dimethylbenz[a]anthracene induced hepatic carcinogenesis in mice. *Asian Pacific Journal of Cancer Prevention*, 13, 2563–2569.
- Shebek, K., Schantz, A. B., Sines, I., Lauser, K., Velegol, S., & Kumar, M. (2015). The flocculating cationic polypeptide from *Moringa oleifera* seeds



- damages bacterial cell membranes by causing membrane fusion. *Langmuir*, 31, 4496–4502.
- Shih, M. C., Chang, C. M., Kang, S. M., & Tsai, M. L. (2011). Effect of different parts (leaf, stem and stalk) and seasons (summer and winter) on the chemical compositions and antioxidant activity of *Moringa oleifera*. *International Journal of Molecular Science*, 12, 6077–6088.
- Shin, S. H., Park, J. H., Kim, G. C., Park, B. S., Gil, Y. G., & Kim, C. H. (2007). The mechanism of apoptosis induced by eugenol in human osteosarcoma cells. *Journal of the Korean Association of Oral and Maxillofacial Surgeons*, 33(1), 20–27.
- Shukla, S., Mathur, R., & Prakash, A. O. (1988). Antifertility profile of the aqueous extract of *Moringa oleifera* roots. *Journal of Ethnopharmacology*, 22, 51–62.
- Siddhuraju, P., & Becker, K. (2003). Antioxidant properties of various solvent extracts of total phenolic constituents from three different agroclimatic origins of drumstick tree (*Moringa oleifera* Lam.) leaves. *Journal of Agricultural and Food Chemistry*, 51, 2144–2155.
- Singh, B. N., Singh, B. R., Singh, R. L., Prakash, D., Dhakarey, R., Upadhyay, G., & Singh, H. B. (2009). Oxidative DNA damage protective activity, antioxidant and anti-quorum sensing potentials of *Moringa oleifera*. *Food and Chemical Toxicology*, 47, 1109–1116.
- Singh, M. K., Paul, J., De, T., & Chakraborti, T. (2015). Bioactivity guided fractionation of *Moringa oleifera* Lam. flower targeting *Leishmania donovani*. *Indian Journal of Experimental Biology*, 53, 747–752.
- Singh, R. S. G., Negi, P. S., & Radha, C. (2013). Phenolic composition, antioxidant and antimicrobial activities of free and bound phenolic extracts of *Moringa oleifera* seed flour. *Journal of Functional Foods*, 5, 1883–1891.
- Singh, U., Wadhvani, A. M., & Johri, B. M. (1983). *Dictionary of Economic Plants in India* (2nd ed.). New Delhi, India: Indian Council of Agricultural Research.
- Sinha, M., Das, D. K., Bhattacharjee, S., Majumdar, S., & Dey, S. (2011). Leaf extract of *Moringa oleifera* prevents ionizing radiation induced oxidative stress in mice. *Journal of Medicinal Food*, 14, 1167–1172.
- Sohani, S. (2018). A multipurpose tree—*Moringa oleifera*: review on nutritive importance and its medicinal evidence. *International Journal of Engineering Sciences and Management*, 8(1), 1–4.
- Sreelatha, S., Jeyachitra, A., & Padma, P. R. (2011). Antiproliferation and induction of apoptosis by *Moringa oleifera* leaf extract on human cancer cells. *Food and Chemical Toxicology*, 49, 1270–1275.
- Srinivasamurthy, S., Yadav, U., Sahay, S., & Singh, A. (2017). Development of muffin by incorporation of dried *Moringa oleifera* (Drumstick) leaf powder with enhanced micronutrient content. *International Journal of Food Science and Nutrition*, 2(4), 173–178.
- Stadlander, T., & Becker, K. (2017). Proximate composition, amino and fatty acid profiles and element compositions of four different *Moringa* species. *Journal of Agricultural Science*, 9(7), 46–57.
- Stevens, C. G., Ugese, F. D., Otitoju, G. T., & Baiyeri, K. P. (2015). Proximate and anti-nutritional composition of leaves and seeds of *moringa oleifera* in Nigeria: A comparative study. *Journal of Tropical Agriculture, Food, Environment and Extension*, 14(2), 9–17.
- Stevens, G. C., Baiyeri, K. P., & Akininagbe, O. (2013). Ethno-medicinal and culinary uses of *Moringa oleifera* Lam. in Nigeria. *Journal of Medicinal Plants Research*, 7, 799–804.
- Stohs, S. J., & Hartman, M. J. (2015). Review of the safety and efficacy of *Moringa oleifera*. *Phytotherapy Research*, 29, 796–804.
- Sudha, P., Asdaq, S. M., Dhamingi, S. S., & Chandrakala, G. K. (2010). Immunomodulatory activity of methanolic leaf extract of *Moringa oleifera* in animals. *Indian Journal of Physiology and Pharmacology*, 54, 133–140.
- Sui, L., Dong, Y., Watanabe, Y., Yamaguchi, F., Hatano, N., Tsukamoto, I., ... Tokuda, M. (2005). The inhibitory effect and possible mechanisms of D-allose on cancer cell proliferation. *International Journal of Oncology*, 27(4), 907–912.
- Sutar, N. G., Bonde, C. G., Patil, V. V., Narkhede, S. B., Patil, A. P., & Kakade, R. T. (2008). Analgesic activity of seeds of *Moringa oleifera* Lam. *International Journal of Green Pharmacy*, 2(2), 108–110.
- Tahany, M. A. A., Hegazy, A. K., Sayed, A. M., Kabieli, H. F., El-Alfy, T., & El-Komy, S. M. (2010). Study on combined antimicrobial activity of some biologically active constituents from wild *Moringa peregrina* Forssk. *Journal of Yeast Fungal Reserach*, 1, 15–24.
- Tahiliani, P., & Kar, A. (2000). Role of *Moringa oleifera* leaf extract in the regulation of thyroid hormone status in adult male and female rats. *Pharmacology Research*, 41, 319–323.
- Tan, W. S., Arulselvan, P., Karthivashan, G., & F akurazi S. (2015). *Moringa oleifera* flower extract suppresses the activation of inflammatory mediators in lipopolysaccharide-stimulated RAW 264.7 macrophages via NF- $\kappa$ B pathway. *Mediators of Inflammation*, 2015(720171), 1–11.
- Teixeira, E. M. B., Carvalho, M. R. B., Neves, V. A., Silva, M. A., & Arantes-Pereira, L. (2014). Chemical characteristics and fractionation of proteins from *Moringa oleifera* Lam. leaves. *Food Chemistry*, 147, 51–54.
- Tende, J. A., Ezekiel, I., & Dikko, A. A. U. (2011). Effect of ethanolic leaves extract of *Moringa oleifera* on blood glucose levels of streptozocin-induced diabetics and normoglycemic Wistar rats. *British Journal of Pharmacology and Toxicology*, 2, 1–4.
- Thakur, R. S., Soren, G., Pathapati, R. M., & Buchineni, M. (2016). Diuretic activity of *Moringa oleifera* leaves extract in swiss albino rats. *The Pharma Innovation Journal*, 5(3), 8–10.
- Thurber, M. D., & Fahey, J. W. (2009). Adoption of *Moringa oleifera* to combat under-nutrition viewed through the lens of the diffusion of innovations theory. *Ecology of Food and Nutrition*, 48, 212–225.
- Tiloke, C., Phulukdaree, A., & Chuturgoon, A. A. (2013). The antiproliferative effect of *Moringa oleifera* crude aqueous extract on cancerous human alveolar epithelial cells. *BMC Complement Alternative Medicine*, 13, 226–233.
- Torondel, B., Opore, D., Brandberg, B., Cobb, E., & Cairncross, S. (2014). Efficacy of *Moringa oleifera* leaf powder as a hand-washing product: A crossover controlled study among healthy volunteers. *BMC Complement Alternative Medicine*, 14, 1–7.
- Tumer, T. B., Rojas-Silva, P., Poulev, A., Raskin, I., & Waterman, C. (2015). Direct and indirect antioxidant activity of polyphenol- and isothiocyanate-enriched fractions from *Moringa oleifera*. *Journal of Agricultural Food Chemistry*, 63, 1505–1513.
- Tuorkey, M. J. (2016). Effects of *Moringa oleifera* aqueous leaf extract in alloxan induced diabetic mice. *Interventional Medicine and Applied Science*, 8, 109–117.
- Ullah, A., Mariutti, R. B., Masood, R., Caruso, I. P., Costa, G. H., Millena de Freitas, C., ... Arni, R. K. (2015). Crystal structure of mature 2S albumin from *Moringa oleifera* seeds. *Biochemical and Biophysical Research Communications*, 468, 365–371.
- Upadhye, K. P., Rangari, V. D., & Mathur, V. B. (2012). Antimigraine activity study of *Moringa oleifera* leaf juice. *International Journal of Green Pharmacy*, 6, 204–207.
- Valdez-Solana, M. A., Veronica, Y., Tellez-Valencia, A., Garcia-Arenas, G., Salas-Pacheco, J., Alba-Romero, J. J., & Sierra-Campos, E. (2015). Nutritional content and elemental and phytochemical analyses of *Moringa oleifera* grown in Mexico. *Journal of Chemistry*, 2015(860381), 1–9.
- Verma, A. R., Vijayakumar, M., Mathela, C. S., & Rao, C. V. (2009). In vitro and in vivo antioxidant properties of different fractions of *Moringa oleifera* leaves. *Food and Chemical Toxicology*, 47, 2196–2201.

- Verma, K. S., & Nigam, R. (2014). Nutritional Assessment of Different parts of *Moringa oleifera* Lam. collected from Central India. *Journal of Natural Product and Plant Resources*, 4(1), 81–86.
- Verma, S. C., Banerji, R., Misra, G., & Nigam, S. K. (1976). Nutritional value of *Moringa*. *Current Science*, 45(21), 769–770.
- Vieira, G. H. F., Mourao, J. A., Angelo, A. M., Costa, R. A., & Vieira, R. H. S. F. (2010). Antibacterial effect (*in vitro*) of *Moringa oleifera* and *Annona muricata* against gram positive and gram negative bacteria. *Revista do Instituto de Medicina Tropical de Sao Paulo*, 52(3), 129–132.
- Vongsak, B., Gritsanapan, W., Wongkrajang, Y., & Jantan, I. (2013). *In vitro* inhibitory effects of *Moringa oleifera* leaf extract and its major components on chemiluminescence and chemotactic activity of phagocytes. *Natural Product Communications*, 8, 1559–1561.
- Vongsak, B., Mangmool, S., & Gritsanapan, W. (2015). Antioxidant activity and induction of mRNA expressions of antioxidant enzymes in HEK-293 cells of *Moringa oleifera* Leaf extract. *Planta Medica*, 81, 1084–1089.
- Vongsak B., Sithisarn P., & Gritsanapan W. (2014). Simultaneous HPLC quantitative analysis of active compounds in leaves of *Moringa oleifera* Lam. *Journal of Chromatography Science*, 52, 641–645.
- Waiyaput, W., Payungporn, S., Issara-Amphorn, J., Nattanan, T., & Panjaworayan, T. (2012). Inhibitory effects of crude extracts from some edible Thai plants against replication of hepatitis B virus and human liver cancer cells. *BMC Complementary and Alternative Medicine*, 12, 1–7.
- Walter, A., Samuel, W., Peter, A., & Josph, O. (2011). Antibacterial activity of *Moringa oleifera* and *Moringa stenopetala* methanol and n-hexane seed extracts on bacteria implicated in water borne diseases. *African Journal of Microbiological Research*, 5, 153–157.
- Wang, L., Chen, X., & Wu, A. (2016). Mini review on antimicrobial activity and bioactive compounds of *Moringa oleifera*. *Medicinal Chemistry (Los Angeles)*, 6, 578–582.
- Waterman, C., Cheng, D. M., Rojas-Silva, P., Poulev, A., Dreifus, J., Lila, M. A., & Raskin, I. (2014). Stable, water extractable isothiocyanates from *Moringa oleifera* leaves attenuate inflammation *in vitro*. *Phytochemistry*, 103, 114–122.
- Waterman, C., Rojas-Silva, P., Tumer, T. B., Kuhn, P., Richard, A. J., Wicks, S., ... Cefalu, W. (2015). Isothiocyanate-rich *Moringa oleifera* extract reduces weight gain, insulin resistance, and hepatic gluconeogenesis in mice. *Molecular Nutrition and Food Research*, 59, 1013–1024.
- Williams, J., Mai, C. T., Mulinare, J., Isenburg, J., Flood, T. J., Ethen, M., ... Kirby, R. S. (2015). Updated estimates of neural tube defects prevented by mandatory folic acid fortification—United States, 1995–2011. *Morbidity and Mortality Weekly Report*, 64(1), 1–5.
- Yabesh, J. E., Prabhu, S., & Vijayakumar, S. (2014). An ethnobotanical study of medicinal plants used by traditional healers in silent valley of Kerala, India. *Journal of Ethnopharmacology*, 154, 774–789.
- Yang, R. Y., Chang, L. C., Hsu, J. C., Weng, B. B., Palada, M. C., Chadha, M. L., & Levasseur, V. (2006). Nutritional and functional properties of *Moringa* leaves—From germplasm, to plant, to food, to health. In *Proceedings of the Moringa and other Highly Nutritious Plant Resources: Strategies, Standards and Markets for a Better Impact on Nutrition in Africa, Ghana, 16–18 November* (pp. 1–8). Ghana: Accra publications.
- Yassa, N., Masoomi, F., Rankouhi, S. E. R., & Hadjiakhoondi, A. (2009). Chemical composition and antioxidant activity of the extract and essential oil of *Rosa damascena* from Iran, Population of Guilan. *DARU Journal of Pharmaceutical Sciences*, 17, 175–180.
- Younus, I., Siddiq, A., Ishaq, H., Anwer, L., Badar, S., & Ashraf, M. (2016). Evaluation of antiviral activity of plant extracts against foot and mouth disease virus *in vitro*. *Pakistan Journal of Pharmacology Science*, 29, 1263–1268.
- Zaffer, M., Ahmad, S., Sharma, R., Mahajan, S., Gupta, A., & Agnihotri, R. K. (2014). Antibacterial activity of bark extracts of *Moringa oleifera* Lam. Against some selected bacteria. *Pakistan Journal of Pharmacology Science*, 27, 1857–1862.
- Zayas-Viera, M. D. M., Vivas-Mejia, P. E., & Reyes, J. (2016). Anticancer effect of *Moringa oleifera* leaf extract in human cancer cell lines. *Journal of Health Disparities Research and Practice*, 9(1), 141–142.
- Zhao, J. H., Zhang, Y. L., Wang, L. W., Wang, J. Y., & Zhang, C. L. (2012). Bioactive secondary metabolites from *Nigrospora* sp. LLGLM003, an endophytic fungus of the medicinal plant *Moringa oleifera* Lam. *World Journal of Microbiology and Biotechnology*, 28, 2107–2112.
- Zhao, S., & Zhang, D. (2011). Supercritical fluid extraction and characterisation of essential oil from *Moringa oleifera* leaves. In *Chemeca 2011—Engineering a Better World, Sydney, Australia, 18–21, (September 2011)*.
- Zheng, Y., Zhang, Y., & Wu, J. (2016). Yield and quality of *Moringa oleifera* under different planting densities and cutting heights in southwest China. *Industrial Crops and Products*, 91, 88–96.

**How to cite this article:** Dhakad AK, Ikram M, Sharma S, Khan S, Pandey VV, Singh A. Biological, nutritional, and therapeutic significance of *Moringa oleifera* Lam. *Phytotherapy Research*. 2019;33:2870–2903. <https://doi.org/10.1002/ptr.6475>