

## Research

## Open Access

# Plants used traditionally to treat malaria in Brazil: the archives of Flora Medicinal

Alexandros S Botsaris\*

Address: IBPM (Brazilian Institute of Medicinal Plants), Rua Gal Urquiza 128, Rio de Janeiro, RJ, 22431-040, Brazil

Email: Alexandros S Botsaris\* - ibpm@ibpm.org.br

\* Corresponding author

Published: 1 May 2007

Received: 9 January 2006

Accepted: 1 May 2007

*Journal of Ethnobiology and Ethnomedicine* 2007, **3**:18 doi:10.1186/1746-4269-3-18

This article is available from: <http://www.ethnobiomed.com/content/3/1/18>

© 2007 Botsaris; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

## Abstract

The archives of Flora Medicinal, an ancient pharmaceutical laboratory that supported ethnomedical research in Brazil for more than 30 years, were searched for plants with antimalarial use. Forty plant species indicated to treat malaria were described by Dr. J. Monteiro da Silva (Flora Medicinal leader) and his co-workers. Eight species, *Bathysa cuspidata*, *Cosmos sulphureus*, *Cecropia hololeuca*, *Erisma calcaratum*, *Gomphrena arborescens*, *Musa paradisiaca*, *Ocotea odorifera*, and *Pradosia lactescens*, are related as antimalarial for the first time in ethnobotanical studies. Some species, including *Mikania glomerata*, *Melampodium divaricatum*, *Galipea multiflora*, *Aspidosperma polyneuron*, and *Coutarea hexandra*, were reported to have activity in malaria patients under clinical observation. In the information obtained, also, there were many details about the appropriate indication of each plant. For example, some plants are indicated to increase others' potency. There are also plants that are traditionally employed for specific symptoms or conditions that often accompany malaria, such as weakness, renal failure or cerebral malaria. Many plants that have been considered to lack activity against malaria due to absence of in vitro activity against *Plasmodium* can have other mechanisms of action. Thus researchers should observe ethnomedical information before deciding which kind of screening should be used in the search of antimalarial drugs.

## I. Background

Flora Medicinal is an ancient and small pharmaceutical laboratory established, in early 1915, by Mr. José Monteiro da Silva, a Medical Doctor in Rio de Janeiro. Mr. Monteiro da Silva was an idealist who believed that the Brazilian rainforest had an enormous potential for research and discovery of new drugs. For more than 40 years Mr. Monteiro da Silva had organized a group of technicians and scientists who made a great number of excursions into Brazilian rainforest, collecting plant specimens and information. Although he had also edited the *Revista da Flora Medicinal*, a scientific paper in which he described his discoveries, a considerable part of his

research remains unpublished. During the '30 s and '40 s, the *Revista da Flora Medicinal* was translated to French and republished by the Institut Pasteur, in Paris, which allowed some of his findings to be used by the international pharmaceutical industry. During his activities, Mr. Monteiro da Silva and his team described more than 200 new medicinal plants from this region. One of his targets was the study of new antimalarial plants, as at his time malaria was a concerning health problem in Brazil. In the following years, quinine, its derivatives and other drugs helped to control malaria. Nowadays, however, its incidence is again growing worldwide, and *Plasmodium falciparum* is getting more resistant to the usual antimalarial

drugs[1]. It is estimated that 62% of *P. falciparum* around the world presents with mono or multi-resistant drug profile[1]. The World Health Organization estimates that there are between 300 and 500 million new cases of malaria worldwide, every year, mostly in Africa, Asia, South Pacific Islands and South America, which causes, at least, 3 million deaths[2,3]. The main drugs developed for malaria and used up to now (quinine alkaloids derived drugs and artemisinin) were discovered based on traditional use and ethnomedical data[4,5]. New efforts to search for novel drugs for treating malaria are very important in countries like Brazil, where many endemic areas still exist[6]. The study of well-documented data such as the archives of Flora Medicinal can point out traditional and probably effective treatments that had not been yet subjected to testing.

## 2. Materials and methods

All documents, including books, hand notes, unpublished studies and the issues of Revista da Flora Medicinal, belonging to the library of Mr. Monteiro da Silva, were examined for information about botanical therapies and plant species used for malaria. Any data or references to plants used for malaria were carefully inserted into a template, and botanical name and classification were re-examined and confirmed with four major plant databases – The Missouri Botanical Garden's VAST[7], the International Plant Names Index[8], the New York Botanical Garden vascular plants database[9] and the Brazilian's Northeast Plants Database[10]. Other information existing in modern databases such as Pubmed (U.S. National Library of Medicine's database that is searchable on the Web) were also examined and compared to other ethnopharmacological studies and current published data.

A review of plants with possible antimalarial activity reported in ethnomedical studies or in pharmacological and biochemical research was also made [11-31].

## 3. Results

The results are summarized on Table 1. Forty [40] plants with possible antimalarial activity were reported and examined by Dr. Monteiro da Silva and his co-workers. The plants were identified by scientific names and families, as well as by vernacular names and usual translations to English, if existent. For each of the species, the parts used for general conditions and symptoms and for treating malaria, as gathered from ethnomedical reports published in Flora Medicinal, are listed. Scientific data about in vitro and in vivo research are also provided.

Most plants, like *Bidens pilosa*, *Cantharanthus roseus*, *Cassia fistula*, *Cinchona calisaya*, *Cuphea ingrata*, *Geissospermum sericeum*, *Jateorrhiza palmata*, *Quassia amara*, *Simaba ferruginea*, and *Strychnos pseudoquina*, were already reported as

antimalarial in previous ethnobotanical studies. Some of these had also their activity against *Plasmodium* tested, as shown on Table 1. Eight species are reported as antimalarial for the first time: *Bathysa cuspidata*, *Cosmos sulphureus*, *Cecropia hololeuca*, *Erisma calcaratum*, *Gomphrena arborescens*, *Musa paradisiaca*, *Ocotea odorifera*, and *Pradosia lactescens*.

A greater proportion of the plants reported as antimalarial belong to the families Asteraceae (six species), Rubiaceae (five), Apocynaceae (four), and Simaroubaceae (four).

## 4. Discussion

Most research for antimalarial new drugs is only focused on direct activity against *Plasmodium* species. But attention to ethnomedical information gathered by Monteiro da Silva suggests that other effects should be investigated. For example, some plants are referred to enhance the action of other herbs, which can indicate an increase on permeability of the *Plasmodium* membrane to antiparasitic substances, or an inhibition of pump mechanisms of eliminating the drugs[58,59]. Considering that one of the common mechanisms of drugs resistance is the reduction of permeability, the development of drugs that enhance parasite permeability could be of valuable help in the treatment of infectious diseases[60,61]. Other possible mechanism of action is interference with parasite enzymes used for protection against antiparasitic drugs[62].

Some plants with noticeable ethnopharmacological use in malaria showed only weak or even no activity against *Plasmodium* in vitro[55]. For example, *Mikania glomerata*, *Melampodium divaricatum*, *Galipea multiflora*, *Aspidosperma polyneuron*, and *Coutarea hexandra* had their antimalarial activity confirmed by clinical observations of medical doctors (Table 1), an information that yields a high probability of accuracy.

Some authors have underestimated the traditional plants used for malaria based exclusively on low activity against *Plasmodium* in vitro or in animal models[55]. This can be a mistake of strategy or even methodology.

There are many explanations for the absence of in vitro activity of an effective antimalarial drug. As an example, the active principle could be formed by hepatic metabolism, or as a result of transformation by gut bacteria. Other possible mechanisms of action include immunomodulation or interference with the invasion of new red blood cells by parasites, which can be species specific. Therefore studies in human subjects, as well as the observance of ethnomedical detailed data, are urged in order to exclude or confirm the activity of herbs traditionally used to treat malaria.

**Table I: Plants with possible antimalarial activity gathered from ethnomedical reports published in Flora Medicinal**

| Scientific name and family   | Vernacular name  | Part used                       | General indications found in ethnomedical studies  | Information regarding use for malaria  | Scientific data on anti-malarial activity  | Origin and geographic distribution   | Ref.    |
|--|--|---------------------------------|--|--|--|--|---------|
| <i>Aniba canelilla</i> (H.B.K.) Mez. (Lauraceae)   | Preciosa, Casca preciosa, Pau rosa, Casca do Maranhão, Rosewood, Brazilian rosewood  | Barks and leaves                | Arthritis, fever, colic, heart problems, dyspepsia, infection, intermittent fevers, weakness, malaria, leukorrhea, chronic discharge. Thoracic, stimulant.                                     | Dr. Monteiro da Silva indicates that Amazon Indians used this plant to treat malaria.  |  | Large tree that occurs in the Amazon and Atlantic Forest.  | (32)    |
| <i>Acanthospermum australe</i> (Loefl.) Kunt. (Asteraceae)                               | Amor de negro, Mata-pasto, Picão da praia, Picão da prata, Paraguayan starburr   | Leaves and roots                | Fever, malaria, diarrhea, erysipelas, anemia, urinary infections, blennorrhea, bronchitis, dyspepsia. Tonic, diaphoretic, eupetic, antidiarrheal, mucilaginous, antimarial, antiblennorrhagic. | Dr. Monteiro da Silva indicates this plant as a substitute for quina and reports that doctors have a good outcome when using this species in malaria.              | Plants of the same genus used to treat malaria in Africa showed antiplasmodial activity against <i>P. falciparum</i> in vitro. | Herbaceous, invasive and ruderal plant that usually invades crops and occurs spontaneously in the Cerrado. | (33,34) |
| <i>Aristolochia cymbifera</i> Mart. & Zucc. (Aristolochiaceae)                           | Jarrinha, Mil homens   | Roots                           | Asthma, fever, diarrhea, dyspepsia, gout, infection, amenorrhea, orchitis, intermittent fevers.  | Dr. Monteiro da Silva reports that this must be associated to <i>Cayaponia tayuya</i> for use in malaria.  |  |  | (35,36) |
| <i>Aspidosperma polyneuron</i> Müll. Arg. (Apocynaceae)                                  | Peroba rosa, Sobre, Peroba amargosa  | Barks                           | Fever, diarrhea. Febrifuge, antimarial, astringent.  | Indicated for malaria in the Flora Medicinal literature. Plant contains alkaloids with antimarial action. Cases of malaria controlled with this bark are reported. |  | Its alkaloids were extensively studied.  | (37)    |
| <i>Bathysa cuspidata</i> (St. Hil.) Hook. f. (Rubiaceae)                                 | Quina do mato  | Barks                           | Febrifuge, bitter tonic, eupptic used as a substitute for quina in malaria   | Indicated for malaria in the Flora Medicinal literature.   |  | Tree that occurs in the Atlantic Rainforest.   | (38)    |
| <i>Bidens pilosa</i> L. (Asteraceae)   | Picão, Picão preto, Erva picão, Cuambu, Farmer's Friend, Cobbler's pegs, Beggar's ticks, Pitchforks, Hairy beggarticks                         | Leaves                          | Jaundice, fever, hepatitis, leukorrhea, diarrhea, pharyngitis, worms, cough, pneumonia, hepatomegaly. Mucilaginous.  | Indicated in many medical texts in the past for malaria. Dr. Monteiro da Silva used for patients that did not respond to quinine.                                  | Preclinical tests revealed strong antiplasmodial activity.   | Plant with worldwide distribution.   | (33,39) |
| <i>Cosmos sulphureus</i> Cav. Syn. <i>Bidens sulphurea</i> (Cav.) Sch. Bip. (Asteraceae) | Picão de flor grande, Picão grande, Beijo de moça, Cosmo amarelo, Yellow cosmos, Klondike Cosmos, Sulphur cosmos, Orange cosmos                | Fruits and aerial parts         | Jaundice, intermittent fever, splenomegaly. Tonic, hepatic, hepatoprotective.  | Indicated for malaria in the Flora Medicinal literature.   |  | Plant bred with ornamental purposes.   | (33)    |
| <i>Cassia fistula</i> L. (Leguminosae Caesalpinoideae)                                   | Chuva de ouro, Cássia amarela, Cássia imperial, Canafistula, Golden shower, Indian laburnum, Purging fistula, Drumstick tree                   | Barks, leaves and seeds         | Poisons, erysipelas. Febrifuge, purgative, emmenagogue, diuretic, hepatic, skin problems.  | Indicated by Dr. Monteiro da Silva as an adjuvant for the treatment of malaria.  |  | Ornamental plant found all over Brazil.  | (32)    |
| <i>Catharanthus roseus</i> (L.) G. Don. (Apocynaceae)                                    | Vinca, Boa noite, Lavadeira, Vinca rósea, Cape periwinkle, Catharanthus, Church flower, Madagascar periwinkle, Red periwinkle, Rosy periwinkle | Aerial parts                    | Diabetes, urinary infection, malaria, intermittent fever. Sudorific, diuretic, hypoglycemic, febrifuge.  | Indicated as a substitute for quina by Dr. Monteiro da Silva.  |  | Ornamental plant used by the pharmacy industry for obtaining alkaloids.                                    | (33)    |
| <i>Cecropia hololeuca</i> Miq. (Cecropiaceae)  | Emabaúba, Imbaúba, Emabaúba branca, Emabaúba prateada, Trumpet tree, Silver embauva, Black embauva, White embauva                              | Leaves, fruit and sprouts juice | Diuretic, antihypertensive, sedative, refreshing, antiinflammatory, thoracic, healing, expectorant antiasthmatic, cough suppressant, resolute, antithermal.                                    | Indicated by Dr. Monteiro da Silva as an adjuvant in malaria with very high fever or neurological symptoms.  |  | Tree that occurs in the Atlantic Rainforest.   | (40)    |
| <i>Cedrela fissilis</i> Vell. (Meliaceae)  | Cedro rosa, Cedro vermelho, South American cedar   | Barks                           | Swamp fever, urinary infection, diarrhea. Aromatic, astringent, diuretic, depurative, febrifuge.   | Indicated for malaria in the Flora Medicinal literature.   |  | Plant that occurs in the Atlantic Forest.  | (41)    |

**Table I: Plants with possible antimalarial activity gathered from ethnomedical reports published in Flora Medicinal (Continued)**

|  |  |                              |  |  |  |  |
|--|--|------------------------------|--|--|--|--|
| <i>Chondodendron platyphyllum</i> (St. Hill.) Miers.<br>(Menispermaceae) | Abútua, Bútua, Uva do mato   | Roots, barks and leaves      | Gases, colic, diarrhea, abdominal pain, verminosis, fever, emesis, nausea, infection, bronchitis, amenorrhea, intermittent fever. Antiasthmatic, bitter tonic, eupeptic.   | Use by Indians from the Tupi-Guarani tribe for treating malaria, reported by Dr. Monteiro da Silva.  | Isoquinolinic alkaloid-rich plant with antiparasitary activity, natural from the Atlantic Forest.  | (42,43)  |
| <i>Cinchona calisaya</i> Wedd. (Rubiaceae)                               | Quina peruviana, Casca dos jesuítas, Quina verdadeira, Ledger quinine, Calisaya, Jesuit's powder, Yellow cinchona                          | Barks                        | Fever, malaria, eczema. Hair tonic.  | Used as the main source of quinin by Dr. Monteiro da Silva. It has quinolinic and quinin derivatives in its composition.   | Originary from the Amazon.   | (38,44)  |
| <i>Coffea arabica</i> L.<br>(Rubiaceae)                                  | Café, Cafeeiro, Coffee, Arabica coffee, Arabian coffee, Abyssinian coffee, Brazilian coffee  | Leaves and seeds             | Colds, intermittent fever. Clears the blood, diuretic, stimulant, antiasthmatic, digestive, hypoglycemic.  | Dr. Monteiro da Silva used the leaves decoction to potentiate other plants with anti-malarial activity.  | Plant with African origin, adapted to Brazil.  | (45)   |
| <i>Coutarea hexandra</i> (Jacq.) Schum.<br>(Rubiaceae)                   | Quina-quina, Quina-brava, Quina-de-pernambuco, Quineira, Murta do mato   | Barks                        | Intermittent fever, gallbladder stones or problems, digestive problems, colic. Antithermal, antimalarial.  | Dr. Monteiro da Silva relates this plant as one of the substitutes for quina and reports cases of malaria cure with its use, some described in the book <i>Botânica Médica Cearense</i> , from Dr. Francisco Dias da Rocha.      | Plant from the Cerrado used for ornamental purposes.   | (33)   |
| <i>Cuphea ingrata</i> Hoehne<br>(Lythraceae)                             | Sete sangrias, Perna de saracura, Mata cana, Pega pinto  | Aerial parts and whole plant | High blood pressure, syphilis, dermatoses, intermittent fever, stomachache, rheumatism, venereal diseases, urethral discharge. Depurative, antisiphilitic, cholesterol-reducing, antihemorrhagic, mucous membrane protector, tonic, analgesic. | Indicated for malaria in the Flora Medicinal literature. According to Dr. Monteiro da Silva co-workers, this plant potentiates other antimalarial extracts and help preventing renal and cerebral complications in severe cases. | Herbaceous and ruderal plant that occurs in almost all regions of Brazil, used also for ornamental purposes. It is described in all South and Central America.               | Personal writings and archives of Mr. Monteiro da Silva. |
| <i>Dipteryx odorata</i> (Aublet) Willd.<br>(Fabaceae)                    | Fava de Tonka, Faveira de cheiro, Imburana de cheiro, Cumaru de cheiro, Cumaru de folha grande, Tonka bean, Cumaru, Coumarou, Tonquin bean | Seeds                        | Antispasmodic, emmenagogue, analgesic, febrifuge, brain stimulating.   | In a review, Dr. Monteiro da Silva points this species as having potential use in malaria based on ethnopharmacological reports obtained in his expeditions.   | Plant from the Amazon that is rich in coumarins, which gives it a special odor, and for this reason it has been used in the food and tobacco industry as an odorizing agent. | (32)   |
| <i>Elephantopus mollis</i> Kunth. (Asteraceae)                           | Erva grossa, Língua de vaca, Pé de elefante, Elephantopus, Elephant's foot, False tobacco, Tobacco weed                                    | Aerial parts                 | Fever, jaundice, gallstone, diarrhea, herpes, syphilis, colds, flu, rheumatism, general pruritus. Tonic, depurative.   | Indicated for malaria in the Flora Medicinal literature. Some doctors suggest that this plant could be tried if no chinchona bark or substitute is available.  | Herbaceous and ruderal plant that is found all over Latin America.   | (41)   |
| <i>Erisma calcaratum</i> (Link) Warm.<br>(Vochysiaceae)                  | Jaboti, Erva de Jaboti, Jabuti, Jabuti-araconha, Jabuti da várzea, Jaboty, Jaboty palm   | Fruits                       | Skin infections, dermatoses, fever, malaria. Oleaginous, resolute.   | Reported in review as a plant used by the Amazonian Indians for the treatment of malaria.  | Medium sized tree that grows along the moist lowlands of the Amazon.   | (32)   |
| <i>Esenbeckia febrifuga</i> (St. Hil.) A. Juss. ex Mart. (Rutaceae)      | Quina do mato, Angustura, Gumarim  | Barks                        | Malaria, intermittent fever, adenitis, constipation, dyspepsia. Bitter tonic, febrifuge.   | Indicated for malaria in the Flora Medicinal literature.   | Tree natural from the Southern and Southeast regions.  | (38)   |
| <i>Galipea multiflora</i> Schultz (Rutaceae)                             | Quina falsa, Jasmim do mato, Ticoró, Guamixinga  | Barks                        | Dyspepsias, gastric atony, fever, infections, malaria. Tonic, astringent, bitter, eupeptic, febrifuge, antidiarrheal.  | Reported as a substitute for quina in the treatment of malaria in a review by Dr. Monteiro da Silva. Effective in malaria, but weaker than Peruvian chinchona.   | Tree that occurs in the Cerrado and Atlantic Forest.   | (32,38)  |

**Table I: Plants with possible antimalarial activity gathered from ethnomedical reports published in Flora Medicinal (Continued)**

|  |  |                                    |  |  |   |  |            |
|--|--|------------------------------------|--|--|---|--|------------|
| <i>Geissospermum sericeum</i> (Sagot)<br>Benth<br>(Apocynaceae)  | Pau pereira, Quinarana, Pau forquilha, Acaririnha                        | Barks                              | Dermatoses, inflammations, swamp fevers. Bitter tonic.   | Reported in a review as being a plant tested and approved by doctors for malaria.  | Alkaloids with activity against <i>Plasmodium falciparum</i> were isolated from trees of the genus <i>Geissospermum</i> . | Species from the Atlantic Forest.                        | (32,42,46) |
| <i>Gomphrena arborescens</i> L.<br>(Amaranthaceae)   | Paratudo, Paratudinho, Perpétua raiz do padre                            | Leaves, flowers and tuberous roots | Weakness, colitis, fevers, mental fatigue, intermittent fevers. Antithermal, antidiarrheal, febrifuge, tonic, emmenagogue, aromatic, euprotective, antitoxic, protector.     | Indicated for malaria in the Flora Medicinal literature. Use in malaria introduced by Brazilian priests that learned it from Indians.  | Plant natural from the central region (Cerrado) of Brazil, sometimes cultivated as ornamental.                            | (47)   |            |
| <i>Himatanthus lancifolius</i> (Mull. Arg.) Wood, Syn.<br><i>Plumeria lancifolia</i> Müll. Arg.<br>(Apocynaceae) | Agoniada, Plumeria, Agonium, Arapuê                                      | Barks                              | Menstrual cramps, fever, hysteria, gastric atony, malaria. Purgative, antispasmodic.   | Use for malaria described by Pekolt among Guarani Indians.   | Plant that occurs in the Cerrado and Atlantic Forest.   | (48)   |            |
| <i>Jateorhiza palmata</i> Miers.<br>(Menispermaceae)   | Calumba, Calunga   | Barks                              | Flatulence, colic, diarrhea, abdominal pain, verminosis, fever, emesis, nausea, infection, hypertension, bronchitis, dyspepsia, digestive atony. Bitter tonic, euprotective. | Plant rich in quinolinic alkaloids with antiparasitic potential.   | Exotic plant, natural from Africa, adapted to Brazil.   | (41)   |            |
| <i>Melampodium divaricatum</i> (L.C. Rich.) DC.<br>(Asteraceae)  | Picão da praia, Fel da terra, Salsa da praia, Butter daisy               | Leaves                             | Fever, malaria, flatulence, stomachache, colics, joint pain, muscular pain, palpitation, vertigo, rheumatism, jaundice, anuria. Diuretic, carminative.                       | Dr. Monteiro da Silva reports many cases of malaria cure using the extract of this plant.  | Worldwide distribution.   | (32,49)  |            |
| <i>Mikania glomerata</i> Spreng. (Asteraceae)  | Guaco, Coração de Jesus, Erva de cobra, Cipó almecega                    | Leaves and flowers                 | Rheumatism, snake poison, intestinal problems, colics, dysmenorrhea, fever, malaria.   | Dr. Pires de Almeida reports to have observed Indians using this plant for malaria with good outcomes.   | Liana that is common in the Atlantic Forest.  | (50,51)  |            |
| <i>Musa paradisiaca</i> L.<br>(Musaceae)   | Banana, Bananeira  | Stem juice                         | Worms, diarrhea, intermittent fever, weakness. Tonic, antidiarrheal, thoracic, expectorant, nutritive.   | Indicated by Dr. Monteiro da Silva to potentiate other plants used in malaria and help in the recovery of patients.  | Exotic plant adapted to Brazil.   | (52)   |            |
| <i>Ocotea odorifera</i> (Vell.) Rohwer<br>Syn. <i>Ocotea pretiosa</i> (Nees) Mez<br>(Lauraceae)                  | Sassafraz, Canela de sassafraz, Sassafraz do Brasil, Brazilian sassafras | Barks and roots                    | Dermatoses, joint pain, fever, rheumatism, syphilis, gout. Sudorific, depurative.  | Indicated for malaria in the Flora Medicinal literature. One of the plants used by Guarany Indians to treat fever and malaria.   | Species form the Atlantic Forest.   | (41)   |            |
| <i>Picrolemma sprucei</i> Hook. f.<br>(Simaroubaceae)  | Caferana, Caferana verdadeira  | Aerial parts and roots             | Malaria, intermittent fevers. Sudorific, depurative, febrifuge, antiinfectious.  | Dr. Monteiro da Silva reports many cases of recovery from malaria after treatment with the extract of this plant.  | Shrub that grows on solid ground in the Amazon.   | (49)   |            |
| <i>Pradosia lactescens</i> (Vell.) Radlk.<br>(Sapotaceae)  | Bunhanhém, Pau de remo, Pau doce, Guaranhém, Monesia                     | Barks                              | Discharge, bronchitis, hemoptysis, diarrhea, ocular inflammation, tuberculosis, cutaneous ulcers, metrorrhagia. Bark provides a milky juice that is astringent and tonic.    | Indicated for malaria in the Flora Medicinal literature. According to Dr. Monteiro da Silva, this plant could be associated to any antimalarial therapeutic drug if the patient is not recovering quickly. | Species from the Atlantic Forest.   | Personal writings and archives of Mr. Monteiro da Silva. |            |

**Table I: Plants with possible antimalarial activity gathered from ethnomedical reports published in Flora Medicinal (Continued)**

|   |   |                       |  |  |  |   |   |
|---|---|-----------------------|--|--|--|---|---|
| <i>Quassia amara</i> L.<br>(Simaroubaceae)  | Quassia, Casca amargosa,<br>Pau amargo, Pau de surinã,<br>Quassia-wood, Surinam<br>quassia, Bitter quassia,<br>Bitterwood | Barks                 | Gastric debility, dyspepsia,<br>blennorrhea, flatulence, fever,<br>malaria, diarrhea, worms. Bitter<br>tonic.                      | According to a survey by Dr.<br>Monteiro da Silva, it is used<br>by Indians from the North of<br>Brazil and from Suriname for<br>treating malaria.   | Extracts showed<br>antimalarial activity in<br>experimental malaria in<br>mice.  | Plant from the Amazonian<br>Rainforest.   | (53,54)   |
| <i>Remijia ferruginea</i> A.<br>St. Hil. (Rubiaceae)                                  | Quina mineira   | Barks                 | Intermittent fever, malaria.   | Cited by Dr. Monteiro da<br>Silva as one of the species<br>popularly used to substitute<br>quina in the treatment of<br>malaria.   | Tested in mice with<br>experimental malaria<br>caused by <i>P. berghei</i> , with<br>reduction of 98% of<br>infected red blood cells.    | Medium sized tree that<br>occurs in Atlantic Rainforest<br>and South Amazonia.                | (33,55)   |
| <i>Simaba ferruginea</i> A.<br>St. Hil.<br>(Simaroubaceae)                            | Calunga   | Barks and roots       | Malaria, fevers, diarrhea. Tonic,<br>eupeptic, febrifuge,<br>antidiarrheal, diuretic.  | Indicated for malaria in the<br>Flora Medicinal literature.<br>According to Peckolt, it was<br>used by Amazonian Indians<br>to treat malaria.  | A quassinoid isolated<br>from <i>Simaba</i> sp showed<br>activity against<br><i>Plasmodium falciparum</i> in<br>vitro.                   | Huge tree from the Amazon<br>Forest.  | (41,56)   |
| <i>Simarouba amara</i> L.<br>(Simaroubaceae)  | Calunga, Marubá, Marupá,<br>Dysentery bark, Bitterwood,<br>Slave wood, Bitter damson                                      | Barks and roots       | Intestinal infections, verminosis,<br>fever, wounds, infected ulcers,<br>abdominal pain. Antidiarrheal,<br>antispasmodic, healing. | Used by Amazonian Indians<br>to treat fever and malaria.<br>According to Dr. Monteiro<br>da Silva, it can be used in<br>cases with neurological<br>signs.  |  | Big size tree that occurs in<br>the Atlantic Rainforest and<br>South Amazonia.                | Personal<br>writings and<br>archives of Mr.<br>Monteiro da<br>Silva.      |
| <i>Strychnos<br/>pseudoquina</i> A. St.<br>Hil. (Loganiaceae)                         | Quina do campo, Quina<br>branca, Quineira, Quina-<br>grossa, Quina do cerrado   | Barks                 | Splenomegaly, hepatomegaly,<br>intermittent fever, malaria,<br>gastric problems. Tonic, bitter,<br>febrifuge, depurative.          | Cited by Dr. Monteiro da<br>Silva as one of the species<br>popularly used to substitute<br>quina in the treatment of<br>malaria. According to<br>Andrade-Neto, its potency<br>is inferior to Peruvian quina<br>bark and must be associated<br>with other antimalarial<br>plants. | In a test with<br>experimental malaria in<br>chicken caused by <i>P.</i><br><i>berhei</i> , no activity was<br>found.                    | Shrub from the Cerrado<br>that produces edible fruits.  | (33,38,55)  |
| <i>Tabebuia avellanedae</i><br>Lor. Ex Griseb.<br>(Bignoniaceae)                      | Ipê roxo, Pau d'arco,<br>Trumpet tree   | Barks                 | Fever, tumors, allergy,<br>weakness, psoriasis.<br>Antiinfectious, antifungic,<br>anticancer, tonic,<br>immunestimulant.           | Indicated for malaria in the<br>Flora Medicinal literature.<br>Should be added to<br>antimalarial regimens for<br>weak patients or in cases of<br>renal failure.   |  | Tree that occurs in the<br>Cerrado and Atlantic<br>Rainforest, with strong<br>medicinal uses. | Personal<br>writings and<br>archives of Mr.<br>Monteiro da<br>Silva.      |
| <i>Tachia guianensis</i><br>Aubl. (Gentianacea)                                       | Jacaruaru, Quassia do Pará,<br>Caferana, Tinguá-abá   | Branches and<br>roots | Infections, abdominal pain,<br>worms, malaria. Digestive,<br>antiinflamatory, febrifuge.   | One of the plants cited by<br>Dr. Monteiro da Silva as<br>having potential for treating<br>malaria. Plant used in<br>Amazon by Indians to treat<br>malaria.  |  |   | Personal<br>writings and<br>archives of Mr.<br>Monteiro da<br>Silva. (47) |
| <i>Tabebuia<br/>impetiginosa</i> Syn.<br><i>Tabebuia roseo-alba</i><br>(Bignoniaceae) | Ipê preto, Ipê roxo, Ipê rosa,<br>Trumpet tree  | Barks                 | Fever, tumors, malaria,<br>parasitosis. Antiinfectious,<br>antifungic, anticancer, tonic,<br>immunestimulant.                      | Indicated for malaria in the<br>Flora Medicinal literature.<br>Should be added to<br>antimalarial regimens for<br>weak patients or in cases of<br>renal failure.   |  | Tree that occurs in the<br>Cerrado and Atlantic<br>Rainforest, with strong<br>medicinal uses. | Personal<br>writings and<br>archives of Mr.<br>Monteiro da<br>Silva.      |
| <i>Xylopia brasiliensis</i><br>Spreng.<br>(Annonaceae)                                | Embira de caçador, Pindaíba   | Seeds and barks       | Stomachaches, flatulence,<br>malaria. Stomachic, carminative,<br>febrifuge.  | Indicated for malaria in the<br>Flora Medicinal literature,<br>but Dr. Monteiro da Silva<br>considered it a weak<br>antimalarial drug.   | <i>Xylopia</i> sp extracts<br>proved active against <i>P.</i><br><i>falciparum</i> with IC <sub>50</sub><br>between 3 and 10 mcg/<br>ml. | Plant that occurs in the<br>Cerrado and Atlantic<br>Forest.                                   | Personal<br>writings and<br>archives of Mr.<br>Monteiro da<br>Silva. (57) |

## References

1. Khan MA, Smego RA Jr, Razi ST, Beg MA: **Emerging drug-resistance and guidelines for treatment of malaria.** *J Coll Physicians Surg Pak* 2004, **14**:319-324.
2. Wernsdorfer G, Wernsdorfer WH: **Malaria at the turn from the 2nd to the 3rd millennium.** *Wien Klin Wochenschr* 2003, **115**:2-9.
3. Heggenhougen HK, Hackethal V, Vivek P: **The behavioural and social aspects of malaria and its control.** In *Special Programme for Research & Training in Tropical Diseases (TDR)* Geneva: UNDP/World Bank/WHO; 2003:12-52. 79-90
4. Farnsworth NR: **Ethnopharmacology and drug development.** *Ciba Found Symp* 1994, **185**:42-51. discussion 51-59
5. Mueller I, Namugi P, Kundi J, Ivivi R, Tandrapah T, Bjorge S, Reeder JC: **Epidemic malaria in the highlands of Papua New Guinea.** *Am J Trop Med Hyg* 2005, **72**:554-560.
6. Kretli AU, Andrade-Neto VF, Brandao MG, Ferrari WM: **The search for new antimalarial drugs from plants used to treat fever and malaria or plants randomly selected: a review.** *Mem Inst Oswaldo Cruz* 2001, **96**:1033-1042.
7. **The Missouri Botanical Garden's VAST** [<http://mobot.mobot.org/W3T/Search/vast.html>]
8. **International Plant Names Index** [<http://www.ipni.org/index.html>]
9. **New York Botanical Garden vascular plants database** [<http://www.nybg.org/bsci/hcpl/vasc/>]
10. **Brazilian's Northeast Plants Database** [<http://umbuzeiro.cnip.org.br/db/pnech/taxa/>]
11. Addae-Kyereme J, Croft SL, Kendrick H, Wright CW: **Antiplasmodial activities of some Ghanaian plants traditionally used for fever/malaria treatment and of some alkaloids isolated from *Pleiocarpa mutica*; in vivo antimalarial activity of pleiocarpine.** *J Ethnopharmacol* 2001, **76**:99-103.
12. Bertani S, Bourdy G, Landau I, Robinson JC, Esterre P, Deharo E: **Evaluation of French Guiana traditional antimalarial remedies.** *J Ethnopharmacol* 2005, **98**:45-54.
13. Bourdy G, Oporto P, Gimenez A, Deharo E: **A search for natural bioactive compounds in Bolivia through a multidisciplinary approach. Part VI. Evaluation of the antimalarial activity of plants used by Isoceno-Guarani Indians.** *J Ethnopharmacol* 2004, **93**:269-277.
14. Bremen JG, Alilio MS, Mills A: **Conquering the intolerable burden of malaria: what's new, what's needed: a summary.** *Am J Trop Med Hyg* 2004, **71**:1-15.
15. Cabral MM, Azambuja P, Gottlieb OR, Kleffmann T, Garcia ES, Schaub GA: **Burchellin: effects on *Triatoma infestans* and on *Trypanosoma cruzi* within this vector.** *Parasitol Res* 2001, **87**:730-735.
16. Carvalho LH, Brandao MG, Santos-Filho D, Lopes JL, Kretli AU: **Antimalarial activity of crude extracts from Brazilian plants studied in vivo in *Plasmodium berghei*-infected mice and in vitro against *Plasmodium falciparum* in culture.** *Braz J Med Biol Res* 1991, **24**:1113-1123.
17. do Ceu de Madureira M, Paula Martins A, Gomes M, Paiva J, Proenca da Cunha A, do Rosario V: **Antimalarial activity of medicinal plants used in traditional medicine in S. Tome and Principe islands.** *J Ethnopharmacol* 2002, **81**:23-29.
18. Courtois M, Mincheva Z, Andreu F, Rideau M, Viaud-Massuard MC: **Synthesis and biological evaluation with plant cells of new fosfomycin analogues containing a benzoxazolone or oxazolopyridinone ring.** *J Enzyme Inhib Med Chem* 2004, **19**:559-565.
19. Etkin NL: **The co-evolution of people, plants, and parasites: biological and cultural adaptations to malaria.** *Proc Nutr Soc* 2003, **62**:311-317.
20. Hay SI, Guerra CA, Tatem AJ, Noor AM, Snow RW: **The global distribution and population at risk of malaria: past, present, and future.** *Lancet Infect Dis* 2004, **4**:327-336.
21. Lee MR: **Plants against malaria. Part I: Cinchona or the Peruvian bark.** *J R Coll Physicians Edinb* 2002, **32**:189-196.
22. MacKinnon S, Durst T, Arnason JT, Angerhofer C, Pezzuto J, Sanchez-Vindas PE, Poveda LJ, Gbeassor M: **Antimalarial activity of tropical Meliaceae extracts and gedunin derivatives.** *J Nat Prod* 1997, **60**:336-341.
23. Milliken WV: *Plants for Malaria; Plants for Fever* Kew: Royal Botanic Gardens; 1997.
24. Muregi FW, Chhabra SC, Njagi EN, Lang'at-Thoruwa CC, Njue WM, Orago AS, Omar SA, Ndige IO: **In vitro antiplasmodial activity of some plants used in Kisii, Kenya against malaria and their chloroquine potentiation effects.** *J Ethnopharmacol* 2003, **84**:235-239.
25. Randrianarivo-Josua M, Rasidimanana VT, Rabarison H, Cheplogo PK, Ratsimbason M, Mulholland DA, Maucleire P: **Plants traditionally prescribed to treat tazo (malaria) in the eastern region of Madagascar.** *Malar J* 2003, **2**:25.
26. Rasoanaivo P, Petitjean A, Ratsimamanga-Urveng S, Rakoto-Ratsimamanga A: **Medicinal plants used to treat malaria in Madagascar.** *J Ethnopharmacol* 1992, **37**:117-127.
27. Rath K, Taxis K, Walz G, Gleiter CH, Li SM, Heide L: **Pharmacokinetic study of artemisinin after oral intake of a traditional preparation of *Artemisia annua* L. (annual wormwood).** *Am J Trop Med Hyg* 2004, **70**:128-132.
28. Sanon S, Ollivier E, Azas N, Mahiou V, Gasquet M, Ouattara CT, Nebie I, Traore AS, Esposito F, Balansard G, Timon-David P, Fumoux F: **Ethnobotanical survey and in vitro antiplasmodial activity of plants used in traditional medicine in Burkina Faso.** *J Ethnopharmacol* 2003, **86**:143-147.
29. Waller RF, Ralph SA, Reed MB, Su V, Douglas JD, Minnikin DE, Cowman AF, Besra GS, McFadden GI: **A type II pathway for fatty acid biosynthesis presents drug targets in *Plasmodium falciparum*.** *Antimicrob Agents Chemother* 2003, **47**:297-301.
30. Weniger B, Robledo S, Arango GJ, Deharo E, Aragon R, Munoz V, Callapa J, Lobstein A, Anton R: **Antiprotozoal activities of Colombian plants.** *J Ethnopharmacol* 2001, **78**:193-200.
31. Zirhi GN, Mambu L, Guedé-Guina F, Bodo B, Grellier P: **In vitro antiplasmodial activity and cytotoxicity of 33 West African plants used for treatment of malaria.** *J Ethnopharmacol* 2005, **98**:281-285.
32. **Revista da Flora Medicinal, Aug. 1938.** .
33. **Revista da Flora Medicinal, July 1945.** .
34. Sanon S, Azas N, Gasquet M, Ollivier E, Mahiou V, Barro N, Cuzin-Ouattara N, Traore AS, Esposito F, Balansard G, Timon-David P: **Antiplasmodial activity of alkaloid extracts from *Pavetta crassipes* (K. Schum) and *Acanthospermum hispidum* (DC), two plants used in traditional medicine in Burkina Faso.** *Parasitol Res* 2003, **90**:314-317.
35. **Revista da Flora Medicinal, Mar. 1935.** .
36. **Revista da Flora Medicinal, Nov. 1948.** .
37. Barroso SM: *Fitoterapia e Medicina Científica* Rio de Janeiro: Laboratório Flora Medicinal; 1953.
38. Costa DAM: *A Malaria e Suas Diversas Modalidades Clínicas* Rio de Janeiro: Lombaerts & Co; 1885.
39. Andrade-Neto VF, Brandao MG, Oliveira FQ, Casali VW, Njaine B, Zalis MG, Oliveira LA, Kretli AU: **Antimalarial activity of *Bidens pilosa* L. (Asteraceae) ethanol extracts from wild plants collected in various localities or plants cultivated in humus soil.** *Phytother Res* 2004, **18**:634-639.
40. **Revista da Flora Medicinal, May 1935.** .
41. Peckolt G, Peckolt F: *História das Plantas Medicinais e Úteis do Brasil* Rio de Janeiro: Companhia Typographica do Brasil; 1853.
42. **Revista da Flora Medicinal, July 1940.** .
43. **Revista da Flora Medicinal, July 1935.** .
44. **Revista da Flora Medicinal, June 1938.** .
45. **Revista da Flora Medicinal, Nov. 1942.** .
46. Steele JC, Veitch NC, Kite GC, Simmonds MS, Warhurst DC: **Indole and beta-carboline alkaloids from *Geissospermum sericeum*.** *J Nat Prod* 2002, **65**:85-88.
47. Caminhão JM: *Elementos de Botânica Médica* Rio de Janeiro: Typografia Nacional; 1877.
48. **Revista da Flora Medicinal, Aug. 1941.** .
49. **Revista da Flora Medicinal, Nov. 1941.** .
50. **Revista da Flora Medicinal, Mar. 1938.** .
51. **Revista da Flora Medicinal, Mar. 1942.** .
52. **Revista da Flora Medicinal, Fev. 1942.** .
53. **Revista da Flora Medicinal, July-Dec. 1952.** .
54. Ajaiyeoba EO, Abalogu UI, Krebs HC, Oduola AM: **In vivo antimalarial activities of *Quassia amara* and *Quassia undulata* plant extracts in mice.** *J Ethnopharmacol* 1999, **67**:321-325.
55. Andrade-Neto VF, Brandao MG, Stehmann JR, Oliveira LA, Kretli AU: **Antimalarial activity of Cinchona-like plants used to treat fever and malaria in Brazil.** *J Ethnopharmacol* 2003, **87**:253-256.

56. Moretti C, Deharo E, Sauvain M, Jardel C, David PT, Gasquet M: **Antimalarial activity of cedronin.** *J Ethnopharmacol* 1994, **43**:57-61.
57. Jenett-Siems K, Mockenhaupt FP, Bienzle U, Gupta MP, Eich E: **In vitro antiplasmodial activity of Central American medicinal plants.** *Trop Med Int Health* 1999, **4**:611-615.
58. Normark BH, Normark S: **Evolution and spread of antibiotic resistance.** *J Intern Med* 2002, **252**:91-106.
59. Sanchez CP, Stein W, Lanzer M: **Trans stimulation provides evidence for a drug efflux carrier as the mechanism of chloroquine resistance in *Plasmodium falciparum*.** *Biochemistry* 2003, **42**:9383-9394.
60. Jones RN: **The current and future impact of antimicrobial resistance among nosocomial bacterial pathogens.** *Diagn Microbiol Infect Dis* 1992, **15**:3S-10S.
61. Fernandes P, Ferreira BS, Cabral JM: **Solvent tolerance in bacteria: role of efflux pumps and cross-resistance with antibiotics.** *Int J Antimicrob Agents* 2003, **22**:211-216.
62. Trotta RF, Brown ML, Terrell JC, Geyer JA: **Defective DNA repair as a potential mechanism for the rapid development of drug resistance in *Plasmodium falciparum*.** *Biochemistry* 2004, **43**:4885-4891.

Publish with **BioMed Central** and every scientist can read your work free of charge

"*BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime.*"

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

